

On configuration space integrals for links

CHRISTINE LESCOP

CNRS
Institut Fourier
B.P.74
38402 Saint-Martin-d'Hères cedex
FRANCE

Email: lescop@ujf-grenoble.fr

URL: <http://www-fourier.ujf-grenoble.fr/~lescop/>

Abstract

We give an introductory survey on the universal Vassiliev invariant called *the perturbative series expansion of the Chern–Simons theory of links in euclidean space*, and on its relation with the Kontsevich integral. We also prove an original geometric property of the *anomaly* of Bott, Taubes, Altschuler, Freidel and D. Thurston, that allowed Poirier to prove that the Chern–Simons series and the Kontsevich integral coincide up to degree 6.

AMS Classification numbers Primary: 57M27

Secondary: 57M25, 17B37, 81T18

Keywords: Kontsevich Integral, Chern–Simons theory, Vassiliev invariants, links, knots, tangles, configuration spaces, quantum invariants, Jacobi diagrams

1 Introduction

There are essentially two universal Vassiliev invariants of links, the *Kontsevich integral*, and the *perturbative series expansion of the Chern–Simons theory* studied by Guadagnini Martellini and Mintchev [10], Bar-Natan [5], Axelrod and Singer [2, 3], Kontsevich [11], Polyak and Viro [22], Bott and Taubes [8], Altschuler and Freidel [1], D. Thurston [23], Yang, Poirier [20]... The question that was raised by Kontsevich in [11] whether the two invariants coincide or not is still open though it is predicted by the Chern–Simons gauge theory that the two invariants should coincide. In [20], Sylvain Poirier reduced this question to the computation of the *anomaly* of Bott, Taubes, Altschuler, Freidel and D. Thurston, which is an element α of the space of Jacobi diagrams $\mathcal{A}(S^1)$.

In this article, we shall begin with an elementary introduction to the *perturbative series expansion of the Chern–Simons theory of links in euclidean space* defined by means of configuration space integrals in a natural and beautiful way. We shall call this series the Chern–Simons series, and we shall denote it by Z_{CS} . Its physical interpretation will not be treated here and we refer the reader to the survey [14] of Labastida for the interpretation of Z_{CS} in the context of the Chern–Simons gauge theory.

Then we shall give a short survey of the substantial Poirier work [20] that allowed the author [18] to define the isomorphism of \mathcal{A} which transforms the Kontsevich integral into the Poirier limit of the Chern–Simons invariant of framed links, as an explicit function of α , and to prove the algebraic property of the anomaly : *The anomaly has two legs*. We end up the article by proving an additional original geometric property (Proposition 7.2) of the anomaly that allowed Poirier to compute the anomaly up to degree 6.

I thank Stéphane Guillermou, Lucien Guillou, Sylvain Poirier and especially the referee for useful comments on these notes. I also thank Tomotada Ohtsuki and Hitoshi Murakami for organizing the very interesting Kyoto conference of September 2001 at the RIMS, and for inviting me to participate.

2 Introduction to configuration space integrals: The Gauss integrals.

In 1833, Carl Friedrich Gauss defined the first example of a *configuration space integral* for an oriented two-component link. Let us formulate his definition in

a modern language. Consider a smooth (C^∞) embedding

$$L : S_1^1 \sqcup S_2^1 \hookrightarrow \mathbb{R}^3$$

of the disjoint union of two circles $S^1 = \{z \in \mathbb{C} \text{ s.t. } |z| = 1\}$ into \mathbb{R}^3 . With an element (z_1, z_2) of $S_1^1 \times S_2^1$ that will be called a *configuration*, we associate the oriented direction

$$\Psi((z_1, z_2)) = \frac{1}{\| \overrightarrow{L(z_1)L(z_2)} \|} \overrightarrow{L(z_1)L(z_2)} \in S^2$$

of the vector $\overrightarrow{L(z_1)L(z_2)}$. Thus, we have associated a map

$$\Psi : S_1^1 \times S_2^1 \longrightarrow S^2$$

from a compact oriented 2-manifold to another one with our embedding. This map has an integral degree $\deg(\Psi)$ that can be defined in several equivalent ways. For example, it is the *differential degree* $\deg(\Psi, y)$ of any regular value y of Ψ , that is the sum of the ± 1 signs of the Jacobians of Ψ at the points of the preimage of y [19, §5]. Thus, $\deg(\Psi)$ can easily be computed from a regular diagram of our two-component link as the differential degree of a unit vector \vec{v} pointing to the reader or as the differential degree of $(-\vec{v})$.

$$\deg(\Psi) = \deg(\Psi, \vec{v}) = \# \times_{2_1} - \# \times_{1_2} = \deg(\Psi, -\vec{v}) = \# \times_{1_2} - \# \times_{2_1}$$

It can also be defined as the following *configuration space integral*

$$\deg(\Psi) = \int_{S^1 \times S^1} \Psi^*(\omega)$$

where ω is the homogeneous volume form on S^2 such that $\int_{S^2} \omega = 1$. Of course, this integral degree is an isotopy invariant of L , and the reader has recognized that $\deg(\Psi)$ is nothing but the *linking number* of the two components of L .

We can again follow Gauss and associate the following similar *Gauss integral* $I(K; \theta)$ to a C^∞ embedding $K : S^1 \hookrightarrow \mathbb{R}^3$. (The meaning of θ will be specified later.) Here, we consider the *configuration space* $C(K; \theta) = S^1 \times]0, 2\pi[$, and the map

$$\Psi : C(K; \theta) \longrightarrow S^2$$

that maps (z_1, η) to the oriented direction of $\overrightarrow{K(z_1)K(z_1 e^{i\eta})}$, and we set

$$I(K; \theta) = \int_{C(K; \theta)} \Psi^*(\omega).$$

Let us compute $I(K; \theta)$ in some cases. First notice that Ψ may be extended to the closed annulus

$$\overline{C}(K; \theta) = S^1 \times [0, 2\pi]$$

by the tangent map K' of K along $S^1 \times \{0\}$ and by $(-K')$ along $S^1 \times \{2\pi\}$. Then by definition, $I(K; \theta)$ is the algebraic area (the integral of the differential degree with respect to the measure associated with ω) of the image of the annulus in S^2 . Now, assume that K is contained in a horizontal plane except in a neighborhood of crossings where it entirely lies in vertical planes. Such a knot embedding will be called *almost horizontal*. In that case, the image of the annulus boundary has the shape of the following bold line in S^2 .



In particular, for each hemisphere, the differential degree of a regular value of Ψ does not depend on the choice of the regular value in the hemisphere. Assume that the orthogonal projection onto the horizontal plane is regular. Then $I(K; \theta)$ is the average of the differential degrees of the North Pole and the South Pole, and it can be computed from the horizontal projection as

$$I(K; \theta) = \# \nearrow - \# \nwarrow .$$

This number, that is called the *writhe* of the projection, can be changed without changing the isotopy class of the knot by local modifications where --- becomes \frown or \smile . In particular, $I(K; \theta)$ can reach any integral value on a given isotopy class of knots, and since it varies continuously on such a class, it can reach any real value on any given isotopy class of knots. Thus, this Gauss integral is NOT an isotopy invariant.

However, we can follow Guadagnini, Martellini, Mintchev [10] and Bar-Natan [5] and associate configuration space integrals to any embedding L of an oriented one-manifold M and to any Jacobi diagram Γ on M . Let us first recall what a Jacobi diagram on a one-manifold is.


3 Definitions of the spaces of Jacobi diagrams

Definition 3.1 Let M be an oriented one-manifold. A *Jacobi diagram* Γ with support M is a finite uni-trivalent graph Γ without simple loop like $\bullet \rightarrow \bullet$ such that every connected component of Γ has at least one univalent vertex, equipped with:

- (1) an isotopy class of injections i of the set U of univalent vertices of Γ also called *legs* of Γ into the interior of M ,

- (2) an *orientation* of every trivalent vertex, that is a cyclic order on the set of the three half-edges which meet at this vertex.

Such a diagram Γ is represented by a planar immersion of $\Gamma \cup M$ where the univalent vertices of Γ are identified with their images under i , the one-manifold M is represented by solid lines, whereas the diagram Γ is dashed. The vertices are represented by big points. The orientation of a vertex is represented by the counterclockwise order of the three dashed half-edges that meet at that vertex. Here is an example of a diagram Γ on the disjoint union $M = S^1 \sqcup S^1$ of two circles:



The *degree* of such a diagram is half the number of all its vertices.

Let $\mathcal{A}_n^{\mathbb{Q}}(M)$ denote the rational vector space generated by the degree n diagrams on M , quotiented out by the following relations AS and STU:

$$\text{AS: } \begin{array}{c} \text{---} \backslash \\ \text{---} \text{---} \text{---} \\ \text{---} / \end{array} + \begin{array}{c} \text{---} / \\ \text{---} \text{---} \text{---} \\ \text{---} \backslash \end{array} = 0 \quad \text{and} \quad \text{STU: } \begin{array}{c} \text{---} \backslash \\ \text{---} \text{---} \text{---} \\ \text{---} / \end{array} = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array} - \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array}$$

Each of these relations relate diagrams which can be represented by immersions that are identical outside the part of them represented in the pictures. For example, AS identifies the sum of two diagrams which only differ by the orientation at one vertex to zero.

Set $\mathcal{A}_n(M) = \mathcal{A}_n^{\mathbb{R}}(M) = \mathcal{A}_n^{\mathbb{Q}}(M) \otimes_{\mathbb{Q}} \mathbb{R}$ and let

$$\mathcal{A}(M) = \prod_{n \in \mathbb{N}} \mathcal{A}_n(M)$$

denote the product of the $\mathcal{A}_n(M)$ as a topological vector space. $\mathcal{A}_0(M)$ is equal to \mathbb{R} generated by the empty diagram. The degree n part of an element $\alpha = (\alpha_n)_{n \in \mathbb{N}}$ of $\mathcal{A}(M)$ will be denoted by α_n .

4 The Chern–Simons series

Let M be an oriented one-manifold and let

$$L : M \longrightarrow \mathbb{R}^3$$

denote a C^∞ embedding from M to \mathbb{R}^3 . Let Γ be a Jacobi diagram on M . Let $U = U(\Gamma)$ denote the set of univalent vertices of Γ , and let $T = T(\Gamma)$ denote the set of trivalent vertices of Γ . A *configuration* of Γ is an embedding

$$c : U \cup T \hookrightarrow \mathbb{R}^3$$

whose restriction $c|_U$ to U may be written as $L \circ j$ for some injection

$$j : U \hookrightarrow M$$

in the given isotopy class $[i]$ of embeddings of U into the interior of M . Denote the set of these configurations by $C(L; \Gamma)$,

$$C(L; \Gamma) = \{c : U \cup T \hookrightarrow \mathbb{R}^3 ; \exists j \in [i], c|_U = L \circ j\}.$$

In $C(L; \Gamma)$, the univalent vertices move along $L(M)$ while the trivalent vertices move in the ambient space, and $C(L; \Gamma)$ is naturally an open submanifold of $M^U \times (\mathbb{R}^3)^T$.

Denote the set of (dashed) edges of Γ by $E = E(\Gamma)$, and fix an orientation for these edges. Define the map $\Psi : C(L; \Gamma) \longrightarrow (S^2)^E$ whose projection to the S^2 factor indexed by an edge from a vertex v_1 to a vertex v_2 is the direction of $\overrightarrow{c(v_1)c(v_2)}$. This map Ψ is again a map between two orientable manifolds that have the same dimension, namely the number of dashed half-edges of Γ , and we can write the *configuration space integral*:

$$I(L; \Gamma) = \int_{C(L; \Gamma)} \Psi^*(\Lambda^E \omega).$$

Bott and Taubes have proved that this integral is convergent [8]. Thus, this integral is well-defined up to sign. In fact, the orientation of the trivalent vertices of Γ provides $I(L; \Gamma)$ with a well-defined sign. Indeed, since S^2 is equipped with its standard orientation, it is enough to orient $C(L; \Gamma) \subset M^U \times (\mathbb{R}^3)^T$ in order to define this sign. This will be done by providing the set of the natural coordinates of $M^U \times (\mathbb{R}^3)^T$ with some order up to an even permutation. This set is in one-to-one correspondence with the set of (dashed) half-edges of Γ , and the vertex-orientation of the trivalent vertices provides a natural preferred such one-to-one correspondence up to some (even!) cyclic permutations of three half-edges meeting at a trivalent vertex. Fix an order on E , then the set of half-edges becomes ordered by (origin of the first edge, endpoint of the first edge, origin of the second edge, \dots , endpoint of the last edge), and this order orients $C(L; \Gamma)$. The property of this sign is that the product $I(L; \Gamma)[\Gamma] \in \mathcal{A}(M)$ depends neither on our various choices nor on the vertex orientation of Γ .

Now, the *perturbative series expansion of the Chern–Simons theory for one-manifold embeddings in \mathbb{R}^3* is the following sum running over all the Jacobi diagrams Γ without vertex orientation¹:

$$Z_{CS}(L) = \sum_{\Gamma} \frac{I(L; \Gamma)}{\sharp \text{Aut} \Gamma} [\Gamma] \in \mathcal{A}(M)$$

where $\sharp \text{Aut} \Gamma$ is the number of automorphisms of Γ as a uni-trivalent graph with a given isotopy class of injections of U into M , but without vertex-orientation for the trivalent vertices.

Let θ denote the Jacobi diagram

$$\theta = \bigcirc \text{---} \bigcirc$$

on S^1 . When L is a knot K , the degree one part of $Z_{CS}(K)$ is $\frac{I(K; \theta)}{2}[\theta]$ and therefore Z_{CS} is not invariant under isotopy. However, the evaluation Z_{CS}^0 at representatives of knots with null Gauss integral is an isotopy invariant that is a universal Vassiliev invariant of knots. (All the finite type knot invariants in the Vassiliev sense (see [4]) factor through it.) This is the content of the following theorem, due independently to Altschuler and Freidel [1], and to D. Thurston [23], after the work of many people including Guadagnini, Martellini and Mintchev [10], Bar-Natan [5], Axelrod and Singer [2, 3], Kontsevich [11], Bott and Taubes [8]...

Theorem 4.1 (Altschuler–Freidel, D. Thurston, 1995) *If $L = K_1 \cup \dots \cup K_k$ is a link, then $Z_{CS}(L)$ only depends on the isotopy class of L and on the Gauss integrals $I(K_i; \theta)$ of its components. In particular, the evaluation*

$$Z_{CS}^0(L) \in \prod_{n \in \mathbb{N}} \mathcal{A}_n(\sqcup_{i=1}^k S_i^1)$$

at representatives of L whose components have zero Gauss integrals is an isotopy invariant of L . Furthermore, Z_{CS}^0 is a universal Vassiliev invariant of links.

Recall that the normalized Kontsevich integral is also a universal Vassiliev knot invariant that is valued in the same target \mathcal{A} . (See [4, 17].) Thus, the still open natural question raised by Kontsevich in [11] is:

¹This sum runs over equivalence classes of Jacobi diagrams, where two diagrams are equivalent if and only if they coincide except possibly for their vertex orientation.

Is the Kontsevich integral of a zero-framed representative of a knot K equal to $Z_{CS}^0(K)$?

On one hand, this Chern–Simons series has a beautiful, very natural and completely symmetric definition. Furthermore, Dylan Thurston directly² proved that the Chern–Simons series is *rational*³ because it behaves like a series of degrees of maps between closed manifolds [23, 20].

On the other hand, the Kontsevich integral fits in with the framework of quantum link invariants and it can be defined in this setting [15, 13]. Therefore, it is explicitly known how to recover quantum link invariants from the Kontsevich integral [15]. Furthermore, the computation of the Kontsevich integral for links can be reduced to the computation of small link pieces called *elementary q -tangles*. In [20], Sylvain Poirier proved that the same can be done for the Chern–Simons series. Let us begin a review of his results.

5 The Poirier extension of Z_{CS} to tangles

A *planar configuration* is an embedding of a finite set X into the plane \mathbb{R}^2 .

In the *ambient space* $\mathbb{R}^3 = \{(x, y, z)\}$, the *horizontal plane* is the plane ($z = 0$), whereas the *blackboard plane* is the plane ($y = 0$). The z -coordinate of a point $(x, y, z) \in \mathbb{R}^3$ is called its *vertical projection*.

A *tangle* is the intersection of the image of a link representative transverse to $\mathbb{R}^2 \times \{\beta, \tau\}$ with a horizontal slice $\mathbb{R}^2 \times [\beta, \tau]$. In particular, it is an embedded cobordism between two planar configurations.

For $\lambda \in]0, 1]$, let h_λ be the homeomorphism of \mathbb{R}^3 that shrinks the horizontal plane with respect to the formula

$$h_\lambda(x_1, x_2, x_3) = (\lambda x_1, \lambda x_2, x_3).$$

Let $L(M)$ be a tangle. Set

$$Z_P(L(M)) = \lim_{\lambda \rightarrow 0} Z_{CS}(h_\lambda \circ L) \in \mathcal{A}(M).$$

Sylvain Poirier proved that this limit exists [20].

²The usual proof of the rationality of the Kontsevich integral of zero-framed links is indirect. It relies on the existence of a rational associator [15].

³For a link L , the degree n part $Z_{CSn}^0(L)$ of $Z_{CS}^0(L)$ belongs to $\mathcal{A}_n^{\mathbb{Q}}(\sqcup_{i=1}^k S_i^1)$.

Let K be an almost horizontal knot embedding, rotated by 90 degrees around a horizontal axis, (so that it is almost contained in some vertical plane) such that $I(K; \theta) = 0$. Then $I(h_\lambda \circ K; \theta) = 0$, for any $\lambda > 0$, and the Poirier limit Z_P is the limit of a constant map. Therefore, Z_P is equal to the Chern–Simons series we started with for these representatives.

In general, we can see that the limit of $I(h_\lambda \circ K; \theta)$ depends on the differential degree of Ψ near the equator of S^2 . Assume that the height function (the third coordinate) of K is a Morse function, (its second derivative does not vanish when the first one does). Then all the horizontal tangent vectors correspond to extrema of the height function. Identify the horizontal plane to \mathbb{C} so that the unit horizontal vector corresponding to an extremum e is of the form $\exp(i\theta_e)$. When λ approaches 0, all the non-horizontal tangent vectors approach the poles, and the image of our annulus boundary (of Section 2) becomes a family of straight meridians intersecting the equator at directions $\exp(i\theta_e)$, $\exp(i(\theta_e + \pi))$ corresponding to extrema e . These meridians cut our sphere like an orange, and the differential degree becomes constant on the boundaries of orange quarters and makes integral jumps at meridians. Thus, it can be seen that

$$\lim_{\lambda \rightarrow 0} I(h_\lambda \circ K; \theta) \equiv \frac{1}{\pi} \left(\sum_{e \text{ minimum}} \theta_e - \sum_{e \text{ maximum}} \theta_e \right) \bmod \mathbb{Z}$$

In particular, this limit is an integer when the horizontal vectors are in the blackboard plane, and does not vary under isotopies that keep the directions of the horizontal vectors fixed. Therefore, we define a *framed* tangle as a tangle whose horizontal vectors are contained in the blackboard plane. Two tangles are said to be *isotopic* if they can be obtained from one another by a composition of

- a possible *rescaling of the height parameter*, that is a composition by $1_{\mathbb{R}^2} \times h$ where h is an increasing diffeomorphism from $[\beta, \tau]$ to another interval of \mathbb{R} , and
- an isotopy that is framed at any time, and that preserves the bottom and the top planar configurations up to homotheties with positive ratio and translations.

The homotheties with positive ratio will be called *dilatations*.

In [20], Sylvain Poirier proved that his limit Z_P is a well-defined functorial isotopy invariant of framed tangles. He also interpreted his limit for braids (that are paths of planar configurations) as the Chen holonomy of a connection on a trivial bundle over the space of planar configurations. He proved that,

unlike the holonomy of the complex Knizhnik-Zamolodchikov connection that provides a similar definition of the Kontsevich integral for braids [4, 17], the holonomy of his real connection converges without any regularisation for paths reaching limit planar configurations. Thus, he was able to extend his invariant to framed cobordisms between limit⁴ configurations.

The Le and Murakami *framed q -tangles* of [15] are framed cobordisms between particular limit linear configurations. In [15], Le and Murakami extended the Kontsevich integral to a monoidal functor of framed q -tangles that behaves "nicely" under the operations of deleting a component, doubling a component that runs from bottom to top, reversing orientations, and under the orthogonal symmetries whose fix point sets are either the vertical axis, or the blackboard plane or the horizontal plane. The framed q -tangles, these operations and the expected "nice" behaviour are precisely defined in [18]. We call a functor that satisfies all these properties of the Kontsevich–Le–Murakami functor a *good*⁵ functor.

The general properties that Sylvain Poirier proved for his limit Z_P in [20] imply the following theorem.

Theorem 5.1 *The restriction of Z_P to framed q -tangles is a good functor.*

Furthermore, Poirier expressed the variation of Z_P under a *framing change*, where \bigcap becomes \bigcirc , in terms of the following constant called the *anomaly*.

6 The anomaly

Let us define the anomaly. Let $v \in S^2$. Let D_v denote the linear map

$$\begin{array}{ccc} D_v : \mathbb{R} & \longrightarrow & \mathbb{R}^3 \\ 1 & \mapsto & v \end{array}$$

Let Γ be a Jacobi diagram on \mathbb{R} . Define $C(D_v; \Gamma)$ and Ψ as in Section 4. Let $\hat{C}(D_v; \Gamma)$ be the quotient of $C(D_v; \Gamma)$ by the translations parallel to D_v and

⁴These limits are precisely defined in [20, Section 10.1] as the boundary points of a suitable compactification of the space of planar configurations, where the planar configurations are considered up to translations and dilatations. In this compactification, points are allowed to collide with each other. But the compactification provides us with magnifying glasses allowing us to see any restricted configuration at the scales of the collisions.

⁵Again, the complete definition can be found in [18].

by the dilatations. Then Ψ factors through $\hat{C}(D_v; \Gamma)$ that has two dimensions less. Now, allow v to run through S^2 and define $\hat{C}(\Gamma)$ as the total space of the fibration over S^2 where the fiber over v is $\hat{C}(D_v; \Gamma)$. The map Ψ becomes a map between two smooth oriented manifolds of the same dimension. Indeed, $\hat{C}(\Gamma)$ carries a natural smooth structure and can be oriented as follows. Orient $C(D_v; \Gamma)$ as before, orient $\hat{C}(D_v; \Gamma)$ so that $C(D_v; \Gamma)$ is locally homeomorphic to the oriented product (translation vector $(0, 0, z)$ of the oriented line, ratio of homothety $\lambda \in]0, \infty[$) $\times \hat{C}(D_v; \Gamma)$ and orient $\hat{C}(\Gamma)$ with the $(\text{base}(= S^2) \oplus \text{fiber})$ convention⁶. Then we can again define

$$I(\Gamma) = \int_{\hat{C}(\Gamma)} \Psi^*(\Lambda^E \omega).$$

Now, the *anomaly*⁷ is the following sum running over all connected Jacobi diagrams Γ on the oriented lines (again without vertex-orientation):

$$\alpha = \sum \frac{I(\Gamma)}{\#\text{Aut}\Gamma} [\Gamma] \in \mathcal{A}(\mathbb{R}).$$

Its degree one part is

$$\alpha_1 = \left[\begin{array}{c} \uparrow \\ \bullet \\ \downarrow \end{array} \right].$$

The central symmetry of \mathbb{R}^3 acts on $\hat{C}(\Gamma)$ (by composition of the configurations). Studying the different orientation changes induced by this action shows that, for any Γ ,

$$I(\Gamma)[\Gamma] = (-1)^{1+E(\Gamma)+T(\Gamma)} I(\Gamma)[\Gamma] = (-1)^{n+1} I(\Gamma)[\Gamma]$$

Therefore, for any integer n ,

$$\alpha_{2n} = 0.$$

In [15], Le and Murakami proved that any good functor that varies like the Kontsevich integral Z_K under a framing change coincides with Z_K for framed links. This theorem allowed Poirier to derive the following corollary from his work.

Theorem 6.1 (Poirier [20]) *If the anomaly α vanishes in degree greater than one, then the Kontsevich integral of zero framed links is equal to their Chern–Simons series Z_{CS}^0 .*

⁶This can be summarized by saying that the S^2 -coordinates replace (z, λ) .

⁷Our convention differs from the Poirier convention of [20] where α denotes half this element.

Say that an element $\beta = (\beta_n)_{n \in \mathbb{N}}$ in $\mathcal{A}(S^1)$ is a *two-leg element* if, for any $n \in \mathbb{N}$, β_n is a combination of diagrams with two univalent vertices. Let β be a two-leg element. Forgetting S^1 from β gives rise to a unique series β^s of diagrams with two distinguished univalent vertices v_1 and v_2 , such that β^s is symmetric with respect to the exchange of v_1 and v_2 . The series β^s is well-defined thanks to the diagrammatic Bar-Natan version [4] of the Poincaré-Birkhoff-Witt theorem.

A *chord diagram* on a one-manifold M is a Jacobi diagram without trivalent vertices. Its (dashed connected) components are just *chords*. The degree n chord diagrams generate $\mathcal{A}_n(M)$. If Γ is a chord diagram, define $\Phi(\beta)([\Gamma])$ by replacing each chord by β^s . Then $\Phi(\beta)$ is a well-defined morphism of topological vector spaces from $\mathcal{A}(M)$ to $\mathcal{A}(M)$ for any one-manifold M , and $\Phi(\beta)$ is an isomorphism as soon as $\beta_1 \neq 0$. See [18].

In [18], I refined the Le and Murakami uniqueness theorem, by characterizing the possible variations of good functors under framing changes. This allowed me to refine the above Poirier theorem as follows.

Theorem 6.2 ([18]) *The anomaly α is an odd⁸ two-leg element of $\mathcal{A}(S^1)$. For any framed link L , the Poirier limit integral $Z_P(L)$ is equal to $\Phi(\alpha)(Z_K(L))$. Thus, for any zero-framed link L , $Z_{CS}^0(L)$ is equal to $\Phi(\alpha)(Z_K(L))$.*

The above result yields an algebraic constraint on the anomaly. Since, conversely, any functor of the form $\Phi(\alpha)(Z_K(L))$, for a real odd two-leg element α , is a good functor, nothing more can be obtained from algebra. We are now going to try to compute the low degree terms of the anomaly in a geometric way.

7 Some geometric properties of the anomaly

Poirier showed that the anomaly can be defined from the logarithm of the holonomy of his connexion for the two-strand braid $\begin{smallmatrix} \nearrow & \nwarrow \\ \searrow & \swarrow \end{smallmatrix}$. This is restated in the following proposition.

Let Γ be a Jacobi diagram on $\{1, 2\} \times \mathbb{R}$. For $(x, y) \in \mathbb{R}^2$, let

$$\begin{array}{rcl} f_{(x,y)} : \{1, 2\} \times \mathbb{R} & \longrightarrow & \mathbb{R}^3 \\ (1, t) & \mapsto & (0, 0, t) \\ (2, t) & \mapsto & (x, y, -t) \end{array}$$

$${}^8 \alpha_{2n} = 0.$$

For $\theta \in [0, 2\pi[$, let $C_\theta(\Gamma)$ denote the quotient of $C(f_{(\cos(\theta), \sin(\theta))}; \Gamma)$ by the translations by some $(0, 0, z)$, with $z \in \mathbb{R}$. And let $C(\Gamma) = \cup_{\theta \in [0, 2\pi[} C_\theta(\Gamma)$. Orient $C(\Gamma)$ by letting the θ -coordinate $\theta \in [0, 2\pi[$ replace⁹ the translation parameter $z \in \mathbb{R}$.

Define the *two-strand anomaly*

$$\tilde{\alpha} = \sum_{\Gamma \text{ connected Jacobi diagram on } \{1,2\} \times \mathbb{R}} \frac{1}{\sharp \text{Aut}(\Gamma)} \int_{C(\Gamma)} \Psi^*(\Lambda^{E(\Gamma)} \omega)[\Gamma].$$

Let $i : \mathcal{A}(\uparrow\downarrow) \longrightarrow \mathcal{A}(\mathbb{R})$ be the linear continuous map induced by the inclusion from $\{1, 2\} \times \mathbb{R} = \uparrow\downarrow$ to $\bigcap = \hat{\uparrow}\hat{\downarrow}$. The map i sends a Jacobi diagram Γ to the diagram with the same dashed graph equipped with the same orientations at trivalent vertices where the embedding of univalent vertices is composed by the above inclusion.

Proposition 7.1 (Poirier [20])

$$\alpha = -i(\tilde{\alpha}).$$

This definition of the anomaly is easier to handle with. Instead of searching for configurations where the univalent vertices are on a common unknown line, we look for configurations where the univalent vertices have the same horizontal coordinates.

Proposition 7.2 *Let n be an integer greater than 2. Let Γ be a degree n connected Jacobi diagram on $\{1, 2\} \times \mathbb{R}$. Set*

$$I(\Gamma) = \int_{C(\Gamma)} \Psi^*(\Lambda^{E(\Gamma)} \omega).$$

If Γ has less than three vertices on some strand, then $I(\Gamma) = 0$.

In particular, the two-strand anomaly $\tilde{\alpha}_n$ in degree n is a combination of connected diagrams with at least 3 vertices on each strand.

Proof Let Γ be a diagram as in the above statement. Let U_i denote the set of its univalent vertices which are on $\{i\} \times \mathbb{R}$, for $i \in \{1, 2\}$. We shall prove that if U_2 contains less than 3 elements, then $I(\Gamma) = 0$.

⁹This means that if the quotient $C_\theta(\Gamma)$ is oriented so that $C(f_{(\cos(\theta), \sin(\theta))}; \Gamma)$ is oriented by the (fiber \oplus base) convention where the base is $C_\theta(\Gamma)$ and the fiber is the oriented vertical translation factor \mathbb{R} of \mathbb{R}^3 , then $C(\Gamma)$ is oriented with the (base = $[0, 2\pi[\oplus$ fiber = $C_\theta(\Gamma)$) convention.

This will be sufficient to conclude by symmetry. Indeed, if Γ is such that U_1 contains less than 3 elements, then the diagram $s(\Gamma)$ obtained from Γ by exchanging the two lines is such that U_2 contains less than 3 elements; and, by considering the action of the central symmetry of \mathbb{R}^3 , by composition of the configurations (up to translations), we can see that $I(\Gamma) = \pm I(s(\Gamma))$.

If U_2 is empty, then all the $(2\sharp E(\Gamma) - 1)$ -manifolds $C_\theta(\Gamma)$ are the same. Thus $\Psi(C(\Gamma))$ is the image under Ψ of one of them, and its volume in $(S^2)^{E(\Gamma)}$ is zero.

In order to study the two remaining cases where U_2 contains one or two elements, we replace $C(\Gamma)$ by a slightly larger smooth configuration space $\hat{C}(\Gamma)$ in which $C(\Gamma)$ is dense, where Ψ extends so that we shall have

$$I(\Gamma) = \int_{\hat{C}(\Gamma)} \Psi^*(\Lambda^{E(\Gamma)} \omega).$$

Fix one vertex u_0 on $\{1\} \times \mathbb{R}$, and let $C_0(f_{(x,y)}; \Gamma)$ denote the subset of $C(f_{(x,y)}; \Gamma)$ made of the configurations that map u_0 to the origin of \mathbb{R}^3 . Orient it as the quotient of $C(f_{(x,y)}; \Gamma)$ by the vertical translations with the ((fiber $(= \mathbb{R})$) \oplus base) convention. Let $\hat{C}(\Gamma)$ denote the $(2\sharp E(\Gamma))$ -dimensional quotient of

$$P(\Gamma) = \cup_{(x,y) \in \mathbb{R}^2} C_0(f_{(x,y)}; \Gamma)$$

by the dilatations with a ratio $\lambda \in]0, \infty[$. The map Ψ is well-defined on this space which contains the additional configurations corresponding to $(0, 0)$ that constitute a codimension 2 subspace of $\hat{C}(\Gamma)$ that will therefore not contribute to the integral.

The orientation of $\hat{C}(\Gamma)$ is obtained as follows. Orient $P(\Gamma)$ with the convention (base $= \mathbb{R}^2$) \oplus (fiber). Then the orientation of $\hat{C}(\Gamma)$ is defined by the (fiber $=]0, \infty[$) \oplus (base) convention.


Now, let us get rid of the case where $U_2 = \{u\}$, by defining a free smooth action of $]0, \infty[$ on $\hat{C}(\Gamma)$ that does not change the image of a configuration under Ψ , so that Ψ will again factor through a map from a $(2\sharp E - 1)$ -dimensional manifold to $(S^2)^E$ and define a zero integral. Let t denote the other end of the edge of u . Let $\mu \in]0, \infty[$. Let $c \in P(\Gamma)$. Define $\mu.c(x) = x$ if $x \neq u$ and $\mu.c(u) = c(t) + \mu(c(u) - c(t))$. This action is compatible with the action of the dilatations (and different since the degree of Γ is greater than 1), $\Psi(\mu.c) = \Psi(c)$, and we are finished with this case.

Let us study the remaining case, $U_2 = \{u_1, u_2\}$. Denote the trivalent vertex connected by an edge to u_1 by t_1 , and denote the trivalent vertex connected

by an edge to u_2 by t_2 . The vertices t_1 and t_2 may coincide. We shall use the symmetry¹⁰ σ that maps a configuration $c \in P(\Gamma)$ to the configuration $\sigma(c)$ defined by:


$$\begin{aligned}\sigma(c)(x) &= c(x) && \text{if } x \notin U_2 \\ \sigma(c)(u_1) &= c(t_1) + c(t_2) - c(u_2) \\ \sigma(c)(u_2) &= c(t_1) + c(t_2) - c(u_1)\end{aligned}$$

as shown in the picture below.



This symmetry factors through the dilatations and reverses the orientation of $\hat{C}(\Gamma)$. Furthermore, $\Psi(\sigma(c))$ is obtained from $\Psi(c)$ by reversing the two unit vectors corresponding to the edges containing u_1 and u_2 and by exchanging them, that is, by a composition by a diffeomorphism of $(S^2)^{E(\Gamma)}$ that preserves $\Lambda^{E(\Gamma)}\omega$. This proves that $I(\Gamma)$ vanishes and this finishes the proof of the proposition. \square


As a corollary, α_n is a combination of Jacobi diagrams with 6 legs when $n \geq 2$. Unfortunately, unlike the previous two-leg condition, this condition is not very restrictive. Nevertheless, this provides another proof¹¹ of the Poirier and Yang result that $\alpha_3 = 0$. There is no connected degree 3 diagram with at least 6 univalent vertices. Recall that for any integer n , $\alpha_{2n} = 0$. Therefore, the next interesting degree is 5. The connected degree 5 diagrams with at least 6 univalent vertices are necessarily trees and their dashed parts have one of the two forms:



¹⁰This symmetry resembles a Bott and Taubes symmetry that is used to prove the isotopy invariance of Z_{CS}^0 .

¹¹Note that with this two-strand definition all configuration space integrals vanish whereas with the one-strand original definition, the different integrals cancel each other without being zero individually. See [20, Section 7].

Now, observe that if Γ is a Jacobi diagram on $\{1, 2\} \times \mathbb{R}$, with two univalent vertices that are connected to the same trivalent vertex, and that lie on the same

vertical line, like in , then $I(\Gamma)$ vanishes. Indeed the two corresponding edges and the vertical vector are coplanar. Therefore, the image of Ψ must lie inside a codimension one subspace of $(S^2)^{E(\Gamma)}$. This additional remark determines the distribution of the univalent vertices of the above graphs on the two vertical lines. Then Sylvain Poirier [21] computed $\tilde{\alpha}_5$ with the help of Maple, and he found that $\alpha_5 = 0$ thanks to AS and STU. As a corollary, all coefficients of the HOMFLY polynomial properly normalized that are Vassiliev invariants of degree less than seven can be explicitly written as combinations of the configuration space integrals of Section 4.

Thus, the following Bar-Natan theorem generalizes to any *canonical*¹² Vassiliev invariant of degree less than 7.

Theorem 7.3 (Bar-Natan [5], 1990) *Let Δ denote the symmetrized Alexander polynomial. For any knot K ,*

$$\frac{\Delta''(K)(1)}{2} = -\frac{1}{3}I(K; \text{triangle}) + \frac{1}{4}I(K; \text{square}) + \frac{1}{24}.$$

This particular coefficient, that is the degree 2 invariant which can be extracted from the Chern–Simons series, has been further studied in [22].

8 Questions

The first question here is of course:

1. Prove or disprove the physicist conjecture:

$$\alpha_i = 0 \text{ for any } i > 1.$$

Let us briefly summarize what remains to be done here. The space $\mathcal{A}(\mathbb{R}) = \mathcal{A}(S^1)$ splits as a natural direct sum where the space of two-leg diagrams is a direct summand, thanks to the diagrammatic Bar-Natan version [4] of the Poincaré-Birkhoff-Witt theorem. Call the natural projection on this summand

¹²Here, *canonical* can be understood as explicitly recovered from the Kontsevich integral like all the quantum invariants.

the *two-leg projection*. According to Theorem 6.2 and Proposition 7.1, it remains to compute the two-leg projection of $i(\tilde{\alpha}_{2n+1})$ for $n \geq 3$, where Proposition 7.2 and Lemma 1.9 in [20] allow us to neglect numerous diagrams in this computation.

After the articles of Axelrod, Singer [2, 3], Bott and Cattaneo [6, 7, 9], Greg Kuperberg and Dylan Thurston have constructed a universal finite type invariant for homology spheres as a series of configuration space integrals similar to Z_{CS}^0 , in [12]. Their construction yields two natural questions:

2. Find a surgery formula for the Kuperberg–Thurston invariant in terms of the above Chern–Simons series.
3. Compare the Kuperberg–Thurston invariant to the LMO invariant [16].

References

- [1] **D. Altschuler and L. Freidel**, *Vassiliev knot invariants and Chern–Simons perturbation theory to all orders*, Comm. Math. Phys. 187 no. 2 (1997) 261–287
- [2] **S. Axelrod and I. M. Singer**, *Chern–Simons perturbation theory*, Proceedings of the XXth International Conference on Differential Geometric Methods in Theoretical Physics, Vol. 1, 2 (New York, 1991), World Sci. Publishing, River Edge, NJ (1992) 3–45
- [3] **S. Axelrod and I. M. Singer**, *Chern–Simons perturbation theory. II*, J. Differential Geom. 39, no. 1 (1994) 173–213
- [4] **D. Bar-Natan**, *On the Vassiliev knot invariants*, Topology 34 (1995) 423–472
- [5] **D. Bar-Natan**, *Perturbative Chern–Simons Theory*, Journal of Knot Theory and its Ramifications 4-4 (1995) 503–548
- [6] **R. Bott and A. S. Cattaneo**, *Integral invariants of 3-manifolds*. J. Differential Geom. 48, no. 1 (1998) 91–133
- [7] **R. Bott and A. S. Cattaneo**, *Integral invariants of 3-manifolds. II*. J. Differential Geom. 53 no. 1 (1999) 1–13
- [8] **R. Bott and C. Taubes**, *On the self-linking of knots*, Jour. Math. Phys. 35 (10) (1994) 5247–5287
- [9] **A. S. Cattaneo**, *Configuration space integrals and invariants for 3-manifolds and knots*. Low-dimensional topology (Funchal, 1998), Contemp. Math., 233, Amer. Math. Soc., Providence, RI, (1999) 153–165
- [10] **E. Guadagnini, M. Martellini and M. Mintchev**, *Chern–Simons field theory and link invariants*, Nucl.Phys B330 (1990) 575–607
- [11] **M. Kontsevich**, *Vassiliev’s knot invariants*, Adv. in Sov. Math 16(2) (1993) 137–150

- [12] **G. Kuperberg and D. P. Thurston**, *Perturbative 3-manifold invariants by cut-and-paste topology*, preprint, math.GT/9912167
- [13] **C. Kassel**, *Quantum groups*. Graduate Texts in Mathematics, Springer-Verlag, New York (1995)
- [14] **J. M. F. Labastida**, *Chern-Simons gauge theory: ten years after*. Trends in theoretical physics, II (Buenos Aires, 1998), AIP Conf. Proc., 484, Amer. Inst. Phys., Woodbury, NY, (1999) 1–40
- [15] **T. T. Q. Le and J. Murakami**, *The universal Vassiliev-Kontsevich invariant for framed oriented links*, Compositio Mathematica 102 (1996) 41–64
- [16] **T. T. Q. Le, J. Murakami and T. Ohtsuki**, *On a universal perturbative invariant of 3-manifolds*, Topology 37 no. 3 (1998) 539–574
- [17] **C. Lescop**, *Introduction to the Kontsevich Integral of Framed Tangles*, June 1999, Summer School in Grenoble, <http://www-fourier.ujf-grenoble.fr/~lescop/>
- [18] **C. Lescop**, *About the uniqueness and the denominators of the Kontsevich integral*, math.GT/0004094, to appear in Journal of Knot Theory and its Ramifications
- [19] **J. W. Milnor**, *Topology from the differentiable viewpoint*. Based on notes by David W. Weaver. The University Press of Virginia, Charlottesville (1965)
- [20] **S. Poirier**, *The configuration space integral for links and tangles in R^3* , preprint, math.GT/0005085
- [21] **S. Poirier**, *A note on the configuration space integral*, in preparation
- [22] **M. Polyak and O. Viro**, *On the Casson knot invariant*, Knots in Hellas '98, 3 (Delphi). J. Knot Theory Ramifications 10, no. 5 (2001) 711–738
- [23] **D. P. Thurston**, *Integral Expressions for the Vassiliev Knot Invariants*, math.QA/9901110
and more in Dror **Bar-Natan**'s bibliography on Vassiliev invariants at <http://www.ma.huji.ac.il/~drorbn>.