

SIMPLE BIRATIONAL EXTENSIONS OF THE POLYNOMIAL RING $\mathbb{C}^{[3]}$

SHULIM KALIMAN, STÉPHANE VÉNÉREAU, AND MIKHAIL ZAIDENBERG

Prépublication de l'Institut Fourier n°532 (2001)
<http://www-fourier.ujf-grenoble.fr/prepublications.html>

ABSTRACT. The Abhyankar-Sathaye Problem asks whether any biregular embedding $\varphi : \mathbb{C}^k \hookrightarrow \mathbb{C}^n$ can be rectified, that is, whether there exists an automorphism $\alpha \in \text{Aut } \mathbb{C}^n$ such that $\alpha \circ \varphi$ is a linear embedding. Here we study this problem for the embeddings $\varphi : \mathbb{C}^3 \hookrightarrow \mathbb{C}^4$ whose image $X = \varphi(\mathbb{C}^3)$ is given in \mathbb{C}^4 by an equation $p = f(x, y)u + g(x, y, z) = 0$, where $f \in \mathbb{C}[x, y] \setminus \{0\}$ and $g \in \mathbb{C}[x, y, z]$. Under certain additional assumptions we show that, indeed, the polynomial p is a variable of the polynomial ring $\mathbb{C}^{[4]} = \mathbb{C}[x, y, z, u]$ (i.e., a coordinate of a polynomial automorphism of \mathbb{C}^4). This is an analog of a theorem due to Sathaye [30] which concerns the case of embeddings $\mathbb{C}^2 \hookrightarrow \mathbb{C}^3$. Besides, we generalize a theorem of Miyanishi [24, Thm. 2] giving, for a polynomial p as above, a criterion for as when $X = p^{-1}(0) \simeq \mathbb{C}^3$.

CONTENTS

Introduction	2
1. Preliminaries	4
1.1. Affine modifications of UFD's	4
1.2. Acyclic varieties	6
1.3. Digest on Makar-Limanov and Derksen invariants	9
1.4. Variables in polynomial rings	12
2. Simple modifications of acyclic 3-folds along cylindrical divisors	15
2.1. Simple affine modifications	16
2.2. Preserving acyclicity: necessary conditions	16
2.3. Preserving acyclicity: a criterion	27
3. Simple affine modifications of \mathbb{C}^3 diffeomorphic to \mathbb{R}^6	29
3.1. Exotic simple modifications of \mathbb{C}^3	30
3.2. Simple affine modifications of \mathbb{C}^3 isomorphic to \mathbb{C}^3	36
4. Simple birational extensions of $\mathbb{C}^{[3]}$ as variables in $\mathbb{C}^{[4]}$	38
4.1. Partial positive results	38
4.2. Simple modifications of \mathbb{C}^3 rectifiable in \mathbb{C}^5	44
4.3. On Sathaye-Wright's Theorem	46
References	48

Research of the first author partially supported by the NSA grant MDA904-00-1-0016.
The third author is grateful to the IHES and to the MPI at Bonn (where a part of the work was done) for their hospitality and excellent working conditions.

1991 *Mathematics Subject Classification*: 14R10, 14R25.

Key words: affine space, polynomial ring, variable, affine modification, birational extension.

INTRODUCTION

Generalizing a theorem of A. Sathaye [30] it is proven in [15] that if a surface $X = p^{-1}(0) \subseteq \mathbb{C}^3$ with $p = fu + g \in \mathbb{C}^{[3]}$ and $f, g \in \mathbb{C}[x, y]$ is acyclic (that is, $\tilde{H}_*(X; \mathbb{Z}) = 0$) then p is a variable of the polynomial ring $\mathbb{C}^{[3]}$ i.e., a coordinate of an automorphism $\alpha \in \text{Aut } \mathbb{C}^3$. Thus X can be rectified, and so is isomorphic to \mathbb{C}^2 . This does not hold any more in \mathbb{C}^4 (even with $f \in \mathbb{C}[x]$). Indeed [28, 19] the Russell cubic 3-fold

$$X := \{p = x^2u + x + y^2 + z^3 = 0\} \subseteq \mathbb{C}^4$$

is an exotic \mathbb{C}^3 i.e., is diffeomorphic to \mathbb{R}^6 and non-isomorphic to \mathbb{C}^3 . In 2.28, 3.21 and 3.6 below we give a criterion for as when a 3-fold $X = p^{-1}(0) \subset \mathbb{C}^4$ with

$$(1) \quad p = f(x, y)u + g(x, y, z) \in \mathbb{C}^{[4]} \quad (f \in \mathbb{C}^{[2]} \setminus \{0\}, g \in \mathbb{C}^{[3]})$$

is acyclic (resp., is isomorphic to \mathbb{C}^3 resp., is an exotic \mathbb{C}^3). In particular, we show in 2.11 that if X is acyclic then actually it is diffeomorphic to \mathbb{R}^6 . If furthermore (3.21) X is isomorphic to \mathbb{C}^3 then any fiber $X_\lambda := p^{-1}(\lambda)$ ($\lambda \in \mathbb{C}$) of the polynomial p is isomorphic to \mathbb{C}^3 as well, and moreover (with an appropriate choice of coordinates (x, y)) all fibers of the morphism

$$\rho := (x, p) : \mathbb{C}^4 \rightarrow \mathbb{C}^2$$

are reduced and isomorphic to \mathbb{C}^2 . We do not know whether in that case a polynomial p in (1) must be a variable of the polynomial ring $\mathbb{C}^{[4]}$, and ρ must be a trivial family. However, in section 4 in many cases we provide affirmative answers to these questions and give simple concrete examples where the answers remain unknown. Due to the Quillen-Suslin Theorem, the latter question would be answered in positive if the following conjecture [6, (3.8.5)] (cf. [31, 16]) were true for $n = 2 = \dim S$:

Dolgachev-Weisfeiler Conjecture. *Let $f : X \rightarrow S$ be a flat affine morphism of smooth schemes with every fiber isomorphic (over the residue field) to an affine space \mathbb{A}_k^n . Then f is locally trivial in the Zariski topology (i.e., is a fiber bundle).*

Whereas the former question (as whether p is a variable of the polynomial ring \mathbb{C}^4) is a particular case (with $n = 4$ and $k = 3$) of the famous

Abhyankar-Sathaye Embedding Problem: *Is it true that any biregular embedding $\mathbb{C}^k \hookrightarrow \mathbb{C}^n$ is rectifiable i.e., is equivalent to a linear one under the action of the group $\text{Aut } \mathbb{C}^n$ on \mathbb{C}^n ?*

Geometrically, the situation can be regarded as follows (cf. [15]). The morphism $\sigma : (x, y, z, u) \mapsto (x, y, z)$ represents the 3-fold X as a birational modification of $Y := \mathbb{C}^3$. The latter essentially consists in replacing the divisor $D := D(f) \subseteq Y$ by the exceptional one $E := \{f = g = 0\} \subseteq X$. All the important properties of X can be recovered in terms of the restriction $\sigma|_E : E \rightarrow D$. In our setting both E and D are cylindrical surfaces, namely $E = C \times \mathbb{C}$ and $D = \Gamma \times \mathbb{C}$, where $C := D(f) \cap D(g) \subset Y$ is the center of the blowing up σ and $\Gamma := f^{-1}(0) \subseteq \mathbb{C}_{x,y}^2$. This makes it possible to formulate the criteria mentioned above in terms of the natural projection $\pi : C \rightarrow \Gamma$, $(x, y, z) \mapsto (x, y)$, and enables us in concrete examples to verify these criteria.

Let us briefly describe the content of the paper. Section 1 contains preliminaries; it can be omitted at the first reading and consulted when necessary. However, some results obtained here (and used later on in the proofs) are of independent interest.

For instance, this concerns 1.3 where we treat the question for as when a birational extension of a UFD is again a UFD. Furthermore, generalizing an observation due to V. Shpilrain and J.-T. Yu [32, 33] we claim in 1.31-1.32 that for arbitrary polynomials $p, q \in \mathbb{C}[x_1, \dots, x_n][y]$ the hypersurfaces in \mathbb{C}^{n+1} given respectively by the equations

$$y = q(p(y)) \quad \text{and} \quad y = p(q(y))$$

are isomorphic and moreover, 1-stably equivalent (see 4.21). We also use the fact (see 1.12) that a one-point compactification of an acyclic smooth affine variety is a homology manifold which is a homology sphere and satisfies the Alexander duality.

In section 2 we study the topology of the 3-folds X as above. More generally, we work with a 3-fold $X = \{p = fu + g = 0\} \subset Y \times \mathbb{C}$, where $f, g \in \mathbb{C}[Y]$ with Y being a smooth acyclic affine 3-fold and $D := f^{-1}(0) \subseteq Y$ being a cylinder $D = \Gamma \times \mathbb{C}$ over an affine curve Γ (whereas in subsection 2.3 Y itself is supposed to be a cylinder over an acyclic affine surface Z i.e., $Y = Z \times \mathbb{C}$, with $\Gamma \subseteq Z$). The main results of section 2 (see 2.11, 2.27 and 2.28) provide a criterion for as when such a 3-fold X is diffeomorphic to \mathbb{R}^6 .

In subsection 3.1 we determine when $X = p^{-1}(0) \subseteq \mathbb{C}^4$ with $p = fu + g$ as in (1) is an exotic \mathbb{C}^3 . The main tool used here is Derksen's version of the Makar-Limanov invariant [19, 4] described in subsection 1.3.

Subsection 3.2 is devoted to a study of embeddings $\mathbb{C}^3 \hookrightarrow \mathbb{C}^4$ given by an equation $p = fu + g = 0$ with $f \in \mathbb{C}^2 \setminus \mathbb{C}$ and $g \in \mathbb{C}^3$. In 3.21 we show that in appropriate new coordinates in the (x, y) -plane, the x -coordinate restricted to any fiber of p gives a \mathbb{C}^2 -fibration. On the other hand, the restriction of p to any hyperplane $x = \text{const}$ is a variable of the polynomial ring $\mathbb{C}[y, z, u]$ (in the latter case we say in brief that p is a *residual x -variable*).

A complete analog of Theorem 7.2 in [15] cited at the beginning holds if the polynomial $p \in \mathbb{C}^4$ is linear with respect to two (and not just one) variables. Indeed (3.24) if the 3-fold X :

$$p = a(x, y)u + b(x, y)v + c(x, y) = 0$$

in \mathbb{C}^4 is smooth and acyclic then $p \in \mathbb{C}^4$ is a variable. We give a simple criterion (in terms of the coefficients $a, b, c \in \mathbb{C}[x, y]$) for as when this is the case.

In section 4 we concentrate on the Abhyankar-Sathaye Problem for our particular class of embeddings. The main results 4.2, 4.16 of subsection 4.1 provide sufficient conditions for as when a residual x -variable $p = fu + g \in \mathbb{C}^4$ as in (1) is indeed a variable. In 2.8 we define a canonical factorization $f = f_{\text{horiz}} f_{\text{slant}} f_{\text{vert}}$, and we show (4.2) that an embedding $\mathbb{C}^3 \xrightarrow{\cong} X := p^{-1}(0) \subseteq \mathbb{C}^4$ can be rectified if either the polynomial f_{horiz} is reduced, or $f_{\text{vert}} = 1$. Moreover, in appropriate new coordinates (x, y) we have $f_{\text{non-horiz}} := f_{\text{slant}} f_{\text{vert}} \in \mathbb{C}[x]$, and p is an x -variable i.e., a variable in $\mathbb{C}[x]^{[3]}$ over $\mathbb{C}[x]$. For instance (see 4.2, 4.3) this is the case if $\deg_z g \leq 1$, or f is a power of an irreducible polynomial, or else $f \in \mathbb{C}[x]$ (the latter strengthens a result of M. Miyanishi [24, Thm. 2], where it is supposed in addition that $g \in \mathbb{C}[y, z]$). As another examples, we show (see 4.17) that the polynomials

$$p_1 := xy^2u + y + xz + xyz^2 \quad \text{and} \quad p_3 := xy^2u + y + x^2z + x^3yz^2$$

are variables of \mathbb{C}^4 . However, we do not know whether or not so is

$$p_2 := xy^2u + y + x^2z + xyz^2$$

(whereas $p_2^{-1}(\mu) \simeq \mathbb{C}^3 \ \forall \mu \in \mathbb{C}$ and p_2 is a residual x -variable and a $\mathbb{C}(x)$ -variable, see 4.9 and 4.18).

In subsection 4.2 we establish (see 4.23) that every embedding $\mathbb{C}^3 \hookrightarrow \mathbb{C}^4$ given by an equation $p = fu + g = 0$ as in (1) can be rectified in \mathbb{C}^5 .

In the last subsection 4.3 (attributed to the second author) we generalize a theorem of D. Wright [37] which says that Sathaye's Theorem holds for the embeddings $\mathbb{C}^2 \hookrightarrow \mathbb{C}^3$ given by an equation $p = fu^n + g = 0$ with $f, g \in \mathbb{C}[x, y]$ and $n \in \mathbb{N}$. Namely, it is shown in 4.27 that a residual x -variable of the form $p = fu^n + g \in \mathbb{C}^{[4]}$, where $f, g \in \mathbb{C}[x, y, z]$ and $n \geq 2$, actually is an x -variable.

1. PRELIMINARIES

1.1. Affine modifications of UFD's. We start by recalling the notion of affine modification [15]; at the same time, we introduce the notation that will be used throughout the paper.

Notation 1.1. Let Y be a reduced, irreducible affine variety over \mathbb{C} , $A = \mathbb{C}[Y]$ be the algebra of regular functions on Y , $I \subseteq A$ be a non-trivial ideal and $f \in I$ be a non-zero element of I . The affine modification of the variety Y along the divisor $D_f = f^*(0)$ with center I is the affine variety $X = \text{spec } A'$, where

$$A' := A[I/f] = \{a' = a_k/f^k \mid a_k \in I^k\} \subseteq \text{Frac } A.$$

The inclusion $A \hookrightarrow A'$ corresponds to a birational morphism $\sigma : X \rightarrow Y$ with the exceptional divisor $E = \sigma^{-1}(D) = (f \circ \sigma)^{-1}(0) \subseteq X$, where $D := \text{supp } D_f$. The restriction $\sigma|(X \setminus E) : X \setminus E \rightarrow Y \setminus D$ is an isomorphism.

Let $D = \bigcup_{i=1}^n D_i$ resp., $E = \bigcup_{i=1}^{n'} E_i$ be the decomposition into irreducible components, which we assume to be Cartier divisors. Letting

$$(2) \quad \sigma^*(D_i) = \sum_{j=1}^{n'} m_{ij} E_j, \quad i = 1, \dots, n,$$

we consider the $n \times n'$ multiplicity matrix $M_\sigma = (m_{ij})$ with non-negative integer entries. Clearly, $m_{ij} > 0 \Leftrightarrow \sigma(E_j) \subseteq D_i$.

The following simple observation will be useful in 1.4 below. We denote $\text{reg } E_j = E_j \setminus \text{sing } E_j$ and $\text{reg } D_i = D_i \setminus \text{sing } D_i$.

Lemma 1.2. *In the notation as above, suppose that the affine varieties X and Y are smooth. If $m_{ij} = 1$ then $\sigma(E_j) \not\subseteq \text{sing } D_i$. Moreover $m_{ij} = 1$ if and only if $\sigma(\text{reg } E_j) \subseteq \text{reg } D_i$ and σ sends the analytic discs in X transversal to E_j at a point $Q \in \text{reg } E_j$ biholomorphically onto analytic discs in Y transversal to D_i at the point $P := \sigma(Q) \in \text{reg } D_i$.*

Proof. We may assume that $m_{ij} > 0$ that is, $\sigma(E_j) \subseteq D_i$. For a point $Q \in E_j$ with the image $P = \sigma(Q) \in D_i$, we let $U \ni P$ resp., $V \ni Q$ be a neighborhood such that $D_i \cap U = f_i^*(0)$ resp., $E_j \cap V = h_j^*(0)$, where f_i resp., h_j is a holomorphic function in U resp., V . Then we have

$$f_i \circ \sigma|_V = h_j^{m_{ij}} \cdot h \quad \text{with } h(Q) \neq 0,$$

which gives the equality of 1-forms:

$$(3) \quad d(f \circ \sigma)(Q) = m_{ij} h_j^{m_{ij}-1}(Q) h(Q) \cdot dh_j(Q).$$

Assuming that $Q \in \text{reg } E_j$ ($\Leftrightarrow dh_j(Q) \neq 0$) and $m_{ij} = 1$, from (3) we obtain:

$$d(f \circ \sigma)(Q) = h(Q) \cdot dh_j(Q) \neq 0,$$

whence $df_i(P) \neq 0$ and $d\sigma(T_Q X) \not\subseteq T_P D_i$. This yields the implication " \Rightarrow ". On the other hand, if $P \in \text{reg } D_i$ ($\Leftrightarrow df_i(P) \neq 0$) and $d\sigma(T_Q X) \not\subseteq T_P D_i$ then $d(f \circ \sigma)(Q) \neq 0$, and so by (3) $m_{ij} = 1$, which gives " \Leftarrow ". \square

For an algebra B , denote by B^* its group of invertible elements.

Proposition 1.3. *In the notation as in 1.1 above, assume moreover that the algebras $A = \mathbb{C}[Y]$ and $A' = \mathbb{C}[X]$ are UFD's, and $A^* = A'^*$. Then $n = n'$ and the multiplicity matrix M_σ is unimodular.*

Proof. Since the algebras A and A' are UFD's there exist irreducible elements $f_1, \dots, f_n \in A$ resp., $h_1, \dots, h_{n'} \in A'$ such that $D_i = f_i^*(0)$, $i = 1, \dots, n$, resp., $E_j = h_j^*(0)$, $j = 1, \dots, n'$. Clearly, $h_j = (a_j/b_j) \circ \sigma$ with coprime $a_j, b_j \in A$, $j = 1, \dots, n'$. The regular functions a_j, b_j do not vanish in $Y \setminus D$, and hence their irreducible factors are proportional to some of the elements f_1, \dots, f_n . Thus for each $j = 1, \dots, n'$ there exists $\gamma_j \in A^*$ such that

$$(4) \quad h_j = \left(\gamma_j \prod_{i=1}^n f_i^{m'_{ji}} \right) \circ \sigma$$

with $m'_{ji} \in \mathbb{Z}$. On the other hand, for each $i = 1, \dots, n$ there exists $\delta_i \in A'^* = A^*$ such that

$$(5) \quad f_i \circ \sigma = \delta_i \prod_{k=1}^{n'} h_k^{m_{ik}}.$$

Plugging successively (4) and (5) one into another and taking into account the assumption that the algebras A' and A are UFD's and σ is a birational morphism, we obtain the equalities

$$(6) \quad \sum_{k=1}^{n'} m_{ik} m'_{kj} = \delta_{ij}, \quad i, j = 1, \dots, n,$$

$$(7) \quad \sum_{i=1}^n m'_{ji} m_{ik} = \delta_{jk}, \quad j, k = 1, \dots, n',$$

that is, $M'_\sigma \cdot M_\sigma = I_n$ and $M_\sigma \cdot M'_\sigma = I_{n'}$, where I_k denotes the identity matrix of order k . Hence $n = n'$ and $M'_\sigma = M_\sigma^{-1}$, so M_σ is unimodular. \square

Proposition 1.4. *Suppose that the varieties X and Y as in 1.1 are smooth, and the multiplicity matrix M_σ is an upper triangular unipotent square matrix:*

$$n = n', \quad m_{ii} = 1, \quad i = 1, \dots, n, \quad m_{ij} = 0 \quad \forall i > j.$$

Then $\sigma_ : \pi_1(X) \rightarrow \pi_1(Y)$ is an isomorphism.*

Proof. Letting $G = \pi_1(Y \setminus D)$, $G' = \pi_1(X \setminus E)$ and $\sigma' = \sigma|_{X \setminus E}$, we have that $\sigma'_* : G' \rightarrow G$ is an isomorphism which sends the subgroup

$$H' := \langle \langle \alpha_{E_1}, \dots, \alpha_{E_n} \rangle \rangle \subseteq G'$$

into the subgroup

$$H := \langle\langle \alpha_{D_1}, \dots, \alpha_{D_n} \rangle\rangle \subseteq G,$$

where for a hypersurface Z in a complex manifold X , α_Z denotes a vanishing loop of Z in $X \setminus Z$, whereas for a group G and elements $a_1, \dots, a_k \in G$, $\langle\langle a_1, \dots, a_k \rangle\rangle$ denotes the minimal normal subgroup of G generated by a_1, \dots, a_k . Moreover, $\sigma_* : \pi_1(X) \rightarrow \pi_1(Y)$ is a surjection with kernel $\ker \sigma_* \cong H/\sigma'_*(H')$ (see e.g., [15, the proof of Prop. 3.1]). Thus we must show that $\sigma'_*(H') = H$.

For $i = 1, \dots, n$ denote

$$H_i := \langle\langle \alpha_{D_1}, \dots, \alpha_{D_i} \rangle\rangle \subseteq G \quad \text{resp.}, \quad H'_i := \langle\langle \alpha_{E_1}, \dots, \alpha_{E_i} \rangle\rangle \subseteq G',$$

so that $H_n = H$ and $H'_n = H'$. Clearly, $\sigma'_*(\alpha_{E_1}) \sim \alpha_{D_1}$ (where \sim stands for conjugation), whence $\sigma'_*(H'_1) = H_1$. We show by induction that $\sigma'_*(H'_i) = H_i$ for all $i = 1, \dots, n$.

Assume that this equality holds for $i = k - 1 < n$. As $m_{kk} = 1$, by 1.2 we may conclude that for a general point $Q \in E_k$, $P := \sigma(Q)$ is a smooth point of D_k , and σ sends biholomorphically a smooth analytic disc transversal to the divisor E_k at Q onto a transversal disc to D_k at P . As the matrix M_σ is upper triangular we obtain $\sigma'_*(H'_i) \subseteq H_i$ ($i = 1, \dots, n$) and furthermore, $\sigma'_*(\alpha_{E_k}) \sim \alpha_{D_k} \pmod{H_{k-1}}$. As $\sigma'_*(H'_{k-1}) = H_{k-1}$ it follows that also $\sigma'_*(H'_k) = H_k$, therefore $\sigma'_*(H') = H$, as desired. \square

1.2. Acyclic varieties. Recall that the acyclicity of a topological space X means that its reduced homology vanishes: $\tilde{H}_*(X) := \tilde{H}_*(X; \mathbb{Z}) = 0$. The following proposition is an immediate corollary of the above results.

Proposition 1.5. *Assume that the affine varieties X and Y as in 1.1 above are smooth and acyclic. Then*

- (a) $n = n'$ and the multiplicity matrix M_σ is unimodular.

If moreover, M_σ is an upper triangular unipotent matrix then

- (b) $\sigma_* : \pi_1(X) \rightarrow \pi_1(Y)$ is an isomorphism.
(c) X is contractible if and only if so is Y .
(d) In the latter case the varieties X and Y are both diffeomorphic to the affine space \mathbb{R}^{2m} provided that $m := \dim_{\mathbb{C}} Y \geq 3$.

Proof. By [8, 1.18-1.20] (see also [11, 3.2]) for any smooth acyclic affine variety Z the algebra $\mathbb{C}[Z]$ is UFD and its invertible elements are constants. Thus (a) follows from 1.3. (b) directly follows from 1.4. In virtue of the Hurewicz and Whitehead theorems [7, Ch.2, §11.5, §14.2], an acyclic manifold is contractible if and only if it is simply connected. Hence (c) follows from (b). In turn, (d) follows from the Dimca-Ramanujam theorem [3, 39]. \square

1.6. In section 3.2 we will apply the following corollary (see 1.7 below) of Miyanishi's characterization of \mathbb{C}^3 [24, 25]. On the other hand, this corollary also follows from [31] and [16], as stated in [16, Cor. 0.2]. Moreover, by [12, L. III] it would be enough to suppose in 1.7 that only general (rather than all) fibers of x were isomorphic to \mathbb{C}^2 .

Theorem 1.7. *Let X be a smooth acyclic affine 3-fold. Then $X \simeq \mathbb{C}^3$ if and only if there exists a regular function $x \in \mathbb{C}[X]$ with all fibers isomorphic to \mathbb{C}^2 .*

1.8. It is known [15, Thm.1.1] that any birational morphism $\sigma : X \rightarrow Y$ of affine varieties is an affine modification. The divisor $D \subseteq Y$ of modification and the exceptional divisor $E \subseteq X$ of σ can be defined as minimal reduced divisors such that the restriction $\sigma|_{X \setminus E} : X \setminus E \rightarrow Y \setminus D$ is an isomorphism. In the next proposition and its corollary we provide conditions (more general than those in [15, Thm. 3.1]) which guarantee preservation of the homology group under modification. (Note that the divisors \hat{E} and \hat{D} below do not need to satisfy the assumption of minimality.)

Proposition 1.9. *Given affine varieties \hat{X}, \hat{Y} and decompositions*

$$\hat{X} = (\hat{X} \setminus \hat{E}) \cup \hat{E} \quad \text{and} \quad \hat{Y} = (\hat{Y} \setminus \hat{D}) \cup \hat{D},$$

where $\hat{E} \subseteq \hat{X}$ and $\hat{D} \subseteq \hat{Y}$ are reduced divisors, let $\sigma : \hat{X} \rightarrow \hat{Y}$ be a birational morphism which respects these decompositions. Suppose that the following hold.

- (i) \hat{E}, \hat{D} are topological manifolds, $\hat{E} \subseteq \text{reg } \hat{X}$, $\hat{D} \subseteq \text{reg } \hat{Y}$, and $\sigma^*(\hat{D}) = \hat{E}$.
- (ii) The induced homomorphisms

$$(\sigma|_{\hat{E}})_* : H_*(\hat{E}) \rightarrow H_*(\hat{D}) \quad \text{and} \quad (\sigma|_{\hat{X} \setminus \hat{E}})_* : H_*(\hat{X} \setminus \hat{E}) \rightarrow H_*(\hat{Y} \setminus \hat{D})$$

are isomorphisms.

Then $\sigma_* : H_*(\hat{X}) \rightarrow H_*(\hat{Y})$ is an isomorphism as well; in particular, \hat{X} is acyclic if and only if \hat{Y} is so.

Proof. We apply the Thom isomorphism [5, 7.15] to the pairs (\hat{X}, \hat{E}) resp., (\hat{Y}, \hat{D}) (notice that locally near \hat{E} resp., \hat{D} these are pairs of topological manifolds). As $\sigma^*(\hat{D}) = \hat{E}$ and $(\sigma|_{\hat{E}})_* : H_0(\hat{E}) \xrightarrow{\cong} H_0(\hat{D})$, σ maps the irreducible components of \hat{E} into those of \hat{D} providing a one-to-one correspondence, and (as in 1.2) sends their transverse classes [5, Ch. VIII] to the corresponding transverse classes. By functoriality of the cap-product [5, VII.12.6] for every $i \geq 0$ the following diagram is commutative:

$$\begin{array}{ccc} H_i(\hat{X}, \hat{X} \setminus \hat{E}) & \xrightarrow[t_{\hat{X}}]{\cong} & \tilde{H}_{i-2}(\hat{E}) \\ \downarrow \sigma_* & & \downarrow \sigma_* \\ H_i(\hat{Y}, \hat{Y} \setminus \hat{D}) & \xrightarrow[t_{\hat{Y}}]{\cong} & \tilde{H}_{i-2}(\hat{D}) \end{array}$$

(where t stands for the Thom isomorphism, and the homology groups in negative dimensions are zero). This allows to replace the relative homology groups in the exact sequences of pairs as to obtain the following commutative diagram:

$$\begin{array}{ccccccccc} \dots & \rightarrow & \tilde{H}_{i-1}(\hat{E}) & \rightarrow & \tilde{H}_i(\hat{X} \setminus \hat{E}) & \rightarrow & \tilde{H}_i(\hat{X}) & \rightarrow & \tilde{H}_{i-2}(\hat{E}) & \rightarrow & \tilde{H}_{i-1}(\hat{X} \setminus \hat{E}) & \rightarrow & \dots \\ & & \downarrow \sigma_* & & \downarrow \sigma_* & & \downarrow \sigma_* & & \downarrow \sigma_* & & \downarrow \sigma_* & & \\ \dots & \rightarrow & \tilde{H}_{i-1}(\hat{D}) & \rightarrow & \tilde{H}_i(\hat{Y} \setminus \hat{D}) & \rightarrow & \tilde{H}_i(\hat{Y}) & \rightarrow & \tilde{H}_{i-2}(\hat{D}) & \rightarrow & \tilde{H}_{i-1}(\hat{Y} \setminus \hat{D}) & \rightarrow & \dots \end{array}$$

By (ii) the four vertical arrows (as shown at the diagram) are isomorphisms, whence by the 5-lemma, the middle one is so as well, as stated. \square

Actually, the divisors E and D we deal with in section 2 below are not always topological manifolds. However, in our setting we can apply 1.9 by decomposing further as follows.

Corollary 1.10. *The same conclusion as in 1.9 holds if (instead of (i), (ii)) we assume that there are decompositions $\hat{E} = E' + E''$ and $\hat{D} = D' + D''$ satisfying the following conditions:*

- (i') $E' \setminus E''$, $D' \setminus D''$, E'' and D'' are topological manifolds, $\hat{E} \subseteq \text{reg } \hat{X}$, $\hat{D} \subseteq \text{reg } \hat{Y}$, and $\sigma^*(D' \setminus D'') = E' \setminus E''$, $\sigma^*(D'') = E''$.
- (ii') The induced homomorphisms $(\sigma_{\hat{X} \setminus \hat{E}})_* : H_*(\hat{X} \setminus \hat{E}) \rightarrow H_*(\hat{Y} \setminus \hat{D})$,
 $(\sigma|_{E' \setminus E''})_* : H_*(E' \setminus E'') \rightarrow H_*(D' \setminus D'')$ and $(\sigma|_{E''})_* : H_*(E'') \rightarrow H_*(D'')$
are isomorphisms.

Proof. Indeed, 1.9 implies that (under our assumptions)

$$(\sigma|_{\hat{X} \setminus E''})_* : H_*(\hat{X} \setminus E'') \rightarrow H_*(\hat{Y} \setminus D'')$$

is an isomorphism. Now (with \hat{E} and \hat{D} in 1.9 replaced by E'' resp., D''), 1.9 implies that $\sigma_* : H_*(\hat{X}) \rightarrow H_*(\hat{Y})$ is an isomorphism as well. \square

1.11. Recall [23, Ch. 5, §5.3], [9, Ch. 2, §8.1]¹ that a simplicial polyhedron P is called a *homology n -manifold* if for any point $p \in P$ we have $H_*(P, P \setminus \{p\}) \cong \tilde{H}_*(S^n)$ ².

Proposition 1.12. *Let X be a smooth acyclic affine variety of complex dimension n . Then the one-point compactification \dot{X} of X is a homology $2n$ -manifold which is a homology $2n$ -sphere: $H_*(\dot{X}) \cong H_*(S^{2n})$. In particular, the Alexander duality holds for \dot{X} .*

Proof. Notice first that X is diffeomorphic to the interior of a compact manifold, say, X_R with boundary ∂X_R (indeed, one can take for X_R the intersection of X with a ball of a large enough radius R in an affine space $\mathbb{C}^N \supseteq X$). Hence $\dot{X} \simeq X_R / \partial X_R \simeq X_R \cup_{\partial X_R} C(\partial X_R)$ (with CY denoting the cone over Y), and any triangulation of X_R naturally extends to those of \dot{X} .

By the Poincaré-Lefschetz duality for a manifold with boundary [23, 5.4.13],

$$H^{2n-i}(X_R) \cong H_i(X_R, \partial X_R) \cong \tilde{H}_i(X_R / \partial X_R) \cong \tilde{H}_i(\dot{X}),$$

whence \dot{X} is a homology $2n$ -sphere. Using the acyclicity of X_R and of $C(\partial X_R)$ and applying the Mayer-Vietoris sequence to the decomposition $\dot{X} = X_R \cup_{\partial X_R} C(\partial X_R)$, we see that $H_i(\dot{X}) \cong H_{i-1}(\partial X_R)$. Thus the smooth manifold ∂X_R is a homology $(2n-1)$ -sphere.

Clearly, $H_*(\dot{X}, \dot{X} \setminus \{x\}) \cong \tilde{H}_*(S^{2n})$ for any point $x \in X = \dot{X} \setminus \{\infty\}$. For the vertex $x = \infty$ of the cone $C(\partial X_R)$ and for any $i \in \mathbb{N}$, by excision and from the exact homology sequence of a pair we obtain:

$$H_i(\dot{X}, \dot{X} \setminus \{\infty\}) \cong H_i(C(\partial X_R), C(\partial X_R) \setminus \{\infty\})$$

¹We are grateful L. Guillou for useful discussions on homology manifolds and references.

²Or equivalently, $H_*(lk(p)) \cong H_*(S^{n-1})$, where $lk(p)$ denotes the link of p in P , or else, for any q -simplex σ in P , $H_*(lk(\sigma)) \cong H_*(S^{n-q-1})$.

$$\cong \tilde{H}_{i-1}(\partial X_R) \cong \tilde{H}_{i-1}(S^{2n-1}) \cong \tilde{H}_i(S^{2n}).$$

Thereby \dot{X} is a homology manifold and is a homology $2n$ -sphere. For any point $x \in \dot{X}$, from the exact homology sequence of the pair $(\dot{X}, \dot{X} \setminus \{x\})$ it follows that $\dot{X} \setminus \{x\}$ is acyclic. Now the proof of the Alexander duality for the usual sphere [23, 5.3.19] goes *mutatis mutandis* for \dot{X} (cf. [2, Thm. 6.4], [36, p. 176]). \square

1.3. Digest on Makar-Limanov and Derksen invariants. These invariants (introduced in [19, 13, 4]) allow in certain cases to distinguish space-like affine varieties from the affine spaces. On this purpose, we use them in subsection 3.1 (to establish 3.6). Let us first recall the following notions and facts.

1.13. Locally nilpotent derivations. Let A be an affine domain over \mathbb{C} . A derivation $\partial \in \text{Der } A$ of A is called *locally nilpotent* (LND for short) if for each $a \in A$ there exists $n = n(a, \partial) \in \mathbb{N}$ such that $\partial^{(n)}(a) = 0$; the set of all non-zero locally nilpotent derivations of the algebra A is denoted by $\text{LND}(A)$. Given $\partial \in \text{LND}(A)$, the function $\text{deg}_{\partial}(a) := \min\{n(a, \partial) - 1\}$ is a degree function on A . The kernel $A^{\partial} = \ker \partial$ of ∂ is a ∂ -invariant subalgebra of A ; its elements are called ∂ -constants. For the proof of the following lemma see e.g., [19, 20, 13, 4], [39, §7] or [12, 5.1(6)].

Lemma 1.14. *The following statements hold:*

- (a) $\text{tr.deg } [A : A^{\partial}] = 1$.
- (b) *The subalgebra A^{∂} is algebraically closed.*
- (c) *It is factorially closed i.e., $uv \in A^{\partial} \setminus \{0\} \implies u, v \in A^{\partial}$.*
- (d) *If $u^k + v^l \in A^{\partial} \setminus \{0\}$ for some $k, l \geq 2$ then $u, v \in A^{\partial}$.*
Moreover, $p(u, v) \in A^{\partial} \setminus \mathbb{C} \implies u, v \in A^{\partial}$ for any polynomial $p \in \mathbb{C}^{[2]}$ with general fibers being irreducible and non-isomorphic to \mathbb{C} .

1.15. Invariants. The *Makar-Limanov invariant* of the algebra A is the subalgebra

$$\text{ML}(A) = \bigcap_{\partial \in \text{LND}(A)} A^{\partial} \subset A,$$

whereas the *Derksen invariant*:

$$\text{Dk}(A) = \mathbb{C} \left[\bigcup_{\partial \in \text{LND}(A)} A^{\partial} \right] \subset A$$

is the subalgebra of A generated by the ∂ -constants of all locally nilpotent derivations on A . If $\text{ML}(A) = \mathbb{C}$ (resp., $\text{Dk}(A) = A$) then we say that the corresponding invariant is trivial. This is, indeed, the case for a polynomial algebra $A = \mathbb{C}^{[n]}$ ($n \in \mathbb{N}$).

1.16. Specializations. Let X be an affine variety, and set $A = \mathbb{C}[X]$. To study the locally nilpotent derivations on A , it is possible to proceed by induction on the dimension of X . Namely, let $\partial \in \text{LND}(A)$, and let $u \in \ker \partial$ be non-zero. As $\partial(u - c) = 0 \forall c \in \mathbb{C}$, the principal ideal $(u - c)$ of the algebra A is invariant under ∂ , and so ∂ descends to the quotient $B_c = A/(u - c) = \mathbb{C}[S_c]$, where $S_c = u^{-1}(c)$ is a fiber of u . For a general $c \in \mathbb{C}$, this specialization ∂_c is a non-zero locally nilpotent derivation of the algebra B_c (see 1.17 below). Clearly, the restriction to S_c of any ∂ -constant $v \in \ker \partial$ is a ∂_c -constant.

1.17. \mathbb{C}_+ -actions. Otherwise, the above specialization can be described via the natural correspondence between the locally nilpotent derivations of the algebra A and the regular actions of the additive group \mathbb{C}_+ on the variety $X = \text{spec } A$ (e.g., see [27, 39]). Indeed, the subalgebra $\ker \partial$ coincides with the algebra of invariants A^φ of the associated \mathbb{C}_+ -action $\varphi = \varphi_\partial$. If u is a φ -invariant then clearly, the \mathbb{C}_+ -action $\varphi|_{S_c}$ on a fiber $S_c = \text{spec } B_c$ of u is associated with the above specialization ∂_c . Hence $\partial_c \in \text{LND}(B_c)$ if and only if the \mathbb{C}_+ -action $\varphi|_{S_c}$ is non-trivial.

1.18. Jacobian derivations [13, 14]. For an n -tuple of polynomials $p_1, \dots, p_{n-1}, q \in \mathbb{C}^{[n]}$, the Jacobian

$$\partial(q) := \text{jac}(p_1, \dots, p_{n-1}, q) = \frac{\partial(p_1, \dots, p_{n-1}, q)}{\partial(x_1, \dots, x_n)}$$

(regarded as a function of q , whereas the polynomials p_1, \dots, p_{n-1} are fixed) gives a derivation on the polynomial algebra $\mathbb{C}^{[n]}$. This derivation is non-zero provided that the polynomials p_1, \dots, p_{n-1} are algebraically independent. For $p := p_1$, the principal ideal $(p) \subseteq \mathbb{C}^{[n]}$ is invariant under ∂ , whence ∂ descends to a derivation of the quotient algebra $A := \mathbb{C}^{[n]}/(p) = \mathbb{C}[X]$, where $X := p^*(0) \subset \mathbb{C}^n$ ³. Denote by $\text{JLND}(A)$ the set of all locally nilpotent Jacobian derivations of the algebra A . Two derivations $\partial, \partial' \in \text{Der}(A)$ are called *equivalent* if $A^\partial = A^{\partial'}$. Actually, two derivations are equivalent iff they generate the same degree function [14]: $\text{deg}_\partial = \text{deg}_{\partial'}$. We have the following theorem.

Theorem 1.19. ([13, 14]) *If $p \in \mathbb{C}^{[n]}$ is a non-constant irreducible polynomial then every locally nilpotent derivation $\partial \in \text{LND}(A)$ (where as above, $A = \mathbb{C}^{[n]}/(p)$) is equivalent to a Jacobian locally nilpotent derivation $\partial' \in \text{JLND}(A)$. Henceforth, we have*

$$\text{ML}(A) = \bigcap_{\partial \in \text{JLND}(A)} A^\partial \quad \text{and} \quad \text{Dk}(A) = \mathbb{C} \left[\bigcup_{\partial \in \text{JLND}(A)} A^\partial \right].$$

1.20. Weight degree functions and the associated graded algebras (see e.g., [13, 14, 39]). A *weight degree function* on the algebra $A = \mathbb{C}[X] = \mathbb{C}^{[n]}/(p)$ is defined by assigning weights $d_i = d(x_i) \in \mathbb{R}$ to the variables $x_1, \dots, x_n \in \mathbb{C}^{[n]}$ and letting for $a \in A$:

$$d_A(a) := \inf \{d(q) \mid q \in \mathbb{C}^{[n]}, q|_X = a\}.$$

Here as usual

$$d(q) = \max \{d(m) \mid m \in M(q)\}$$

with $M(q)$ denoting the set of the monomials $m = c_m \prod_{i=1}^n x_i^{\alpha_i}$ of q , where $d(m) := \sum_{i=1}^n \alpha_i d_i$. The weight degree function $d_A : A \rightarrow \mathbb{R} \cup \{-\infty\}$ defines an ascending filtration $\{A_t\}_{t \in \mathbb{R}}$ of the algebra A with $A_t := \{a \in A \mid d_A(a) \leq t\}$. We also let $A'_t := \{a \in A \mid d_A(a) < t\} \subset A_t$, and we consider the *associated graded algebra*

$$\hat{A} = \bigoplus_{t \in \mathbb{R}} A_t/A'_t$$

(actually, the set of non-zero homogeneous components A_t/A'_t of the algebra \hat{A} is at most countable).

³To simplify the exposition we suppose that X is a hypersurface, whereas the results of [14] hold in a more general setting.

1.21. Associated homogeneous derivations. For a polynomial $q \in \mathbb{C}^{[n]}$, we consider its d -principal part (in other words, the *principal d -quasihomogeneous part*) $\hat{q} := \sum_{m \in M(q), d(m)=d(q)} m$. For an element $a \in A$, we let \hat{a} to be its image in the graded algebra \hat{A} (clearly, $\hat{a} \in A_t/A'_t$ with $t = d_A(a)$). Notice that $\hat{a} = \hat{q}|_{\hat{X}}$ for a polynomial $q \in \mathbb{C}^{[n]}$ such that $q|_X = a$ and $d(q) = d_A(a)$; the latter equality holds if and only if $\hat{q}|_{\hat{X}} \neq 0$.

If $\partial \in \text{Der}(A)$ is a derivation then the degree

$$d_A(\partial) := \inf \{d_A(a) - d_A(\partial a) \mid a \in A\}$$

is finite [14]. Letting

$$\hat{\partial}\hat{a} := \widehat{\partial a} \quad \text{if} \quad d_A(a) - d_A(\partial a) = d_A(\partial) \quad \text{and} \quad \hat{\partial}\hat{a} = 0 \quad \text{otherwise,}$$

and extending $\hat{\partial}$ in a natural way to a homogeneous derivation of the graded algebra \hat{A} , we obtain a correspondence $\text{Der}(A) \rightarrow \text{Der}_{\text{gr}}(\hat{A})$. It has the following properties.

Theorem 1.22. ([13, 14]) *Let $p \in \mathbb{C}^{[n]}$ be a polynomial with a non-constant, irreducible d -principal part \hat{p} . Set $\hat{X} = \hat{p}^{-1}(0) \subset \mathbb{C}^n$. Then the following hold.*

- (a) $\hat{A} \simeq \mathbb{C}[\hat{X}]$.
- (b) If $\partial \in \text{LND}(A)$ then $\hat{\partial} \in \text{LND}(\hat{A})$, and for any $a \in A^\partial$, $\hat{a} \in \hat{A}^{\hat{\partial}}$.
- (c) ⁴ For any non-zero Jacobian derivation

$$\partial = \frac{\partial(p, p_2, \dots, p_{n-1}, *)}{\partial(x_1, \dots, x_n)}$$

of the algebra $A = \mathbb{C}[X]$ there exists an equivalent one

$$\partial' = \frac{\partial(p, p'_2, \dots, p'_{n-1}, *)}{\partial(x_1, \dots, x_n)}$$

such that the d -principal parts $\hat{p}, \hat{p}'_2, \dots, \hat{p}'_{n-1}$ are algebraically independent.

- (d) If the latter condition holds then the associated derivation $\hat{\partial}' \in \text{Der}(\hat{A})$ of the associated graded algebra \hat{A} is also a Jacobian one, namely,

$$\hat{\partial}' = \frac{\partial(\hat{p}, \hat{p}'_2, \dots, \hat{p}'_{n-1}, *)}{\partial(x_1, \dots, x_n)}.$$

1.23. Graded invariants. For a graded algebra \hat{A} , we denote by $\text{Dk}_{\text{gr}}(\hat{A})$ the following ‘graded’ version of the Derksen invariant:

$$\text{Dk}_{\text{gr}}(\hat{A}) = \mathbb{C} \left[\bigcup_{\partial \in \text{LND}_{\text{gr}}(\hat{A})} \hat{A}^\partial \right].$$

The way we use in the next section the Derksen invariant (similar to that of [4, 13, 19, 39]) is based on the next simple lemma.

Lemma 1.24. *Given an irreducible hypersurface $X \subseteq \mathbb{C}^n$, suppose that the algebra $A = \mathbb{C}[X]$ is equipped with a weight degree function d_A such that the hypersurface $\hat{X} \subset \mathbb{C}^n$ is also irreducible. If*

$$\text{Dk}_{\text{gr}}(\hat{A}) \subset \hat{A}_{\leq 0} := \bigoplus_{t \leq 0} A_t/A'_t \subset \hat{A}$$

⁴A similar fact remains true for any affine domain [21].

then

$$\mathrm{Dk}(A) \subset A_0 = \{a \in A \mid d_A(a) \leq 0\}.$$

Henceforth, $X \not\cong \mathbb{C}^{n-1}$ unless $A = A_0$.

Proof. By 1.22 (a),(b) for every $\partial \in \mathrm{LND}(A)$ and for every ∂ -constant $a \in A^\partial$, we have $\hat{a} \in \ker \hat{\partial}$, where $\hat{\partial} \in \mathrm{LND}_{\mathrm{gr}}(\hat{A})$, whence $\hat{a} \in \hat{A}_{\leq 0}$. Therefore, $d_A(a) \leq 0$ i.e., $a \in A_0$, as stated. \square

1.4. Variables in polynomial rings.

1.25. For a commutative ring B , we let $B^{[n]} = B[y_1, \dots, y_n]$. A polynomial $p \in B^{[n]}$ is called a B -variable (or simply a variable if no ambiguity occurs) if it is a coordinate ($p = p_1$) of an automorphism $\beta = (p_1, p_2, \dots, p_n) \in \mathrm{Aut}_B B^{[n]}$ (thus $\beta|_B = \mathrm{id}_B$). The set of all B -variables of $B^{[n]}$ is denoted as $\mathrm{Var}_B B^{[n]}$. For $B = \mathbb{C}[x_1, \dots, x_n]$, a B -automorphism is called in brief an (x_1, \dots, x_n) -automorphism, and a B -variable is also called an (x_1, \dots, x_n) -variable.

Remark 1.26. It is easily seen that a polynomial $p \in \mathbb{C}^{[n]}$ is an (x_1, \dots, x_k) -variable (where $0 \leq k \leq n-1$) if and only if $X := p^{-1}(0) \simeq \mathbb{C}^{n-1}$ and there exists $\partial \in \mathrm{LND}(\mathbb{C}^{[n]})$ with $x_1, \dots, x_k \in \ker \partial$ such that $\partial p = 1$ (i.e., p is a slice of ∂). Indeed, $\partial p = 1$ implies that X can be identified with the orbit space $\mathbb{C}^n / \varphi_\partial = \mathrm{spec}(\mathbb{C}^{[n]})^{\varphi_\partial} = \mathrm{spec}(\ker \partial)$ of the associated \mathbb{C}_+ -action φ_∂ on \mathbb{C}^n (see 1.17), and so $\mathbb{C}^{[n]} = (\ker \partial)[p] \cong \mathbb{C}^{[n-1]}[p]$.

Notice that as $\mathbb{C}^n \simeq X \times \mathbb{C}$, the assumption $X \simeq \mathbb{C}^{n-1}$ above is superfluous provided that the Zariski Cancellation Conjecture holds.

1.27. For a polynomial $q = q(x, y, z, \dots) \in \mathbb{C}^{[n]}$, we denote by q_λ ($\lambda \in \mathbb{C}$) the specialization $q(\lambda, y, z, \dots) \in \mathbb{C}^{[n-1]}$.

We say that a polynomial $p(x, y_1, \dots, y_n) \in \mathbb{C}[x][y_1, \dots, y_n]$ is a residual x -variable if for every $\lambda \in \mathbb{C}$, the specialization p_λ is a variable of $\mathbb{C}[y_1, \dots, y_n]$. It is easily seen that any x -variable is a residual x -variable.

1.28. Let $f \in B$. An element $b \in B$ is said to be invertible (resp., nilpotent) mod f if its image $[b] \in B/(f)$ is so. Thus b is nilpotent mod f if and only if $b \in \mathrm{rad}(f)$. Observe that if B is UFD and $f = \prod_{i=1}^n f_i^{a_i}$ is a canonical factorization then $b \in \mathrm{rad}(f) \Leftrightarrow b \in (f^{\mathrm{red}})$, where $f^{\mathrm{red}} := \prod_{i=1}^n f_i \in B$.

To detect variables in polynomial rings, the following results will be useful. The statement (a) below is due to P. Russell [28, Prop. 2.2], and is reproved in [35] basing on (b) (see [35, Prop. 1.4]).

Proposition 1.29. For a commutative ring B , the following hold.

- (a) For any $f, b_0, b_1 \in B$ and $q \in B[z]$ with b_1 invertible mod f and q nilpotent mod f , we have $p := fu + b_0 + b_1z + q(z) \in \mathrm{Var}_B B[z, u]$.
- (b) For $f \in B$, we let $B_f := B/(f)$, and we denote by $\rho : B \rightarrow B_f$ the canonical surjection. If $p \in \mathrm{Var}_B B[z, u]$ is such that $\rho(p(z, 0)) \in \mathrm{Var}_{B_f} B_f[z]$ then also $p_f(z, u) := p(z, fu) \in \mathrm{Var}_B B[z, u]$.
- (c) Let $B = \mathbb{C}[x]$ and $f \in B$. If $p \in \mathrm{Var}_B B[y, z, u]$ is such that for every root x_0 of f , the specialization $p_{x_0}(y, z, 0)$ is a variable of $\mathbb{C}[y, z]$ ⁵ then also $p_f(y, z, u) := p(y, z, fu) \in \mathrm{Var}_B B[y, z, u]$.

⁵Or equivalently, $\rho(p(y, z, 0)) \in \mathrm{Var}_{B_f} B_f[y, z]$.

Proof of (c). It suffices to prove (c) for $f = x$ and then conclude by induction on the degree of f . Let $\gamma \in \text{Aut}_B B[y, z, u]$ be such that $\gamma(y) = p(x, y, z, u)$. Denote γ_0 the specialization of γ at $x = 0$. Clearly, $\gamma_0(y) = p(0, y, z, u)$ is a variable of $\mathbb{C}[y, z, u]$; in particular,

$$\mathbb{C}[y, z, u]/(p(0, y, z, u)) \simeq \mathbb{C}^{[2]}.$$

If ϕ denotes this isomorphism then

$$\mathbb{C}[y, z, u]/(p(0, y, z, u), u) \simeq \mathbb{C}^{[2]}/(\phi(u)).$$

By our assumption, $p(0, y, z, 0)$ is a variable of $\mathbb{C}[y, z]$, and so ⁶

$$\mathbb{C}[y, z, u]/(p(0, y, z, u), u) \simeq \mathbb{C}[y, z]/(p(0, y, z, 0)) \simeq \mathbb{C}^{[1]}.$$

Hence by the Abhyankar-Moh-Suzuki Theorem, $\phi(u)$ is a variable in $\mathbb{C}^{[2]}$. Therefore we may assume that $\phi(u) = u$ i.e., there is a $\mathbb{C}[u]$ -isomorphism

$$\mathbb{C}[u][y, z]/(p(0, y, z, u)) \simeq \mathbb{C}[u]^{[1]}.$$

By the Abhyankar-Moh-Suzuki theorem as generalized by Russell and Sathaye [29, Thm. 2.6.2], $p(0, y, z, u)$ is a $\mathbb{C}[u]$ -variable of $\mathbb{C}[u][y, z]$. Let α_0 be the corresponding $\mathbb{C}[u]$ -automorphism of $\mathbb{C}[u][y, z]$ such that $\alpha_0(y) = p(0, y, z, u)$, and denote by $\bar{\gamma}$ the composition

$$\bar{\gamma} := \gamma\gamma_0^{-1}\alpha_0 \in \text{Aut}_B B[y, z, u].$$

Then we get

$$\bar{\gamma}(y) = \gamma\gamma_0^{-1}\alpha_0(y) = \gamma\gamma_0^{-1}(p(0, y, z, u)) = \gamma(y) = p(x, y, z, u)$$

and

$$\bar{\gamma}(u) = \gamma\gamma_0^{-1}\alpha_0(u) = \gamma\gamma_0^{-1}(u) \equiv u \pmod{x}.$$

If σ_u is the (x, y, z) -automorphism of $\mathbb{C}(x)[y, z][u]$ defined by $\sigma_u(u) = xu$ then the composition $\sigma_u\bar{\gamma}\sigma_u^{-1}$ (which *a priori* is an x -automorphism of $\mathbb{C}(x)[y, z, u]$) actually is a B -automorphism of $B[y, z, u]$. Thus $\sigma_u\bar{\gamma}\sigma_u^{-1}(y) = p(x, y, z, xu)$ is an x -variable of $\mathbb{C}[x][y, z, u]$. \square

Corollary 1.30. *Let $p \in \mathbb{C}[x][y, z, u]$ be an x -variable.*

- (a) *If for $q(x) \in \mathbb{C}[x]$, $p_q := p(x, y, z, q(x)u)$ is a residual x -variable then it actually is an x -variable.*
- (b) *Consequently, $p(x, y, z, q(x)u)$ is an x -variable if and only if $p(x, y, z, q_{\text{red}}(x)u)$ is so.*

Proof. (a) As p_q is a residual x -variable, for every root x_0 of q , $p_{x_0,0} := p(x_0, y, z, 0)$ is a variable of $\mathbb{C}[y, z, u]$. In particular, $\mathbb{C}[y, z, u]/(p_{x_0,0}) = \mathbb{C}[y, z]/(p_{x_0,0}) \otimes \mathbb{C}[u] \cong \mathbb{C}^{[2]}$. It follows that $\mathbb{C}[y, z]/(p_{x_0,0}) \cong \mathbb{C}^{[1]}$, and then by the Abhyankar-Moh-Suzuki Theorem $p_{x_0,0}$ is a variable of $\mathbb{C}[y, z]$. It follows from 1.29(c) that $p(x, y, z, q(x)u)$ is an x -variable, as stated.

The proof of (b) relies on (a) and on the fact that $p(x, y, z, q(x)u)$ and $p(x, y, z, q_{\text{red}}(x)u)$ are simultaneously residual x -variables. \square

The proof of the following results is inspired by those of [32, Thm. 1.1] and [33, Thm. 1.4].

Proposition 1.31. *For any commutative ring B , the following hold.*

⁶Hereafter under (f, g) we mean the ideal generated by f and g .

- (a) For arbitrary pair of polynomials $p, q \in B[y]$ there exists an automorphism $\gamma = \gamma_{p,q} \in \text{Aut}_B B[y, v]$ such that

$$\gamma((y - q(p(y)), v)) = (y - p(q(y)), v).$$

- (b) In particular, for $q(y) = -ay$ with $a \in B$ we have:

$$\gamma((y + ap(y), v)) = (y + p(ay), v).$$

Moreover, if $a^k | p(0)$ then there exists an automorphism $\gamma_k \in \text{Aut}_B B[y, v]$ such that

$$\gamma_k((y + ap(y), v)) = (y + \frac{p(a^{k+1}y)}{a^k}, v).$$

Proof. It is not difficult to verify that the desired automorphism γ and its inverse γ^{-1} can be defined as follows :

$$\begin{cases} \gamma(y) = v + q(y) \\ \gamma(v) = y - p(v + q(y)) \end{cases} \quad \begin{cases} \gamma^{-1}(y) = v + p(y) \\ \gamma^{-1}(v) = y - q(v + p(y)) \end{cases}.$$

Iterating γ yields an appropriate γ_k as needed in (b). \square

Corollary 1.32. (a) For any pair $p, q \in B[y]$ the homomorphism

$$\begin{array}{ccc} \phi : B[y]/(y - q(p(y))) & \longrightarrow & B[y]/(y - p(q(y))) \\ y & \longmapsto & q(y) \end{array}$$

is an isomorphism.

- (b) In particular, for any $a \in B$ and $p \in B[y]$ the homomorphism

$$\begin{array}{ccc} \phi : B[y]/(y + ap(y)) & \longrightarrow & B[y]/(y + p(ay)) \\ y & \longmapsto & ay \end{array}$$

is an isomorphism. Moreover, if $a^k | p(0)$ then also

$$\begin{array}{ccc} \phi_k : B[y]/(y + ap(y)) & \longrightarrow & B[y]/(y + \frac{p(a^{k+1}y)}{a^k}) \\ y & \longmapsto & a^{k+1}y \end{array}$$

is an isomorphism.

Proof. We just apply 1.31 and the obvious isomorphism $B[y]/(f(y)) \simeq B[y, v]/(f(y), v)$. \square

1.33. We denote by $\text{SAut } \mathbb{C}^{[n]} \subset \text{Aut } \mathbb{C}^{[n]}$ the subgroup which consists of the automorphisms α with the Jacobian $\text{jac}(\alpha) = 1$.

For the polynomial extension $B^{[2]}$ of the algebra $B := \mathbb{C}[x]$, we have the following fact.

Proposition 1.34. (a) For any set of distinct points $\lambda_1, \dots, \lambda_n \in \mathbb{C}$, the homomorphism of multi-specialization

$$\begin{array}{ccc} \Phi : \text{SAut}_B B[y, z] & \longrightarrow & (\text{SAut } \mathbb{C}[y, z])^n \\ \gamma & \longmapsto & (\gamma_{\lambda_1}, \dots, \gamma_{\lambda_n}) \end{array}$$

is surjective.

(b) *The multi-specialization map*

$$\begin{aligned} \Phi' : \text{Var}_B B[y, z] &\longrightarrow (\text{Var } \mathbb{C}[y, z])^n \\ p &\longmapsto (p_{\lambda_1}, \dots, p_{\lambda_n}) \end{aligned}$$

is surjective.

Proof. By the Jung-van der Kulk decomposition theorem it is enough to show that the image of Φ contains the elements $g_\alpha \in (\text{SAut } \mathbb{C}[y, z])^n$, where

$$g_\alpha := (\text{Id}, \dots, \text{Id}, \alpha, \text{Id}, \dots, \text{Id}),$$

and $\alpha \in \text{SAut } \mathbb{C}[y, z]$ is a triangular automorphism of the form

$$\text{either } \begin{cases} \alpha(y) = y + r(z) \\ \alpha(z) = z \end{cases} \quad \text{or} \quad \begin{cases} \alpha(y) = y \\ \alpha(z) = z + s(y). \end{cases}$$

Indeed, any automorphism $\beta_c : (y, z) \mapsto (cy + r(z), c^{-1}z)$ with $c \in \mathbb{C}^*$ can be decomposed as $\beta_c = \delta_c \circ \alpha$ with $\alpha : (y, z) \mapsto (y + c^{-1}r(z), z)$ as above and $\delta_c : (y, z) \mapsto (cy, c^{-1}z)$. In turn, we have $\delta_c = \gamma_1 \circ \gamma_{-c}$, where $\gamma_c : (y, z) \mapsto (cz, -c^{-1}y)$ can be written as $\gamma_c = \alpha'_c \circ \alpha''_c \circ \alpha'_c$ with

$$\alpha'_c : (y, z) \mapsto (y, z - c^{-1}y) \quad \text{and} \quad \alpha''_c : (y, z) \mapsto (y + cz, z).$$

Without loss of generality we may assume that $g_\alpha = (\alpha, \text{Id}, \dots, \text{Id})$. Let $\varphi \in B = \mathbb{C}[x]$ be such that $\varphi(\lambda_1) = 1$, $\varphi(\lambda_2) = \dots = \varphi(\lambda_n) = 0$, and consider the triangular automorphism $\gamma \in \text{Aut}_B B[y, z]$, where

$$\begin{cases} \gamma(y) = y + \varphi(x)r(z) \\ \gamma(z) = z \end{cases} \quad \text{resp.,} \quad \begin{cases} \gamma(y) = y \\ \gamma(z) = z + \varphi(x)s(z). \end{cases}$$

It is easily seen that, indeed, $\Phi(\gamma) = g_\alpha$. This proves (a).

(b) By (a) it suffices to show that every variable $p = \alpha(y) \in \text{Var}_\mathbb{C} \mathbb{C}[y, z]$ with $\alpha \in \text{Aut}_\mathbb{C} \mathbb{C}[y, z]$ can be regarded as a coordinate function of an automorphism $\alpha' \in \text{SAut } \mathbb{C}[y, z]$. Indeed, let $\alpha' := \alpha \circ \sigma$ with $\sigma : (y, z) \mapsto (y, j^{-1}z)$, where $j := \text{jac}(\alpha)$. Then $\alpha'(y) = \alpha(y) = p$ and $\alpha' \in \text{SAut } \mathbb{C}[x, y]$. Thus (b) follows. \square

Remark 1.35. We would like to use this opportunity to indicate a flaw in the proof of Theorem 7.2 in [15] (which generalizes the Sathaye Theorem, as mentioned in the Introduction). Namely, proving Claim 3 of Proposition 7.1 in [15] and carrying induction by the multiplicity of a root $x = 0$ of the polynomial p , it has been forgotten to extend it over all the roots. Instead, one might apply 1.34 (or 1.29(b); cf. [35, Thm. 3.6]) which would simplify the proof considerably.

2. SIMPLE MODIFICATIONS OF ACYCLIC 3-FOLDS ALONG CYLINDRICAL DIVISORS

We focus below on a special case of affine modifications called simple birational extensions (see 2.1 below), applied to acyclic affine 3-folds. Our aim in this sections is to give a criterion for as when the acyclicity is preserved under such a modification (cf. [15, Thm. 3.1]). In the special case when the divisor of modification is a cylinder, we obtain in Theorem 2.11 below necessary conditions for preserving the acyclicity. In Theorem 2.27 (cf. also 2.28) we show that these conditions are also sufficient, provided that the given acyclic 3-fold is a cylinder as well.

2.1. Simple affine modifications.

Definition 2.1. A *simple birational extension* of a domain A (over \mathbb{C}) is an algebra $A' := A[g/f]$, where $f, g \in A$ are such that the ideal $I = (f, g) \subset A$ is of height 2 (i.e., the center of modification $C = V(I) = D_f^{\text{red}} \cap D_g^{\text{red}}$ is of codimension 2 in $Y := \text{spec } A$). We also call $X := \text{spec } A'$ a *simple affine modification* of the affine variety Y .

In the sequel A is UFD, and the above condition simply means that $f, g \in A$ are coprime. More generally [15, Prop. 1.1], any affine modification can be obtained as $A' = A[g_1/f_1, \dots, g_n/f_n]$ with $f_i, g_i \in A$; here again, if A is UFD then we may suppose f_i, g_i being coprime ($i = 1, \dots, n$).

2.2. Observe that the variety $X = \text{spec } A'$ can be realized as the hypersurface in $Y \times \mathbb{C}$ with the equation $fu + g = 0$ (where u is a coordinate in \mathbb{C}), and the blowup morphism is just the first projection: $\sigma = \text{pr}_1|_X : X \rightarrow Y$. The exceptional divisor $E = \sigma^{-1}(C) = \sigma^{-1}(D) = \{f = 0\} \subseteq X$ is cylindrical: $E \simeq C \times \mathbb{C}$ (cf. [15, Prop. 1.1]).

We have the following simple but useful lemma.

Lemma 2.3. *In the notation as above, assume that the affine variety $Y := \text{spec } A$ is smooth. Then the simple affine modification $X = \text{spec } A[g/f]$ is also smooth if and only if*

- (i) *the divisor $D_g = g^*(0)$ is smooth and reduced at each point of the center $C = D_f^{\text{red}} \cap D_g^{\text{red}}$, and*
- (ii) *D_f and D_g meet transversally at those points of C where the divisor D_f is also smooth and reduced.*

Furthermore, if X is smooth then $df = 0$ at each singular point of the center C .

Proof. The second assertion easily follows from the first one. To prove the first one, let $F = 0$ be the equation of the hypersurface $X \subseteq Y \times \mathbb{C}$ with $F := uf - g \in \mathbb{C}[Y][u]$ being irreducible. As $X \setminus E \simeq Y \setminus D$ is smooth, we should only control the smoothness of X at the points of the exceptional divisor

$$E = \{f = g = 0\} = C \times \mathbb{C} \subseteq Y \times \mathbb{C}.$$

At a point $Q = (P, u) \in E$, we have:

$$dF = udf + dg + fdu = udf + dg = 0$$

if and only if $dg = -udf$ is proportional to df . Now the assertion easily follows. \square

2.2. Preserving acyclicity: necessary conditions. In this subsection we adopt the following convention and notation.

Convention 2.4. (i) X and Y denote smooth, acyclic affine 3-folds ⁷ such that

- (ii) $A' = \mathbb{C}[X]$ is a simple birational extension of the algebra $A = \mathbb{C}[Y]$ i.e., $A' = A[g/f]$ with coprime elements $f, g \in A$, and
- (iii) $D := D_f^{\text{red}} \simeq \Gamma \times \mathbb{C}$ (where Γ is an affine curve) is a cylindrical divisor.

⁷Hence the algebras A and A' are UFD's, see the proof of 1.5.

2.5. Denote by $\pi : D \rightarrow \Gamma$ the morphism induced by the canonical projection $\Gamma \times \mathbb{C} \rightarrow \Gamma$. By abuse of notation, we equally denote by π the restriction $\pi|_C : C \rightarrow \Gamma$. Thus we have the following commutative diagram:

$$\begin{array}{ccc} E \simeq C \times \mathbb{C} & \longrightarrow & C \\ \downarrow \sigma & & \downarrow \pi \\ D \simeq \Gamma \times \mathbb{C} & \longrightarrow & \Gamma \end{array}$$

If $C = \bigcup_{i=1}^{n'} C_i$ resp., $\Gamma = \bigcup_{j=1}^n \Gamma_j$ is the irreducible decomposition then the irreducible components of the divisor E resp., D are $E_i \simeq C_i \times \mathbb{C}$, resp., $D_j \simeq \Gamma_j \times \mathbb{C}$. As in 1.1 we denote by $M_\sigma = (m_{ij})$ the multiplicity matrix of σ . Recall that by 1.1, 1.2 and 2.3, $m_{ij} \geq 1$ if and only if $C_j \subset D_i^{\text{red}}$, and $m_{ij} = 1$ if and only if the surfaces $D_i^{\text{red}}, D_j^{\text{red}} \subseteq Y$ meet transversally at general points of the curve C_j .

The terminology in the following definitions comes from the real picture corresponding to our situation.

2.6. Notice that for each $j = 1, \dots, n$ there exists $i \in \{1, \dots, n\}$ such that $\pi(C_j) \subseteq \Gamma_i$, and this index $i = i(j)$ is unique unless $\pi|_{C_j} = \text{const}$.

An irreducible component C_j of the curve C is called *vertical* if $\pi|_{C_j} = \text{const}$ (i.e. $\deg(\pi|_{C_j}) = 0$) and *non-vertical* otherwise (thus the vertical components of C are disjoint and each of them is isomorphic to \mathbb{C}). The uniqueness of the index $i = i(j)$ for a non-vertical component C_j and the unimodularity of M_σ (see 1.5) imply that the j -th column of the matrix M_σ is the i -th vector of the standard basis $(\bar{e}_1, \dots, \bar{e}_n)$ in \mathbb{R}^n , and two different non-vertical components C_j and $C_{j'}$ of C project into two different irreducible components Γ_i resp., $\Gamma_{i'}$ of Γ . Hence up to reordering, we may assume that C_1, \dots, C_k are the non-vertical components of C and $\pi(C_i) \subseteq \Gamma_i, i = 1, \dots, k$. Then we have

$$M_\sigma = \left(\begin{array}{c|c} I_k & B \\ \hline 0 & B' \end{array} \right)$$

with a unimodular matrix B' . Consequently, for every $i = 1, \dots, k$, the surfaces D_i^{red} and D_g^{red} meet transversally at general points of the curve C_i .

2.7. The irreducible components $\Gamma_1, \dots, \Gamma_k$ are also called *non-vertical* resp., $\Gamma_{k+1}, \dots, \Gamma_n$ are called *vertical*. Among the non-vertical components C_i resp., Γ_i we distinguish those with $\deg(\pi|_{C_j}) = 1$ which we call *horizontal* and those with $\deg(\pi|_{C_j}) \geq 2$ which we call *slanted*. We reorder again to obtain that C_1, \dots, C_h resp., $\Gamma_1, \dots, \Gamma_h$ are the horizontal components of C resp., Γ , and C_{h+1}, \dots, C_k resp., $\Gamma_{h+1}, \dots, \Gamma_k$ are the slanted ones. An irreducible component of C resp., Γ which is not horizontal is referred to as a *non-horizontal* component; C_{horiz} denotes the union of all horizontal components of C . In the same way we define $C_{\text{non-horiz}}, C_{\text{vert}}, C_{\text{non-vert}}, C_{\text{slant}}, \Gamma_{\text{horiz}}, \Gamma_{\text{non-horiz}}, D_{\text{horiz}}$, etc. Thus we have:

$$\Gamma_{\text{horiz}} = \bigcup_{i=1}^h \Gamma_i, \quad \Gamma_{\text{slant}} = \bigcup_{i=h+1}^k \Gamma_i, \quad \Gamma_{\text{vert}} = \bigcup_{i=k+1}^n \Gamma_i,$$

and similarly

$$C_{\text{horiz}} = \bigcup_{i=1}^h C_i, \quad C_{\text{slant}} = \bigcup_{i=h+1}^k C_i, \quad C_{\text{vert}} = \bigcup_{i=k+1}^n C_i.$$

2.8. Let furthermore $f = \prod_{j=1}^n f_j^{a_j} = f_{\text{horiz}} \cdot f_{\text{non-horiz}}$ be factorizations such that

$$f_j^*(0) = D_j^{\text{red}} \quad (j = 1, \dots, n), \quad (f_{\text{horiz}})^{-1}(0) = D_{\text{horiz}} \simeq \Gamma_{\text{horiz}} \times \mathbb{C}$$

and

$$(f_{\text{non-horiz}})^{-1}(0) = D_{\text{non-horiz}} \simeq \Gamma_{\text{non-horiz}} \times \mathbb{C}.$$

2.9. An irreducible component is called *isolated* if it is a connected component.

Let us give a typical example which illustrates our definitions.

Example 2.10. Letting $Y = \mathbb{C}^3$ with coordinates x, y, z , set $f = xy, g = y + xz$. Then $X = \{xyu + y + xz = 0\} \subset \mathbb{C}^4 = Y \times \mathbb{C}$, $D = \{xy = 0\} \subset \mathbb{C}^3$, $\sigma(x, y, z, u) = (x, y, z)$ and $E = \{xy = y + xz = 0\} \subseteq \mathbb{C}^4$. Therefore, $\Gamma = \{xy = 0\} \subset \mathbb{C}^2$, $\Gamma_{\text{horiz}} = \{y = 0\}$ and $\Gamma_{\text{vert}} = \{x = 0\}$ whereas $C = C_{\text{vert}} \cup C_{\text{horiz}}$ with $C_{\text{horiz}} = \{y = z = 0\}$ and $\Gamma_{\text{vert}} = \{x = y = 0\}$.

The main result of this subsection is the following theorem.

Theorem 2.11. *Let X and Y be smooth acyclic affine 3-folds satisfying the conditions (ii) and (iii) of 2.4. Then (in the notation of 2.6-2.9) the following hold.*

- (α) $\pi|(C_{\text{horiz}} \setminus C_{\text{non-horiz}}) : C_{\text{horiz}} \setminus C_{\text{non-horiz}} \rightarrow \Gamma_{\text{horiz}} \setminus \Gamma_{\text{non-horiz}}$ is an isomorphism.
- (β) The slanted components C_{h+1}, \dots, C_k and $\Gamma_{h+1}, \dots, \Gamma_k$ are isolated and homeomorphic to \mathbb{C} .
- (γ) $f_{\text{non-horiz}} = p \circ f_{h+1}$ with $p \in \mathbb{C}[z]$, and every non-horizontal component of Γ is homeomorphic to the affine line \mathbb{C} . In other words,

$$f_{\text{non-horiz}} = c \cdot \prod_{i=h+1}^n (f_{h+1} - \lambda_i)^{a_i} \quad (c \in \mathbb{C}^*, \lambda_{h+1} = 0)$$

with $\pi(f_{h+1}^{-1}(\lambda_i)) = \Gamma_i$ homeomorphic to \mathbb{C} ($i = h+1, \dots, n$).

- (δ) Up to a further reordering, the multiplicity matrix has the following form:

$$M_\sigma = \left(\begin{array}{c|c|c} I_h & 0 & B_0 \\ \hline 0 & I_{k-h} & 0 \\ \hline 0 & 0 & I_{n-k} \end{array} \right)$$

Consequently (by 1.5), $\sigma_* : \pi_1(X) \rightarrow \pi_1(Y)$ is an isomorphism. Moreover, X is contractible (and hence diffeomorphic to \mathbb{R}^6) if and only if Y is so.

The rest of this subsection is devoted to the proof of Theorem 2.11. It is convenient to introduce the following terminology and notation.

2.12. Let F be a curve. We say that a point $P \in F$ is *multibranch* (resp., *unibranch*) if it is a center of $\mu_P = \mu_P(F) > 1$ (resp., $\mu_P = 1$) local analytic branches of F . We denote by F^{norm} the normalization of F and by \bar{F} its smooth complete model. The points of $\bar{F} \setminus F^{\text{norm}}$ are called the *punctures* of F . A morphism of curves $\rho : F \rightarrow G$

can be lifted to the normalizations resp., the completions; we denote the lift by $\rho^{\text{norm}} : F^{\text{norm}} \rightarrow \Gamma^{\text{norm}}$ resp., $\bar{\rho} : \bar{F} \rightarrow \bar{G}$.

2.13. Consider further an irreducible smooth curve F of genus g with n punctures, and let the 1-cycles $a_1, \dots, a_g, b_1, \dots, b_g$ on F provide a symplectic basis of the group $H_1(\bar{F}) = H_1(\bar{F}; \mathbb{Z})$. Then there is an injection $H_1(\bar{F}) \hookrightarrow H_1(F)$ onto the subgroup generated by the classes $[a_1], \dots, [a_g], [b_1], \dots, [b_g]$; we may identify the group $H_1(\bar{F})$ with its image in $H_1(F)$. The group $H_1(F)$ being freely generated by the classes $[a_1], \dots, [a_g], [b_1], \dots, [b_g], [c_1], \dots, [c_{n-1}]$, where c_1, \dots, c_n are simple 1-cycles around the punctures of F , we have a (non-canonical) decomposition

$$(8) \quad H_1(F) \simeq H_1(\bar{F}) \oplus G(F) \quad \text{with} \quad G(F) := \langle [c_1], \dots, [c_n] \rangle \simeq \mathbb{Z}^{n-1}$$

($\sum_{i=1}^n [c_i] = 0$ being the only relation between the generators $[c_1], \dots, [c_n]$ of the group $G(F)$).

2.14. We denote by $S_\Gamma = S_\Gamma^{(0)} \cup S_\Gamma^{(1)}$ (where $S_\Gamma^{(0)} \cap S_\Gamma^{(1)} = \emptyset$) the finite subset of the curve Γ such that a point P belongs to S_Γ if and only if it satisfies at least one of the following three conditions:

- (i) $P = \pi(C_i)$ for a vertical component C_i of C ($\Leftrightarrow P \in S_\Gamma^{(1)}$);
- (ii) $P = \pi(Q)$ for a multibranch point Q of C ;
- (iii) P is a multibranch point of Γ .

2.15. Set $S_C = \pi^{-1}(S_\Gamma) \subseteq C$ and $S_C^{(i)} = \pi^{-1}(S_\Gamma^{(i)})$, $i = 0, 1$. Thus the analytic set S_C contains the union $S_C^{(1)} = C_{\text{vert}}$ of vertical components of C , whereas the residue set $S_C^{(0)} = S_C \setminus S_C^{(1)}$ is finite.

It is easily seen that if $Q \in S_C^{(0)}$ then over any local analytic branch B_i of Γ at the point $P := \pi(Q)$ there is at least one local analytic branch $A_j \subseteq \pi^{-1}(B_i) \cap D_j^{\text{red}}$ of C at Q . Indeed, the surfaces $\pi^{-1}(B_i) \simeq B_i \times \mathbb{C}$ and D_j meet at Q , and so the polynomial $g|_{\pi^{-1}(B_i)} \in \mathcal{O}[B_i][z]$ vanishes at $Q =: (P, z_0)$, but its specialization at P is nonzero, as $Q \in S_C^{(0)}$ means that no vertical component of C passes through Q . Consequently $\mu_Q(C) \geq \mu_P(\Gamma)$ and $P \notin \Gamma_{\text{vert}}$, whence $\pi^{-1}(\Gamma_{\text{vert}}) \subseteq S_C^{(1)} = C_{\text{vert}}$.

2.16. We let $\Gamma^* = \Gamma \setminus S_\Gamma$ (more generally, $\Gamma_{\text{something}}^* = \Gamma_{\text{something}} \cap \Gamma^*$) and $C^* = C \setminus S_C = \pi^{-1}(\Gamma^*) = \pi^{-1}(\Gamma_{\text{non-vert}}^*) \subseteq C$.

2.17. For a complex hermitian manifold M and a closed analytic subset T of M , by a *link* $lk_P(T)$ of T at a point $P \in T$ we mean the intersection $T \cap S_\varepsilon$ of T with a small enough sphere S_ε in M centered at P . We also call *link* the corresponding homology class $[lk_P(T)] \in H_*(T \setminus \{P\}) = H_*(T \setminus \{P\}; \mathbb{Z})$, and we still denote it simply by $lk_P(T)$.

2.18. We denote by H_Γ resp., H_C the subgroup of the group $H_1(\Gamma^*) = H_1(\Gamma^*; \mathbb{Z})$ resp., $H_1(C^*)$ generated by the links⁸ $lk_P(\Gamma)$ resp., $lk_Q(C)$ of the points $P \in S_\Gamma$ resp., $Q \in S_C^{(0)}$. Notice that $lk_P(\Gamma) = \sum_{i=1}^\mu lk_P(B_i)$, where B_1, \dots, B_μ are the

⁸Here we are in the special case when $M = X$ resp., Y and $T = C$ resp. Γ .

local branches of Γ at P and $\mu := \mu_P(\Gamma)$. Clearly, $H_\Gamma \subseteq G(\Gamma^*)$ resp., $H_C \subseteq G(C^*)$, and so we obtain (non-canonical) isomorphisms

$$(9) \quad H_1(C^*)/H_C \simeq H_1(\overline{C^*}) \oplus [G(C^*)/H_C]$$

resp.,

$$(10) \quad H_1(\Gamma^*)/H_\Gamma \simeq H_1(\overline{\Gamma^*}) \oplus [G(\Gamma^*)/H_\Gamma].$$

The next proposition is our main technical tool in the proof of Theorem 2.11.

Proposition 2.19. *Under the assumptions as in 2.4, consider the restriction $\pi = \pi|_{C^*} : C^* \rightarrow \Gamma^*$. Then $\pi_*(H_C) \subseteq H_\Gamma$, and π induces the following isomorphisms:*

$$(11) \quad \widehat{\pi}_* : H_1(C^*)/H_C \xrightarrow{\cong} H_1(\Gamma^*)/H_\Gamma,$$

$$(12) \quad \overline{\pi}_* : H_1(\overline{C^*}) \xrightarrow{\cong} H_1(\overline{\Gamma^*}),$$

$$(13) \quad \widetilde{\pi}_* : G(C^*)/H_C \xrightarrow{\cong} G(\Gamma^*)/H_\Gamma.$$

The proof is based on Lemmas 2.21 and 2.22 below. Let us introduce the following notation.

2.20. Denote

$$S_E = \sigma^{-1}(S_C) = S_C \times \mathbb{C} \subseteq E \quad \text{resp.}, \quad S_E^{(i)} = \sigma^{-1}(S_C^{(i)}) \quad (i = 0, 1)$$

and

$$S_D = \pi^{-1}(S_\Gamma) \subseteq D, \quad \text{so that} \quad S_D \simeq S_\Gamma \times \mathbb{C} \subseteq \Gamma \times \mathbb{C} \simeq D.$$

Furthermore, set

$$E^* = \sigma^{-1}(C^*) = E \setminus S_E = C^* \times \mathbb{C} \quad \text{resp.}, \quad D^* = \pi^{-1}(\Gamma^*) = D \setminus S_D \simeq \Gamma^* \times \mathbb{C}$$

and

$$X^* = X \setminus S_E \quad \text{resp.}, \quad Y^* = Y \setminus S_D.$$

Thus

$$X \setminus E = X^* \setminus E^* \quad \text{resp.}, \quad Y \setminus D = Y^* \setminus D^*,$$

and so we have an isomorphism

$$\sigma|_{X^* \setminus E^*} : X^* \setminus E^* \rightarrow Y^* \setminus D^*.$$

Lemma 2.21. *There are monomorphisms*

$$(14) \quad \rho_X : H_3(X^*) \rightarrow H_1(C^*) \quad \text{resp.}, \quad \rho_Y : H_3(Y^*) \rightarrow H_1(\Gamma^*)$$

such that $\pi_* \rho_X = \rho_Y \sigma_*$ and

$$\widehat{\pi}_* : H_1(C^*)/\rho_X(H_3(X^*)) \rightarrow H_1(\Gamma^*)/\rho_Y(H_3(Y^*))$$

is an isomorphism.

Proof. The map of pairs $\sigma|_{X^*} : (X^*, X^* \setminus E^*) \rightarrow (Y^*, Y^* \setminus D^*)$ induces the following commutative diagram (where the horizontal lines are exact homology sequences of pairs):

$$\begin{array}{ccccccc}
\dots & \longrightarrow & H_i(X^* \setminus E^*) & \longrightarrow & H_i(X^*) & \xrightarrow{i_*} & H_i(X^*, X^* \setminus E^*) & \xrightarrow{\partial_*} & H_{i-1}(X^* \setminus E^*) & \longrightarrow & \dots \\
& & \downarrow \wr \sigma_* & & \downarrow \sigma_* & & \downarrow \sigma_* & & \downarrow \wr \sigma_* & & \\
\dots & \longrightarrow & H_i(Y^* \setminus D^*) & \longrightarrow & H_i(Y^*) & \xrightarrow{i_*} & H_i(Y^*, Y^* \setminus D^*) & \xrightarrow{\partial_*} & H_{i-1}(Y^* \setminus D^*) & \longrightarrow & \dots
\end{array} \tag{*1}$$

Claim. (a) $H_3(X^* \setminus E^*) = H_3(Y^* \setminus D^*) = 0$ and (b) $H_2(X^*) = H_2(Y^*) = 0$.

Proof of the claim. (a) In virtue of the isomorphism

$$\sigma_* : H_3(X^* \setminus E^*) = H_3(X \setminus E) \xrightarrow{\cong} H_3(Y \setminus D) = H_3(Y^* \setminus D^*)$$

it is enough to show the vanishing of one of these groups, say, of $H_3(X \setminus E)$. Let as before, \bar{F} denote the one-point compactification of a topological space F . As X is acyclic (whence by 1.12, the Alexander duality can be applied to \bar{X}) and moreover E is closed in X , the Alexander duality gives an isomorphism

$$(15) \quad H_3(X \setminus E) \cong H^2(\dot{E}, \{\infty\}).$$

Consider the homeomorphisms $E \approx C \times \mathbb{R}^2$, $\dot{E} \approx (\dot{C} \times \mathbb{R}^2)/(\dot{C} \vee \mathbb{R}^2)$ and replace here \mathbb{R}^2 by S^2 . Since $(\dot{C} \times S^2, \dot{C} \vee S^2)$ has the homotopy type of a pair of cell complexes, by [5, 4.4] we get

$$H^2(\dot{E}, \{\infty\}) \cong \tilde{H}^2(\dot{E}) \cong H^2(\dot{C} \times S^2, \dot{C} \vee S^2).$$

The Künneth formula for cohomology [22, (11.2)] yields a monomorphism

$$\mu : \sum_{p+q=2} H^p(\dot{C}, \{\infty\}) \otimes H^q(S^2, \{\infty\}) =: H \rightarrow H^2(\dot{C} \times S^2, \dot{C} \vee S^2)$$

with the cokernel

$$\text{coker } \mu = \sum_{p+q=3} \text{Tor}(H^p(\dot{C}, \{\infty\}), H^q(S^2, \{\infty\})).$$

As

$$H^0(\dot{C}, \{\infty\}) = H^0(S^2, \{\infty\}) = H^1(S^2, \{\infty\}) = 0$$

we have $H = 0$. The group $H^*(S^2, \{\infty\})$ being torsion free, we also have $\text{coker } \mu = 0$, and so $H^2(\dot{C} \times S^2, \dot{C} \vee S^2) = 0$ as well. Thus in view of (15), $H_3(X \setminus E) = 0$. This proves (a).

(b) By the Alexander duality we obtain

$$H_2(X^*) = H_2(X \setminus S_E) \cong H^3(\dot{S}_E, \{\infty\}) = H^3(\dot{S}_E).$$

The topological space \dot{S}_E is homeomorphic to a bouquet of 4-spheres S^4 and 2-spheres S^2 . The 4-spheres are provided by the one-point compactification of the product $S_C^{(1)} \times \mathbb{C} = C_{\text{vert}} \times \mathbb{C}$ (recall (see 2.6) that the components of the curve C_{vert} are disjoint and each one is isomorphic to \mathbb{C}), whereas the 2-spheres S^2 are provided by the one-point compactification of the product $S_C^{(0)} \times \mathbb{C}$.

Hence $H^3(\dot{S}_E) \cong H_2(X^*) = 0$. Similarly, we have $H_2(Y^*) = 0$. This proves the claim. \square

In virtue of the above claim, $(*_1)$ (with $i = 3$) leads to the commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & H_3(X^*) & \xrightarrow{i_*} & H_3(X^*, X^* \setminus E^*) & \xrightarrow{\partial_*} & H_2(X^* \setminus E^*) \longrightarrow 0 \\
& & \downarrow \sigma_* & & \downarrow \sigma_* & & \downarrow \wr \sigma_* \\
0 & \longrightarrow & H_3(Y^*) & \xrightarrow{i_*} & H_3(Y^*, Y^* \setminus D^*) & \xrightarrow{\partial_*} & H_2(Y^* \setminus D^*) \longrightarrow 0
\end{array} \quad (*_2)$$

Next we apply the Thom isomorphism [5, 7.15] to the pairs of manifolds (X^*, E^*) resp., (Y^*, D^*) (cf. the proof of 1.9). Indeed, the curve C^* resp., Γ^* being locally unibranch (whence homeomorphic to its normalization), $E^* = C^* \times \mathbb{C}$ resp., $D^* = \Gamma^* \times \mathbb{C}$ is a topological manifold. Notice that (in virtue of 1.2 and 2.6) σ^* maps the transverse classes [5, Ch. VIII] of E^* into those of D^* . By functoriality of the cap-product [5, VII.12.6], for every $i \geq 2$ the following diagram is commutative:

$$\begin{array}{ccc}
H_i(X^*, X^* \setminus E^*) & \xrightarrow[t_X]{\cong} & \tilde{H}_{i-2}(E^*) \\
\downarrow \sigma_* & & \downarrow \sigma_* \\
H_i(Y^*, Y^* \setminus D^*) & \xrightarrow[t_Y]{\cong} & \tilde{H}_{i-2}(D^*)
\end{array} \quad (*_3)$$

By making use of $(*_3)$ together with the following commutative diagram (see 2.5):

$$\begin{array}{ccc}
H_*(E^*) & \xrightarrow{\cong} & H_*(C^*) \\
\downarrow \sigma_* & & \downarrow \pi_* \\
H_*(D^*) & \xrightarrow{\cong} & H_*(\Gamma^*)
\end{array} \quad (*_4)$$

we may replace in $(*_2)$ the group $H_3(X^*, X^* \setminus E^*)$ resp., $H_3(Y^*, Y^* \setminus D^*)$ by the group $H_1(C^*)$ resp., $H_1(\Gamma^*)$ to obtain a commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & H_3(X^*) & \xrightarrow{\rho_X} & H_1(C^*) & \xrightarrow{\partial_*} & H_2(X^* \setminus E^*) \longrightarrow 0 \\
& & \downarrow \sigma_* & & \downarrow \pi_* & & \downarrow \wr \\
0 & \longrightarrow & H_3(Y^*) & \xrightarrow{\rho_Y} & H_1(\Gamma^*) & \xrightarrow{\partial_*} & H_2(Y^* \setminus D^*) \longrightarrow 0
\end{array} \quad (*_5)$$

where $\rho_X := \sigma_* \circ t_X \circ i_*$ resp., $\rho_Y := \pi_* \circ t_Y \circ i_*$. The diagram $(*_5)$ yields the assertions of the lemma. \square

Lemma 2.22. $\rho_X(H_3(X^*)) = H_C$ and $\rho_Y(H_3(Y^*)) = H_\Gamma$.

Proof. We start by constructing an appropriate free base of the \mathbb{Z} -module $H_3(X^*)$ (resp., $H_3(Y^*)$). By the Alexander duality we have isomorphisms

$$\tilde{H}_3(X^*) \cong H^2(\dot{S}_E) \cong H^2(\dot{S}_E^{(0)}) \cong \mathbb{Z}^{b_0(S_C^{(0)})},$$

where $S_E^{(0)} := S_C^{(0)} \times \mathbb{C}$ (similarly, $\tilde{H}_3(Y^*) \cong \mathbb{Z}^{b_0(S_r)}$). Thus $H_3(X^*) = \tilde{H}_3(X^*) \cong H^2(\dot{S}_E^{(0)})$ is a free \mathbb{Z} -module, and so the universal coefficient formula provides yet another form of the Alexander duality:

$$H_3(X^*) \cong H^2(\dot{S}_E^{(0)}) \cong H_2(\dot{S}_E^{(0)}) \quad (\text{resp., } H_3(Y^*) \cong H_2(\dot{S}_D))$$

(see e.g., [7, Ch. 2, §15.5]). Now a free base of $H_3(X^*)$ can be reconstructed as follows. On each component $T_Q := \sigma^{-1}(Q) = \{Q\} \times \mathbb{C} \simeq \mathbb{C}$ of $S_E^{(0)}$ (where $Q \in S_C^{(0)}$) fix a point $Q' \in T_Q$, and fix a complex 2-ball $B_{Q'}^2$ in X transversal to T_Q at Q' . Then the 3-cycle $[s_{Q'}] \in H_3(X^*)$ which corresponds to the 2-cycle $[T_Q] \in H_2(\dot{S}_E^{(0)})$ by the Alexander duality, can be represented by a small 3-sphere (say) $s_{Q'}$ in $B_{Q'}^2$, centered at Q' [7, Ch. 2, §17]. These cycles $[s_{Q'}] \in H_3(X^*)$ with $\pi(Q') = Q$, where Q runs over $S_C^{(0)}$ form the desired base.

It is convenient to fix the choice of a point $Q' \in T_Q$ as follows. Consider the curve $C' \subseteq X$ given by the equations $u = f \circ \sigma = 0$ (thus $C' = E \cap D_u^{\text{red}}$). It is easily seen that the restriction $\sigma|_{C'} : C' \rightarrow C$ is an isomorphism. Its inverse $C \rightarrow C' \subseteq E$ is a section of the projection $\sigma|_E : E = C \times \mathbb{C} \rightarrow C$. Letting $Q' := T_Q \cap C'$ and placing the transversal ball $B_{Q'}^2$ into the divisor $D_u^{\text{red}} = \{u = 0\} \subseteq X$ (notice that by 2.3, Q' is a smooth point of $D_u^{\text{red}} \simeq D_g^{\text{red}}$), we let $\delta_{Q'} := s_{Q'} \cap C'$. Then $\sigma(\delta_{Q'}) =: \delta_Q = lk_Q(C)$ is a link of the curve C at the point $Q \in C$, and so $\delta_Q \in H_C$. We claim that $\rho_X([s_{Q'}]) = [\delta_Q] \in H_1(C^*)$, or in other words, that $i_*([s_{Q'}]) = t_X^{-1}([\delta_{Q'}])$.

Indeed, consider the Zariski open dense subsets $X^{**} := X^* \setminus \text{sing } E^* \subseteq X^*$ resp., $\text{reg } E^* := E^* \setminus \text{sing } E^* \subseteq E^*$. Let U be a small tubular neighborhood of the closed submanifold $\text{reg } E^* \subseteq X^{**}$ with a retraction $\tau : U \rightarrow \text{reg } E^*$. We have $\delta_{Q'} \subseteq \text{reg } E^*$, and the class $[\tau^{-1}(\delta_{Q'})] \in H_3(X^{**}, X^{**} \setminus \text{reg } E^*) = H_3(U, U \setminus \text{reg } E^*)$ corresponds to the class $[\delta_{Q'}]$ under the Thom isomorphism $H_3(X^{**}, X^{**} \setminus \text{reg } E^*) \xrightarrow{\cong} H_1(\text{reg } E^*)$ [7, Ch. 4, §30]. The Thom isomorphism being functorial under open embeddings [5, VIII.11.5], we have $t_X^{-1}([\delta_{Q'}]) = [\tau^{-1}(\delta_{Q'})] \in H_3(X^*, X^* \setminus E^*)$.

The divisors D_u^{red} and E^* in X^* are transversal, and so the 3-sphere $s_{Q'}$ is transversal to the divisor E^* along the 1-cycle $\delta_{Q'} \subseteq E^*$. Hence $s_{Q'} \cap U$ and $\tau^{-1}(\delta_{Q'})$ represent the same relative homology class in $H_3(U, U \setminus \text{reg } E^*) = H_3(X^{**}, X^{**} \setminus \text{reg } E^*)$. This proves the equality $\rho_X([s_{Q'}]) = [\delta_Q]$.

Therefore, we have shown that $\rho_X(H_3(X^*)) \subseteq H_C$, and every element $[\delta_Q]$ of a free base of the \mathbb{Z} -module H_C is the image of an element $[s_{Q'}] \in H_3(X^*)$ with $\sigma(Q') = Q$. This proves the first equality of the lemma; the proof of the second one is similar. \square

Proof of Proposition 2.19. In virtue of 2.21 and 2.22 we have $\pi_*(H_C) \subseteq H_\Gamma$ and $\hat{\pi}_*$ in (11) is, indeed, an isomorphism. Furthermore, as $H_1(\bar{\Gamma}) = H_1(\Gamma)/G(\Gamma^*)$ and $H_1(\overline{C^*}) = H_1(C)/G(C^*)$ (see 2.13), the homomorphism $\bar{\pi}_* : H_1(\overline{C^*}) \rightarrow H_1(\bar{\Gamma})$ factors through $\hat{\pi}_*$, whence $\bar{\pi}_*$ is a surjection. To show that it is also injective, consider a nonzero element $[\alpha] \in H_1(\overline{C^*})$ generated by a 1-cycle α in C_i^* (where $i \leq k$) such that the 1-cycle $\beta := \pi(\alpha)$ in Γ_i^* gives a zero element of $H_1(\bar{\Gamma}_i)$. Hence the class $[\beta] \in H_1(\Gamma_i^*)$ is contained in the subgroup $G(\Gamma_i^*)$. It is easily seen that $\pi_*(G(C_i^*))$ contains a subgroup of finite index in $G(\Gamma_i^*)$. Thus for some $m > 0$ we have $m[\beta] = \pi_*([\gamma])$ with $[\gamma] \in G(C_i^*)$. As $m[\alpha] - [\gamma]$ is still a nonzero element of $H_1(\overline{C^*})$ whose image in $H_1(\Gamma_i^*)$ is zero, we have $\hat{\pi}_*(m[\alpha] - [\gamma]) = 0$, which is wrong since $\hat{\pi}_*$ is an isomorphism. This proves that $\bar{\pi}_*$ in (12) is an isomorphism.

It follows that the decompositions (9) and (10) in 2.18 can be chosen in such a way that π_* respects them. Then by (11) and (12), $\tilde{\pi}_*$ in (13) is also an isomorphism. The proof is completed. \square

2.23. The proof of the next lemma is based on the following simple observation. Let G is a free abelian group of finite rank. For an element $a \in G$, the following conditions are equivalent:

- (i) a is *primitive* (that is, if $a = kb$ with $b \in G$, $k \in \mathbb{Z}$ then $k = \pm 1$);
- (ii) given a free base a_1, \dots, a_n of a \mathbb{Z} -module G , the coordinates in the presentation $a = \sum_{i=1}^n k_i a_i$ are relatively prime.

Lemma 2.24. *Let F be a curve with the irreducible components F_1, \dots, F_L , which are all non-compact, and with the multiple points P_1, \dots, P_M . Denote by p_1, \dots, p_N the punctures of the curve $F^* := F \setminus \{P_1, \dots, P_M\}$, and let B_i be a local branch of \bar{F} centered at p_i . Consider the group $G := G(F^*)/H_F$, where*

$$G(F^*) := \langle lk_{p_i}(B_i) \mid i = 1, \dots, N \rangle \subseteq H_1(F^*)$$

and

$$H_F := \langle lk_{P_j}(F) \mid j = 1, \dots, M \rangle \subseteq G(F^*).$$

Then $G \cong \mathbb{Z}^{N-M-L}$. Furthermore, for each $i = 1, \dots, N$ the class $[lk_{p_i}(B_i)] \in G$ is primitive unless it is zero. The latter holds if and only if p_i is the only puncture of the corresponding component F_l , and F_l is an isolated component of F .

Proof. Denoting $\nu : F^{\text{norm}} \rightarrow F$ a normalization, we let $\nu^{-1}(P_m) = \{p_j\}_{j \in J_m}$ with $J := \coprod_{m=1}^M J_m \subseteq \{1, \dots, N\}$ ⁹. Thus $lk_{P_m}(F) = \sum_{j \in J_m} lk_{p_j}(B_j)$. We also let $\{p_i\}_{i \in I_l}$ be the set of punctures of a component $F_l^* := F_l \cap F^*$ of the curve F^* . For every $m = 1, \dots, M$ (resp., $l = 1, \dots, L$) we pick an index $j_m \in J_m$ (resp., $i_l \in I_l \setminus J$) (notice that the index set $I_l \setminus J$ of the punctures of F_l is non-empty, as F_l is assumed to be non-compact). Then we have:

$$\begin{aligned} G(F^*) &= \langle lk_{p_i}(B_i) \mid i \notin \{i_1, \dots, i_L\} \rangle \\ &= \langle lk_{p_i}(B_i), lk_{P_m}(F) \mid m = 1, \dots, M, i \notin \{i_1, \dots, i_L, j_1, \dots, j_M\} \rangle. \end{aligned}$$

Thus $G(F^*) \cong \mathbb{Z}^{N-L} \cong H_F \oplus \mathbb{Z}^{N-L-M}$ with $H_F \cong \mathbb{Z}^M$, and so

$$(16) \quad G = G(F^*)/H_F = \langle [lk_{p_i}(B_i)] \mid i \notin \{i_1, \dots, i_L, j_1, \dots, j_M\} \rangle \cong \mathbb{Z}^{N-L-M}.$$

The elements $[lk_{p_i}(B_i)] \in G$ of the free base (16) satisfy the condition (ii) of 2.23, whence are primitive. The indices $j_m \in J_m$ being arbitrary, as $\text{card} J_m \geq 2$ the classes $[lk_{p_i}(B_i)] \in G$ with $i \in J$ are all primitive as well. By the same reason, a class $[lk_{p_i}(B_i)] \in G$ with $i \in I_l \setminus J$ is primitive unless $\{i\} = I_l \setminus J$ (i.e., unless p_i is the only puncture of F_l). If such a component F_l meets another one $F_{l'}$ at a multibranch point P_m , then choosing j_m from $J_m \cap I_{l'}$ and decomposing $[lk_{p_i}(B_i)] \in G$ in the base (16) we obtain that at least one of the coordinates equals -1 , hence 2.23(ii) is fulfilled, and so $[lk_{p_i}(B_i)]$ is primitive too. Finally, $[lk_{p_i}(B_i)] \in G$ is not primitive if and only if p_i is the only puncture of F_l and F_l is isolated (in that case $lk_{p_i}(B_i) + \sum_{P_m \in F_l} lk_{P_m}(F) = 0$ in $G(F^*)$, whence $[lk_{p_i}(B_i)] = 0$ in G). \square

In the proof of Theorem 2.11 below (based on 2.19 and 2.24) the role of F in 2.24 is played respectively by the curves Γ and $C_{\text{non-vert}} \setminus C_{\text{vert}}$. As in the proof of

⁹Here \coprod stands for the disjoint union.

2.24, it will be important to bear in mind the freedom of choice when selecting the indices $\{i_1, \dots, i_L, j_1, \dots, j_M\}$ as in (16). From 2.24 we obtain such a corollary.

Corollary 2.25. *In the notation as in 2.19, let $Q \in \bar{C}_i$ ($i \in \{1, \dots, k\}$) be a puncture of C_i^* such that $P := \bar{\pi}(Q) \in \bar{\Gamma}_i$ is not a puncture at infinity of the affine curve Γ_i . Then the following hold.*

- (a) $\text{mult}_Q \bar{\pi} = 1$ (that is, $\bar{\pi}$ is non-ramified at Q);
- (b) P is a puncture of Γ_i^* ;
- (c) the components C_i and Γ_i are horizontal (that is, $1 \leq i \leq h$).

Proof. The curve C_i^* possesses yet another puncture over a place at infinity of Γ_i , whence by (2.24), $[lk_Q(C_i^*)]$ is a primitive element of the group $G(C^*)/H_C$. As $\tilde{\pi}_*$ in (13) is an isomorphism, $\tilde{\pi}_*([lk_Q(C_i^*)]) = \kappa[lk_P(\Gamma_i^*)] \in G(\Gamma^*)/H_\Gamma \setminus \{0\}$ is primitive as well, whence $\kappa = \text{mult}_Q \bar{\pi} = 1$, which yields (a) and (b).

Suppose that there are two different local branches A and A' of \bar{C} over a local branch B of $\bar{\Gamma}$ at P , and let $Q, Q' \in \bar{C}$ be their centers. The same argument as above shows that $\tilde{\pi}_*([lk_Q(A)]) = \tilde{\pi}_*([lk_{Q'}(A')]) = [lk_P(B)]$, where both $[lk_Q(A)], [lk_{Q'}(A')] \in G(C^*)/H_C$ are primitive. Hence

$$(17) \quad [lk_Q(A)] = [lk_{Q'}(A')] \neq 0.$$

It is not difficult to verify that (in contradiction with (17)) there exists a base (16) which includes both $[lk_Q(A)]$ and $[lk_{Q'}(A')]$, unless A and A' are local branches of the curve C at a multibranch point $Q \in S_C^{(0)}$ with $\mu_Q(C) = 2$. But in the latter case $[lk_Q(A)] = -[lk_{Q'}(A')]$, which again contradicts (17). Therefore, there is only one local branch of \bar{C} over B , which yields (c). \square

Now we are ready to prove Theorem 2.11.

Proof of Theorem 2.11. It is convenient to proceed first with the proof of (β) . From 2.25 we get:

$$\Gamma_{\text{slant}} \cap S_\Gamma = \emptyset = C_{\text{slant}} \cap S_C$$

(indeed, Γ_{slant}^* resp., C_{slant}^* cannot have punctures other than the places at infinity). Thus the slanted components C_i and Γ_i ($h+1 \leq i \leq k$) of C resp., Γ are isolated and do not contain multibranch points. In particular, $C_i = C_i^*$ resp., $\Gamma_i = \Gamma_i^*$, and these curves are homeomorphic to their normalizations. Furthermore, the group $G(C_i^*)$ (resp., $G(\Gamma_i^*)$) is a direct summand of $G(C^*)/H_C$ (resp., $G(\Gamma^*)/H_\Gamma$). Hence by 2.19,

$$(18) \quad \tilde{\pi}_*|_{G(C_i^*)} : G(C_i^*) \xrightarrow{\cong} G(\Gamma_i^*)$$

is an isomorphism, as well as

$$(19) \quad \bar{\pi}_*|_{H_1(\bar{C}_i)} : H_1(\bar{C}_i) \xrightarrow{\cong} H_1(\bar{\Gamma}_i).$$

Since $\deg \bar{\pi}|_{\bar{C}_i} > 1$, (19) shows that $H_1(\bar{C}_i) = 0 = H_1(\bar{\Gamma}_i)$ i.e., both \bar{C}_i and $\bar{\Gamma}_i$ are rational curves. Therefore by (8) in 2.13,

$$H_1(C_i^{\text{norm}}) \cong H_1(C_i) = H_1(C_i^*) = G(C_i^*)$$

and

$$H_1(\Gamma_i^{\text{norm}}) \cong H_1(\Gamma_i) = H_1(\Gamma_i^*) = G(\Gamma_i^*).$$

Now (18) yields that

$$\pi^{\text{norm}*} : H_1(C_i^{\text{norm}}) \rightarrow H_1(\Gamma_i^{\text{norm}})$$

is an isomorphism. By a theorem of Hurwitz (see e.g., [18, I.2.1]), a morphism $\rho : F \rightarrow G$ of smooth irreducible affine curves is an isomorphism once the induced homomorphism $\rho_* : H_1(F) \rightarrow H_1(G)$ is an isomorphism, unless $H_1(F) = H_1(G) = 0$ i.e., $F \simeq G \simeq \mathbb{C}$. Thus $C_i^{\text{norm}} \simeq \Gamma_i^{\text{norm}} \simeq \mathbb{C}$, and so the curves C_i and Γ_i are homeomorphic to \mathbb{C} , which proves (β) .

(α) By 2.15 $\pi^{-1}(\Gamma_{\text{vert}}) \subseteq C_{\text{vert}}$, whence $\pi(C_{\text{horiz}} \setminus C_{\text{vert}}) \subseteq \Gamma_{\text{horiz}} \setminus \Gamma_{\text{vert}}$. To show that actually, the latter inclusion is an equality, suppose on the contrary that for a point $P \in \Gamma_{\text{horiz}} \setminus \Gamma_{\text{vert}}$, $\pi^{-1}(P) = \emptyset$. Then all $\mu := \mu_P(\Gamma)$ branches A_1, \dots, A_μ of \tilde{C} over the branches B_1, \dots, B_μ of Γ at P have centers at infinity. By 2.25(b) the curve Γ^* has a puncture over P . Thus $P \in S_\Gamma^{(0)}$ is a multibranch point of $\Gamma_{\text{horiz}} \setminus \Gamma_{\text{vert}}$, and so the primitive classes (see 2.24) $[lk_P(B_j)] \in G(\Gamma^*)/H_\Gamma$ ($j = 1 \dots, \mu$) are subjected to the (only) relation $\sum_{j=1}^\mu [lk_P(B_j)] = 0$. Therefore, the primitive classes $[lk_{Q_j}(A_j)] \in G(C^*)/H_C$ ($j = 1 \dots, \mu$) are also related by

$$(20) \quad \sum_{j=1}^\mu [lk_{Q_j}(A_j)] = 0,$$

where $Q_j \in \tilde{C}$ is the center of the branch A_j . Constructing a free base (16) of the free \mathbb{Z} -module $G(C^*)/H_C$, we may suppose that for any $j = 1, \dots, \mu$, $j \notin \{i_1, \dots, i_L\}$ (and definitely, $j \notin \{j_1, \dots, j_M\}$ as Q_j is a puncture at infinity of C). Thus the classes $[lk_{Q_j}(A_j)]$ ($j = 1 \dots, \mu$) make a part of a free base (16), and so cannot satisfy (20), a contradiction.

Denote $F := C_{\text{horiz}} \setminus C_{\text{non-horiz}} = C_{\text{horiz}} \setminus C_{\text{vert}}$ and $G := \Gamma_{\text{horiz}} \setminus \Gamma_{\text{non-horiz}} = \Gamma_{\text{horiz}} \setminus \Gamma_{\text{vert}}$. We have shown that the morphism $\pi|_F : F \rightarrow G$ of degree 1 is bijective. It follows that it is an isomorphism. Indeed, as D_g and D_f meet transversally along F (see 2.5), the curve F is the zero divisor of the restriction $g|_{G \times \mathbb{C}}$, which is a polynomial of degree one in z , say, $az + b \in \mathbb{C}[G][z]$. Note that a and b have no common zero on G (otherwise the zeros of $g|_{G \times \mathbb{C}} = az + b$ would contain a vertical component). As $\pi|_F$ is surjective, a is nowhere zero, whence $z = -b/a \in \mathbb{C}[G]$, and so $\pi|_F$ is an isomorphism. This proves (α) .

(γ) The group $G(\Gamma^*)/H_\Gamma = \tilde{\pi}_*(G(C^*)/H_C)$ being generated by the classes

$$\tilde{\pi}([lk_{Q_j}(C_i^*)]) \in G(\Gamma_{\text{non-vert}}^*)/[H_\Gamma \cap G(\Gamma_{\text{non-vert}}^*)]$$

(where Q_j runs over the set of punctures of the curve C^*) we have

$$G(\Gamma_{\text{non-vert}}^*)/[H_\Gamma \cap G(\Gamma_{\text{non-vert}}^*)] = G(\Gamma^*)/H_\Gamma.$$

Since in virtue of (β) , $G(\Gamma_{\text{slant}}^*) = 0$ we obtain

$$G(\Gamma^*)/H_\Gamma = G(\Gamma_{\text{non-slant}}^*)/H_\Gamma = G(\Gamma_{\text{horiz}}^*)/[H_\Gamma \cap G(\Gamma_{\text{horiz}}^*)].$$

In particular, for each puncture P of Γ_{vert}^* , we have

$$(21) \quad [lk_P(\bar{\Gamma})] \in G(\Gamma_{\text{horiz}}^*)/[H_\Gamma \cap G(\Gamma_{\text{horiz}}^*)].$$

It follows that on any vertical component of Γ there is only one puncture at infinity. Indeed, if there were a component Γ_i of Γ_{vert} with at least two punctures at infinity, say, P_1 and P_2 , then any free base as in (16) of the \mathbb{Z} -module $G(\Gamma^*)/H_\Gamma$ would contain at least one of the corresponding classes, say, $[lk_{P_1}(\bar{\Gamma})]$, which contradicts (21).

A similar argument shows that the curve Γ_{vert} has no selfintersection. In particular, the components of Γ_{vert} are disjoint, and for each $i = k+1, \dots, n$, Γ_i is homeomorphic to Γ_i^{norm} .

Furthermore, by 2.19

$$\bar{\pi}_* : H_1(\bar{C}^*) \rightarrow H_1(\bar{\Gamma}) = H_1(\bar{\Gamma}_{\text{non-vert}}) \oplus H_1(\bar{\Gamma}_{\text{vert}})$$

is an isomorphism, and $\bar{\pi}_*(H_1(\bar{C}^*)) \subseteq H_1(\bar{\Gamma}_{\text{non-vert}})$. It follows that $H_1(\bar{\Gamma}_{\text{vert}}) = 0$, whence the components of Γ_{vert} are rational. Finally, Γ_i ($i = k + 1, \dots, n$) is a rational curve with one place at infinity and without selfintersections, therefore is homeomorphic to \mathbb{C} .

As for any $i = h + 2, \dots, n$, $D_{h+1} \cap D_i = \emptyset$ and $D_i \simeq \Gamma_i \times \mathbb{C}$ is simply connected, with the notation as in 2.6 we have that the restriction $f_{h+1}|_{D_i}$ does not vanish, and so is constant: $f_{h+1}|_{D_i} =: \lambda_i \in \mathbb{C}$. Thus we obtain a decomposition as in (γ) , which completes the proof of (γ) .

(δ) As the matrix B' in 2.6 is unimodular and by (γ) the vertical components are disjoint, arguing as in 2.6 we can easily see that the morphism $\pi : C \rightarrow \Gamma$ maps any component of C_{vert} into a component of Γ_{vert} , and maps different components of C_{vert} into different components of Γ_{vert} . Moreover, the columns of B' are vectors of the standard basis in \mathbb{R}^{n-k} . Hence up to a reordering we may assume that B' is the unit matrix. The slanted components $\Gamma_{h+1}, \dots, \Gamma_k$ being isolated the last $k - h$ lines of the matrix B are zero, which completes the proof of (δ) . \square

Remark 2.26. Notice that for any vertical component C_i of C ($i = k + 1, \dots, n$), $P_i := \pi(C_i)$ is a smooth point of the curve Γ_i . This follows from 1.2 as $m_{ii} = 1$.

2.3. Preserving acyclicity: a criterion. We have the following partial converse to 2.11.

Theorem 2.27. *Let X and Y be irreducible smooth affine 3-folds which satisfy the condition (ii) of 2.4, as well as the following one (which replaces 2.4(iii)):*

- (iii') $Y = Z \times \mathbb{C}$ is a cylinder over a smooth acyclic affine surface Z , and $D = \Gamma \times \mathbb{C}$ with $\Gamma \subseteq Z$.

Suppose also that the conditions (α) - (δ) of Theorem 2.11 are fulfilled.¹⁰ Then the induced homomorphisms

$$\sigma_* : H_*(X) \rightarrow H_*(Y) \quad \text{and} \quad \sigma_* : \pi_1(X) \rightarrow \pi_1(Y)$$

are isomorphisms. Henceforth, the 3-fold X is acyclic; it is contractible if and only if Y is so.

Proof. The theorem immediately follows from 1.4, 1.10 and 2.32 below. Indeed, the assumption 2.11 (δ) allows to apply 1.4 in order to get that $\sigma_* : \pi_1(X) \rightarrow \pi_1(Y)$ is an isomorphism. In turn, by 2.32 the assumptions of 1.10 are fulfilled, and so by 1.10, $\sigma_* : H_*(X) \rightarrow H_*(Y)$ is an isomorphism as well. \square

From 2.11 and 2.27 we obtain the following criterion for preserving the acyclicity.

Corollary 2.28. *Let X and Y be smooth affine 3-folds satisfying the conditions 2.4(i) and 2.27(iii'). If Y is acyclic resp., contractible then X is so if and only if the conditions 2.11 (α) - (δ) are fulfilled.*

¹⁰See 2.29 below.

Remark 2.29. Assuming in 2.27 that the conditions 2.11(α)-(δ) are fulfilled we require implicitly that the things are as in 2.6-2.8, without supposing acyclicity as in 2.4(i). In fact, 2.6-2.8 refer only to the fact that the multiplicity matrix M_σ is unimodular, which is anyhow foreseen by 2.11(δ).

Actually the proofs of the lemmas below rely only on the conditions (ii), (iii) and the following one.

- (iv) There is a regular function $\varphi \in A = \mathbb{C}[Y]$ such that for every $i = h+1, \dots, n$ the restriction $\varphi|_{D_i}$ is constant, as well as the restriction of φ to any fiber of the morphism $\pi : D \rightarrow \Gamma$, and for any point $P \in S_\Gamma \setminus \Gamma_{\text{vert}}$, both $\Phi_P := \varphi^*(\varphi(\pi^{-1}(P))) \subseteq Y$ and $\Psi_P := \sigma^{-1}(\Phi_P) \subseteq X$ are smooth, reduced surfaces with $\sigma^*(\Phi_P) = \Psi_P$.

Under these assumptions σ_* is an isomorphism in homology (even if we do not suppose as in 2.27 that Y is acyclic).

The next lemma shows that the conditions (ii) and (iii') imply (iv).

Lemma 2.30. *Under the assumptions of 2.27 there exists a (possibly empty) smooth, reduced curve Γ' on Z such that*

- (a) $\Gamma' \cap \Gamma_{\text{non-horiz}} = \emptyset$ and $\Gamma' \supseteq S_\Gamma^{(0)} \setminus \Gamma_{\text{non-horiz}}$,
- (b) the surfaces $\Phi := \Gamma' \times \mathbb{C} \subseteq Y$ and $\Psi := \sigma^{-1}(\Phi) \subseteq X$ are smooth, and
- (c) $\sigma|_\Psi : H_*(\Psi) \rightarrow H_*(\Phi)$ is an isomorphism.

Proof. If $h < n$ (i.e., $\Gamma_{\text{non-horiz}} \neq \emptyset$) then we take for Γ' the union of the fibers of the regular function $f_{h+1}|_Z$ through the points of $S_\Gamma^{(0)} \setminus \Gamma_{\text{non-horiz}}$. Indeed, since the Euler characteristics $e(\Gamma_{h+1}) = e(Z) = 1$ and $\Gamma_{h+1} = (f_{h+1}|_Z)^*(0)$, by [38, 6.2] the fibers of $f_{h+1}|_Z$ are smooth, reduced and irreducible except for, possibly, Γ_{h+1} . Thus Γ' is smooth and reduced. As f_{h+1} is constant on any component Γ_i of $\Gamma_{\text{non-horiz}}$ we have $\Gamma' \cap \Gamma_{\text{non-horiz}} = \emptyset$, whence (a) is fulfilled.

In the case where $h = n$ (i.e., $\Gamma_{\text{non-horiz}} = \emptyset$) the existence of a smooth, reduced curve Γ' satisfying (a) easily follows by Bertini's Theorem.

Evidently, $\Phi = \Gamma' \times \mathbb{C}$ is a smooth surface. Since $\sigma|_{X \setminus E} : X \setminus E \rightarrow Y \setminus D$ is an isomorphism, the surface $\Psi := \sigma^{-1}(\Phi)$ is smooth if it is smooth at the points $R \in \Psi \cap E$. Let $Q = (P, z_0) := \sigma(R) \in C$ with $P \in \Gamma \cap \Gamma' \subseteq \Gamma_{\text{horiz}} \setminus \Gamma_{\text{non-horiz}}$. As Γ' is smooth we can find local coordinates (x, y) on Z centered at P such that (locally) $\Gamma' = \{x = 0\}$. Thus $Q = (P, z_0) = (0, 0, z_0)$, $R = (Q, u_0) = (0, 0, z_0, u_0)$, and (locally)

$$\Psi = \{x = 0\} \cap X = \{f(0, y)u - g(0, y, z) = 0\} \subseteq \mathbb{C}_{y,z,u}^3,$$

where by the condition 2.11(α), (locally) $g(0, y, z) = a(y)z + b(y)$ with $a(0) \neq 0$ (see the proof of 2.11(α)). Therefore, $\partial g / \partial z(Q) = a(0) \neq 0$, whence the surface Ψ is smooth at R . Thus (b) holds.

To show (c) we need the following claim.

Claim. $(\sigma|_\Psi)^*((D_{\text{horiz}} \cap \Phi)^{\text{red}}) = (E_{\text{horiz}} \cap \Psi)^{\text{red}}$.

Proof of the claim. We will use the same local chart and the notation as above. We have $(D_{\text{horiz}} \cap \Phi)^{\text{red}} = \{P\} \times \mathbb{C}_z = y^*(0)$ (y being regarded as a function on Φ). Then locally, $(y \circ \sigma) : (y, z, u) \mapsto y$, whence the subspace $\ker d(y \circ \sigma)|_R = \{y = 0\}$ does not contain the tangent space $T_R X$ (indeed as $a(0) \neq 0$ the differential $d(fu - g)|_R = \dots - a(0)dz$ of the defining polynomial of Ψ in $\mathbb{C}_{y,z,u}^3$ at the point R is

not proportional to the differential dy of $y = y \circ \sigma$ at R). It follows that $d(y \circ \sigma)|_{\Psi}$ does not vanish at the points $R \in \Psi \cap E$, and so locally

$$(E_{\text{horiz}} \cap \Psi)^{\text{red}} = (y \circ \sigma)^*(0) = (\sigma|_{\Psi})^*((D_{\text{horiz}} \cap \Phi)^{\text{red}}),$$

as desired. \square

Notice that $\Phi \cap D_{\text{horiz}} = (\Gamma' \cap \Gamma_{\text{horiz}}) \times \mathbb{C} \subseteq \Phi$ is a disjoint union of affine lines, whereas in virtue of 2.11(α), $\Phi \cap C_{\text{horiz}}$ is a finite set of points, one on every of those lines. Furthermore, $\Psi \cap E_{\text{horiz}} = (\Phi \cap C_{\text{horiz}}) \times \mathbb{C}$.

The proof of (c) is based on 1.9. In the notation as in 1.9 we let $\hat{X} := \Psi$, $\hat{Y} := \Phi$, $\hat{E} := \Psi \cap E_{\text{horiz}}$ and $\hat{D} := \Phi \cap D_{\text{horiz}}$. Then by (b) above, \hat{X} , \hat{Y} , \hat{E} and \hat{D} are smooth varieties, $\sigma(\hat{E}) \subseteq \hat{D}$, $\sigma(\hat{X} \setminus \hat{E}) = \hat{Y} \setminus \hat{D}$, and (by the above claim) $\sigma^*(\hat{D}) = \hat{E}$, that is, the condition (i) of 1.9 is fulfilled. As $\sigma|_{\hat{X} \setminus \hat{E}} : \hat{X} \setminus \hat{E} \rightarrow \hat{Y} \setminus \hat{D}$ is an isomorphism, taking into account the observations following the claim it is easily seen that 1.9(ii) holds as well. Thus by 1.9, $\sigma_* : H_*(\hat{X}) \rightarrow H_*(\hat{Y})$ is an isomorphism, which yields (c). \square

2.31. The proof of the next lemma relies on 1.10, where we let $\hat{E} = E' \cup E''$ and $\hat{D} = D' \cup D''$ with $E' := E_{\text{horiz}}$, $D' := D_{\text{horiz}}$,

$$E'' := E_{\text{non-horiz}} \coprod \Psi \quad \text{and} \quad D'' := D_{\text{non-horiz}} \coprod \Phi,$$

Φ and Ψ being the same as in 2.30(b).

Lemma 2.32. *Under the assumptions and the notation as in 2.27 and 2.31, the conditions (i') and (ii') of 1.10 are fulfilled.*

Proof. In virtue of 2.11(γ), for every $i = h + 1, \dots, n$ both curves C_i and Γ_i are homeomorphic to \mathbb{C} , whence both surfaces $E_i = C_i \times \mathbb{C}$ and $D_i \simeq \Gamma_i \times \mathbb{C}$ are homeomorphic to \mathbb{C}^2 . Therefore $E_{\text{non-horiz}}$ and $D_{\text{non-horiz}}$ are topological manifolds, and $(\sigma|_{E_{\text{non-horiz}}})_* : H_*(E_{\text{non-horiz}}) \rightarrow H_*(D_{\text{non-horiz}})$ is an isomorphism. As the surfaces Φ and Ψ are smooth we have that E'' and D'' are topological manifolds as well, and in view of 2.30(c), $(\sigma|_{E''})_* : H_*(E'') \rightarrow H_*(D'')$ is an isomorphism.

As by our construction

$$S_{\Gamma} \subseteq \Gamma' \cup \Gamma_{\text{non-horiz}} \quad \text{and} \quad S_C \subseteq \pi^{-1}(\Gamma') \cup C_{\text{non-horiz}},$$

the curves $\Gamma_{\text{horiz}} \setminus (\Gamma' \cup \Gamma_{\text{non-horiz}})$ and $C_{\text{horiz}} \setminus (\pi^{-1}(\Gamma') \cup C_{\text{non-horiz}})$ have no multi-branch points. Therefore they are topological manifolds, as well as the surfaces

$$D' \setminus D'' = [\Gamma_{\text{horiz}} \setminus (\Gamma' \cup \Gamma_{\text{non-horiz}})] \times \mathbb{C} \quad \text{and} \quad E' \setminus E'' = [C_{\text{horiz}} \setminus (\pi^{-1}(\Gamma') \cup C_{\text{non-horiz}})] \times \mathbb{C}.$$

By 2.11(α) the projection

$$\pi : C_{\text{horiz}} \setminus (\pi^{-1}(\Gamma') \cup C_{\text{non-horiz}}) \rightarrow \Gamma_{\text{horiz}} \setminus (\Gamma' \cup \Gamma_{\text{non-horiz}})$$

is an isomorphism, whence $\sigma|_{E' \setminus E''} : E' \setminus E'' \rightarrow D' \setminus D''$ is so as well. Now the conditions (i') and (ii') of 1.10 follow. \square

3. SIMPLE AFFINE MODIFICATIONS OF \mathbb{C}^3 DIFFEOMORPHIC TO \mathbb{R}^6

The main result of this section is 3.6, which together with 2.28 provides a criterion for as when a simple affine modification $X \subseteq \mathbb{C}^4$ of $Y = \mathbb{C}^3$ is isomorphic to \mathbb{C}^3 .

3.1. Exotic simple modifications of \mathbb{C}^3 . We keep the terminology and the notation of section 2, and we adopt the following

Convention 3.1. Hereafter

- (i) $Y = \mathbb{C}^3$ with the coordinates x, y, z , and
- (ii) X is a smooth affine 3-fold in \mathbb{C}^4 diffeomorphic to \mathbb{R}^6 , with equation of the form

$$(22) \quad p = f(x, y)u + g(x, y, z) = 0,$$

where $f \in \mathbb{C}[x, y] \setminus \mathbb{C}$, $g \in \mathbb{C}[x, y, z]$ (thus by 2.28, the conditions 2.11(a)-(δ) hold).

3.2. Note that the blowup morphism $\sigma : X \ni (x, y, z, u) \mapsto (x, y, z) \in Y$ represents X as a simple affine modification of $Y = \mathbb{C}^3$ along the cylindrical divisor $D = D_f^{\text{red}} = \Gamma \times \mathbb{C}$ (where $\Gamma := f^{-1}(0) \subset \mathbb{C}^2$) with center $C = \{f = g = 0\} \subset \mathbb{C}^3$ and with the morphism $\pi : C \rightarrow \Gamma$ as in 2.5 given by $\pi : (x, y, z) \mapsto (x, y)$. Hence the assumptions (i)-(iii) of 2.4 are fulfilled.

3.3. As in 2.8 we factorize $f \in \mathbb{C}[x, y]$ into irreducible factors: $f = \prod_{i=1}^n f_i^{a_i}$, and we write g as a polynomial in z :

$$g(x, y, z) = \sum_{j=0}^d b_j(x, y)z^j \quad \text{with} \quad d := \deg_z g.$$

We let $d_i := \deg(\pi|_{C_i})$. Recall (2.6, 2.7) that an irreducible component C_i (resp., $\Gamma_i = f_i^{-1}(0)$) of the curve C (resp., Γ) is vertical (resp., horizontal resp., slanted) if and only if $d_i = 0$ (resp., $d_i = 1$ resp., $d_i \geq 2$). The following lemma is a simple observation, and so we omit the proof.

Lemma 3.4. *For every $i = 1, \dots, n$ we have $b_j \in (f_i) \forall j = d_i + 1, \dots, d$ and $b_{d_i} \notin (f_i)$. Hence p as in (22) admits a unique presentation*

$$p = fu + g_i + f_i h_i$$

with $g_i(x, y, z) := \sum_{j=0}^{d_i} b_j(x, y)z^j \in \mathbb{C}[x, y, z]$ and $h_i \in (z^{d_i+1}) \subseteq \mathbb{C}[x, y, z]$. Furthermore, if $f_{\text{horiz}} \neq \text{const}$ then p can be written as

$$(23) \quad p = f_{\text{horiz}} f_{\text{non-horiz}} u + b_0 + b_1 z + f_{\text{horiz}}^{\text{red}} h_0$$

with $b_0, b_1 \in \mathbb{C}[x, y]$ and $h_0 \in (z^2) \subseteq \mathbb{C}[x, y, z]$, where $b_1|_{\Gamma_{\text{horiz}} \setminus \Gamma_{\text{non-horiz}}}$ has no zero.

3.5. Recall [39] that an *exotic* \mathbb{C}^3 is a smooth affine 3-fold diffeomorphic to \mathbb{R}^6 but non-isomorphic to \mathbb{C}^3 .

The principal result of this subsection is the following theorem.

Theorem 3.6. *If under the assumptions as in 3.1, at least one of the curves $C_{\text{non-horiz}}$ and $\Gamma_{\text{non-horiz}}$ is singular then X is an exotic \mathbb{C}^3 .*

The proof is done in 3.9 and 3.15 below. For a converse result, see 3.21 in the next subsection.

Notice that 3.6 provides a regular way of constructing exotic \mathbb{C}^3 -s as hypersurfaces in \mathbb{C}^4 . Let us give concrete examples.

3.7. Examples. For the Russell cubic 3-fold

$$X = \{x^2u + x + y^2 + z^3 = 0\} \subset \mathbb{C}^4,$$

the curve $\Gamma = \Gamma_{\text{slant}} = \{x = 0\} \subset \mathbb{C}^2$ is smooth and isomorphic to \mathbb{C} , whereas $C = C_{\text{slant}} = \{x = y^2 + z^3 = 0\} \subset \mathbb{C}^3$ (the center of modification) is homeomorphic to \mathbb{C} but singular. It is well known [19, 4, 39] that X represents an exotic algebraic structure on \mathbb{C}^3 . By [13] the Koras-Russell cubic 3-fold

$$X = \{(x^2 + y^3)u + x + z^2 = 0\} \subset \mathbb{C}^4$$

also is an exotic \mathbb{C}^3 ; here both $\Gamma = \Gamma_{\text{slant}} = \{x^2 + y^3 = 0\} \subset \mathbb{C}^2$ and $C = C_{\text{slant}} = \{x^2 + y^3 = 0 = x + z^2\} \subset \mathbb{C}^3$ are homeomorphic to \mathbb{C} but singular.

Remark 3.8. Generalizing a theorem of Sathaye [30], in [15, Thm. 7.2] it is proven that actually, every smooth acyclic surface in \mathbb{C}^3 with equation $p = f(x, y)u + g(x, y) = 0$ (where $f, g \in \mathbb{C}^2$) is isomorphic to \mathbb{C}^2 and rectifiable. Examples 3.7 show that in general, this does not hold anymore in \mathbb{C}^4 without the additional assumption of smoothness of $C_{\text{non-horiz}}$ and $\Gamma_{\text{non-horiz}}$ (cf. 3.6 above).

The proof of 3.6 starts with the following proposition (cf. 3.20 below).

Proposition 3.9. *Let X and Y be as in 3.1. If the curve $\Gamma_{\text{non-horiz}}$ is singular then the Derksen invariant $\text{Dk}(A')$ of the algebra $A' := \mathbb{C}[X]$ is non-trivial, whence $X \not\cong \mathbb{C}^3$.*

The proof is done in 3.10-3.14.

Lemma 3.10. *Under the assumptions as in 3.9, choosing appropriate new coordinates in the (x, y) -plane and rescaling the z -coordinate we may write the polynomial p in the form*

$$(24) \quad p = (x^k - y^l)^m f_{\text{horiz}}(x, y)u + z^e + g_0(x, y, z) + (x^k - y^l)h_0(x, y, z),$$

where $k, l, e \geq 2$, $(k, l) = 1$, $f_{\text{horiz}} \in \mathbb{C}[x, y]$ and $f_{\text{horiz}}(0, 0) = 1$, $g_0, h_0 \in \mathbb{C}[x, y, z]$, $\deg_z g_0 < e$ and $z^e | h_0$.

Proof. In the notation as in 2.6-2.9 we have $h < n$. In virtue of the condition 2.11(γ) we may suppose that the component Γ_{h+1} of $\Gamma_{\text{non-horiz}}$ is a singular plane curve homeomorphic to \mathbb{C} . Hence by the Lin-Zaidenberg Theorem [18], choosing new coordinates in the (x, y) -plane we may assume that $f_{h+1} = x^k - y^l$ with $k, l \geq 2$ and $(k, l) = 1$. As the other fibers $x^k - y^l = c$ ($c \in \mathbb{C} \setminus \{0\}$) of the polynomial f_{h+1} are not homeomorphic to \mathbb{C} , in view of 2.11(γ) we have $h + 1 = n$, and so $f_{\text{non-horiz}} = f_n^m = (x^k - y^l)^m$ with $m := a_n$. By 3.4 the polynomial (22) can be now written as follows:

$$p = (x^k - y^l)^m f_{\text{horiz}}(x, y)u + b_e(x, y)z^e + g_0(x, y, z) + (x^k - y^l)h_0(x, y, z)$$

with $b_e \in \mathbb{C}[x, y]$, $g_0, h_0 \in \mathbb{C}[x, y, z]$, $\deg_z g_0 < e$ and $z^{e+1} | h_0$. In virtue of 2.11(δ) and 2.5 the divisors $D_n = \Gamma_n \times \mathbb{C}$ and D_g^{red} meet transversally at general points of $C_n = D_n \cap D_g^{\text{red}}$. If Γ_n were vertical (i.e., $g(x, y, z) \equiv b_0(x, y) \pmod{(x^k - y^l)}$; see 3.3, 3.4) then the equation $b_0(t^l, t^k) = 0$ would have a unique solution t_0 such that (in virtue of 2.26) $\pi(C_n) = (t_0^l, t_0^k)$ is a smooth point of $\Gamma_n \subseteq \mathbb{C}_{x, y}^2$ i.e., $t_0 \neq 0$. Thus $b_0(t^l, t^k) = c(t - t_0)^r$ which is wrong as the derivative of the left hand side vanishes at $t = 0$. Therefore, $\Gamma_n = \Gamma_{\text{slant}}$, whence $e = \deg \pi|_{C_n} \geq 2$

and $\pi|_{C_n} : C_n \rightarrow \Gamma_n$ is a proper morphism (as each of these curves has only one puncture). It follows that the restriction $b_e|_{\Gamma_n}$ has no zero. As Γ_n is simply connected we have $b_e|_{\Gamma_n} = \text{const} =: b_e^0 \in \mathbb{C}^*$, and so $b_e(x, y) = b_e^0 + (x^k - y^l)c_e(x, y)$ for a certain polynomial $c_e \in \mathbb{C}^{[2]}$. Finally, rescaling the z -coordinate if necessary, we obtain the desired presentation. As by 2.11(δ), $\Gamma_n = \Gamma_{\text{slant}}$ is an isolated component of Γ , the restriction $f_{\text{horiz}}|_{\Gamma_n}$ does not vanish, whence is constant, and we may assume in addition that this constant is $f_{\text{horiz}}(0, 0) = 1$. \square

3.11. We choose the weight degree function d on the algebra A with

$$d_x = -lN, \quad d_y = -kN, \quad d_z = \sqrt{2} \quad \text{and} \quad d_u = klmN + e\sqrt{2},$$

so that the polynomials

$$f_n = f_{h+1} = x^k - y^l \quad \text{and} \quad \hat{p} := (x^k - y^l)^m u + z^e$$

are d -quasihomogeneous. Letting $N \in \mathbb{N}$ to be sufficiently large (thus $d(h_0) < kln$, and hence $d((x^k - y^l)h_0) < 0$) we may assume that \hat{p} as above is the d -principal part of the polynomial p as in (24).

Lemma 3.12. *Let $q \in \mathbb{C}[x, y, z, u] \setminus \mathbb{C}$ be an irreducible d -homogeneous polynomial with $\deg_z q < e$. Then q coincides (up to a constant factor) with one of the following polynomials:*

$$x, y, z, u, \lambda x^k + \mu y^l, \quad \text{where} \quad \lambda, \mu \in \mathbb{C}^*.$$

Proof. Letting $q = \sum_{i=0}^{e-1} a_i(x, y, u)z^i$ with $a_i \in \mathbb{C}[x, y, u]$ ($i = 0, \dots, e-1$) we claim that there can be only one nonzero coefficient a_i . Assuming on the contrary that $a_i, a_{i+j} \neq 0$ for some $i, i+j \in \{0, \dots, e-1\}$ with $1 \leq j \leq e-1$ we would have $d(a_i z^i) = d(a_{i+j} z^{i+j})$. Taking into account the equality $d_y = \frac{k}{l}d_x$ we derive

$$\begin{aligned} d(a_i z^i) - d(a_{i+j} z^{i+j}) &= j d_z = j\sqrt{2} = \alpha d_u + \beta d_x = \alpha e\sqrt{2} + \beta' d_x \\ (25) \quad &\Rightarrow (j - \alpha e)\sqrt{2} = \beta' d_x \in \mathbb{Q} \end{aligned}$$

for certain $\alpha \in \mathbb{Z}_{\geq 0}$ and $\beta, \beta' \in \mathbb{Q}$. But $\alpha e - j \in \mathbb{Q} \setminus \{0\}$, whence (25) leads to a contradiction.

Therefore, the irreducible polynomial q must coincide (up to a constant factor) either with z or with a polynomial $a_0 \in \mathbb{C}[x, y, u]$. Let $q = a_0(x, y, u) = \sum_{i=0}^t c_i(x, y)u^i$ with $c_i \in \mathbb{C}[x, y]$ ($i = 0, \dots, t$). The weights d_x (resp., d_y) and d_u being independent over \mathbb{Q} the same argument as above shows that at most one summand $c_i(x, y)u^i$ may be different from zero.

Thus once again, the irreducible polynomial q coincides (up to a constant factor) either with u or with a polynomial $c_0 \in \mathbb{C}[x, y]$. In the latter case, being d -homogeneous the polynomial q must coincide (up to a constant factor) with one of the polynomials $x, y, \lambda x^k + \mu y^l$ ($\lambda, \mu \in \mathbb{C}^*$), as stated. \square

3.13. Let an irreducible polynomial $p \in \mathbb{C}^{[4]}$ be as in (24). Set as before $X = p^{-1}(0) \subset \mathbb{C}^4$, $\hat{X} = \hat{p}^{-1}(0) \subset \mathbb{C}^4$ and $\hat{A} = \mathbb{C}[\hat{X}]$ where \hat{p} is the d -principal part of p . Denote by $\rho_{\hat{A}}$ the canonical surjection

$$\rho_{\hat{A}} : \mathbb{C}^{[4]} \rightarrow \mathbb{C}^{[4]}/(\hat{p}) = \hat{A}'$$

and by $\hat{x} = \rho_{\hat{A}}(x), \dots, \hat{u} = \rho_{\hat{A}}(u) \in \hat{A}'$ the traces on \hat{X} of the coordinate functions.

The polynomial \hat{p} being irreducible, by 1.22(a) \hat{A}' is just the graded algebra associated with the filtered algebra A' , a filtration being provided by the degree function d on A' .

Lemma 3.14. *In the notation as in 3.13 we have*

$$\text{ML}_{\text{gr}}(\hat{A}') = \text{Dk}_{\text{gr}}(\hat{A}') = \mathbb{C}[\hat{x}, \hat{y}] \subset \hat{A}'_{\leq 0}.$$

Therefore (by 1.24)

$$\text{Dk}(A') \subset A'_0 \neq A',$$

so that $A' \not\cong \mathbb{C}^3$ and $X \not\cong \mathbb{C}^3$, which proves 3.9.

Proof. The Jacobian derivation

$$\partial_0 := \frac{\partial(\hat{p}, x, y, *)}{\partial(x, y, z, u)}$$

on the algebra \hat{A}' being homogeneous and locally nilpotent with $\hat{A}'^{\partial_0} = \ker \partial_0 = \mathbb{C}[\hat{x}, \hat{y}]$, it is sufficient to show that $\hat{x}, \hat{y} \in \ker \hat{\partial}$ for any $\hat{\partial} \in \text{LND}_{\text{gr}}(\hat{A}')$. Indeed, by 1.14 both $\mathbb{C}[\hat{x}, \hat{y}]$ and $\ker \hat{\partial}$ are algebraically closed subalgebras of \hat{A}' of transcendence degree 2, thereby they coincide provided that $\mathbb{C}[\hat{x}, \hat{y}] \subset \ker \hat{\partial}$.

The derivation $\hat{\partial}$ being homogeneous, the subalgebra $\ker \hat{\partial} \subset \hat{A}'$ is generated by homogeneous elements (i.e., by the restrictions to \hat{X} of d -homogeneous polynomials on \mathbb{C}^4). Let $a = q|_{\hat{X}} \in \ker \hat{\partial}$ (with a d -homogeneous $q \in \mathbb{C}^4$) be nonconstant. We may assume that $\deg_z q < e$ (otherwise we replace the polynomial q by the rest of the Euclidean division of q by the z -monic d -homogeneous polynomial $\hat{p} = z^e + (x^k - y^l)^m u$).

The kernel $\hat{A}'^{\hat{\partial}} = \ker \hat{\partial}$ being factorially closed (see 1.14(c)), the irreducible factors of q restricted to \hat{X} belong to this kernel as well. Therefore, $\ker \hat{\partial}$ is generated by the traces of irreducible d -homogeneous polynomials q with $\deg_z q < e$. By 1.14(d) we have $\hat{x}, \hat{y} \in \ker \hat{\partial}$ provided that $\lambda \hat{x}^k + \mu \hat{y}^l \in \ker \hat{\partial}$ for some $\lambda, \mu \in \mathbb{C}^*$. Due to 3.12 and to the above argument on algebraic closeness, $\ker \hat{\partial}$ coincides with the subalgebra of \hat{A}' generated by one of the following pairs:

$$(\hat{x}, \hat{y}), \quad (\hat{x}, \hat{u}), \quad (\hat{y}, \hat{u}), \quad (\hat{x}, \hat{z}), \quad (\hat{y}, \hat{z}), \quad (\hat{z}, \hat{u}).$$

If $\hat{z} \in \ker \hat{\partial}$ then by 1.14 and the equality

$$(26) \quad \hat{z}^e + (\hat{x}^k + \hat{y}^l)^m \hat{u} = 0$$

also $\hat{x}, \hat{y}, \hat{u} \in \ker \hat{\partial}$, whence $\hat{\partial} = 0$, a contradiction. This eliminates the last three cases.

If $\ker \hat{\partial} = \mathbb{C}[\hat{x}, \hat{u}]$ ¹¹ then (26) yields the equality

$$(27) \quad lm \deg_{\hat{\delta}} \hat{y} = e \deg_{\hat{\delta}} \hat{z}.$$

On the other hand, as $\ker \hat{\partial} = \mathbb{C}[\hat{x}, \hat{u}]$, by 1.19 $\hat{\partial}$ is equivalent to the Jacobian derivation

$$\hat{\partial}_1 := \frac{\partial(\hat{p}, x, u, *)}{\partial(x, y, z, u)}.$$

It is easily seen that

$$\hat{\partial}_1(\hat{y}) = -(\hat{p})'_z|_{\hat{X}} = -\hat{z}^{e-1},$$

¹¹The case where $\ker \hat{\partial} = \mathbb{C}[\hat{y}, \hat{u}]$ can be eliminated by a similar argument.

and so as $\deg_{\partial} = \deg_{\partial_1}$ (see 1.18) we get

$$(28) \quad \deg_{\hat{\delta}} \hat{y} = (e - 1) \deg_{\hat{\delta}} \hat{z} + 1.$$

From (27) and (28) we obtain:

$$\left[\left(1 - \frac{1}{lm}\right)e - 1 \right] \deg_{\hat{\delta}} \hat{z} = -1.$$

But this is impossible because $l, e \geq 2, m \geq 1$, and so $\left(1 - \frac{1}{lm}\right)e \geq 1$. This completes the proof of the lemma, as well as those of 3.9. \square

Next we consider the remaining possibility in 3.6.

Proposition 3.15. *Let X and Y be as in 3.1. Suppose that $\Gamma_{\text{non-horiz}}$ is a smooth curve, whereas there exists a singular component, say, C_{h+1} of the curve C_{slant} . Then once again, the Derksen invariant $\text{Dk}(A')$ is non-trivial, whence $X \not\cong \mathbb{C}^3$.*

The proof is done in 3.16-3.19 below.

Lemma 3.16. *Under the assumptions as in 3.15, after applying an appropriate tame automorphism of $\mathbb{C}_{x,y,z}^3$ the polynomial p can be presented in the form*

$$(29) \quad p = x^m h_0(x, y, z)u + y^k + z^l + xh_1(x, y, z)$$

with $k, l, m \geq 2, (k, l) = 1, h_0, h_1 \in \mathbb{C}[x, y, z]$ and $h_0(0, y, z) = \text{const} = 1$.

Proof. The curve $\Gamma_{h+1} \subset \mathbb{C}_{x,y}^2$ (being smooth and homeomorphic to \mathbb{C}) is isomorphic to \mathbb{C} . Thus by the Abhyankar-Moh-Suzuki Theorem, choosing appropriate new coordinates in the (x, y) -plane we may suppose that $f_{h+1} = x$. We factorize $f = f_{h+1}^m \tilde{f} = x^m \tilde{f}$ with $m := a_{h+1}$. As C_{h+1} is singular, by 2.3 the gradient $\text{grad} f$ has to vanish at some point of $\Gamma_{h+1} = \{x = 0\}$. The component Γ_{h+1} of Γ being isolated (see 2.11(β)) this implies that $m \geq 2$. Furthermore the polynomial $\tilde{f}(0, y)$ does not vanish, whence is a non-zero constant; by rescaling the x -coordinate we may assume this constant being 1.

As the curve $C_{h+1} = \{x = 0 = g(0, y, z)\} \subseteq \mathbb{C}_{y,z}^2$ is homeomorphic to \mathbb{C} and singular, by the Lin-Zaidenberg Theorem (after performing an appropriate automorphism of the plane $\mathbb{C}_{y,z}^2$) we may suppose that C_{h+1} is given by $x = y^k + z^l = 0$ with $k, l \geq 2$ and $(k, l) = 1$. In virtue of 2.5 and the condition 2.11(δ) the divisors $D_{h+1} = \{x = 0\} \subset \mathbb{C}_{x,y,z}^3$ and D_g^{red} meet transversally at general points of the curve C_{h+1} , whence $g(0, y, z) = y^k + z^l$ (up to a constant factor which can be put equal to 1) i.e., $g = y^k + z^l + xh_1(x, y, z)$. Taking for $h_0(x, y, z)$ the polynomial obtained from \tilde{f} as a result of the latter coordinate change, we obtain the desired presentation (29) with $h_0(0, y, z) = \tilde{f}(0, y) = 1$. \square

3.17. We consider the weight degree function d on the algebra $A' := \mathbb{C}[X]$ with

$$d_x = -1, \quad d_y = d_z = 0 \quad \text{and} \quad d_u = m.$$

As the d -principal part $\hat{p} = x^m u + y^k + z^l$ of the polynomial p as in (29) is irreducible, by 1.22 $\hat{A}' := \mathbb{C}[\hat{X}]$ is just the graded algebra associated with the filtration on A' defined by the degree function d . Notice that in \hat{A}' the following relation holds:

$$(30) \quad \hat{x}^m \hat{u} + \hat{y}^k + \hat{z}^l = 0.$$

With this notation we have the following lemma.

Lemma 3.18. $\text{Dk}_{\text{gr}}(\hat{A}') \subset \hat{A}'_{\leq 0}$, and so (by 1.24) $\text{Dk}(A') \subset A'_0 \neq A'$. Thereby $A' \not\subset \mathbb{C}^{[3]}$ and $X \not\subset \mathbb{C}^3$, which yields 3.15.

Proof. We must show that $\ker \hat{\partial} \subset \hat{A}'_{\leq 0}$ whenever $\hat{\partial} \in \text{LND}_{\text{gr}}(\hat{A}')$. Assume that there exist $\hat{\partial} \in \text{LND}_{\text{gr}}(\hat{A}')$ and $\hat{a} \in \ker \hat{\partial}$ such that $\hat{a} \notin \hat{A}'_{\leq 0}$. Furthermore, by 1.14(c) we may suppose that this element \hat{a} is non-decomposable. We have $\hat{a} = q|_{\hat{X}}$ for some d -homogeneous polynomial $q = \sum_{i,j} a_{ij}x^i u^j \in \mathbb{C}^{[4]}$ (with $a_{ij} \in \mathbb{C}[y, z]$). Moreover, taking into account (30) we may suppose that $i < m$ whenever $j > 0$.

Claim. *In the above expression for q there is only one non-zero monomial.*

Proof of the claim. Indeed, if there were two of them, say, if $a_{i_1 j_1} \neq 0$ and $a_{i_2 j_2} \neq 0$ then we would have

$$(31) \quad d(q) = m j_1 - i_1 = m j_2 - i_2 \quad \implies \quad m(j_1 - j_2) = i_1 - i_2.$$

Assuming that $i_1 > i_2$, by (31) we get $j_1 > j_2$, and vice versa. Thus $j_1 > 0$, and then by our assumption $i_1 < m$. Hence also $i_1 - i_2 < m$, which contradicts (31). This proves the claim. \square

Since the element $\hat{a} = q(\hat{x}, \hat{y}, \hat{z}, \hat{u}) = a_{ij}(\hat{y}, \hat{z})\hat{x}^i \hat{u}^j$ is supposed to be non-decomposable and of positive d -degree, we have $\hat{a} = \hat{u} \in \ker \hat{\partial}$. Thus $\hat{\partial}$ can be specified to a locally nilpotent derivation ∂_1 of the algebra $B = \mathbb{C}[S]$, where $S := \{x^m + y^k + z^l = 0\} = \{u = 1\} \cap \hat{X} \subset \mathbb{C}^3$ (see 1.16).

By 1.14(a), $\text{tr.deg}(\ker \hat{\partial}) = 2$, whence there is a homogeneous $\hat{\partial}$ -constant \hat{b} such that the elements $\hat{a} = \hat{u}, \hat{b} \in \hat{A}'$ are algebraically independent. As above we obtain that either $\hat{b} = b(\hat{y}, \hat{z})$ for some irreducible polynomial $b \in \mathbb{C}^{[2]} \setminus \mathbb{C}$, or $\hat{b} = \hat{x}$. In the latter case by (30) we have $\hat{y}^k + \hat{z}^l \in \ker \hat{\partial}$, and thus by 1.14(d) also $\hat{y}, \hat{z} \in \ker \hat{\partial}$. Therefore $\hat{\partial} = 0$, which is impossible. Finally, we conclude that $b(\hat{y}, \hat{z}) \in \ker \hat{\partial}$ for a certain polynomial $b \in \mathbb{C}^{[2]} \setminus \mathbb{C}$, and so the restriction $b|_S$ is a ∂_1 -constant.

Now the proof can be completed by applying the next lemma (cf. [4, 16, 39]). \square

Lemma 3.19. *Let $B = \mathbb{C}[S]$ where $S := \{x^m + y^k + z^l = 0\} \subset \mathbb{C}^3$ and $k, l, m \geq 2$, $\text{gcd}(k, l) = 1$, and let $\partial_1 \in \text{LND}(B)$. Then $b|_S \notin \ker \partial_1$ whenever $b \in \mathbb{C}[y, z] \setminus \mathbb{C}$.*

Proof. We define the weight degree function \hat{d} on the algebra B by letting

$$\hat{d}_x = kl, \quad \hat{d}_y = lm, \quad \hat{d}_z = km.$$

Actually $x^m + y^k + z^l$ is a \hat{d} -homogeneous polynomial, B is a graded algebra, and we may consider the associated homogeneous derivation $\hat{\partial}_1 \in \text{LND}_{\text{gr}}(B)$. Assuming that for a polynomial $b \in \mathbb{C}[y, z] \setminus \mathbb{C}$, $b|_S \in \ker \partial_1$ we will get a \hat{d} -homogeneous polynomial $\hat{b}_1 \in \mathbb{C}[y, z] \setminus \mathbb{C}$ such that $\hat{b}_1|_S \in \ker \hat{\partial}_1$ as well. By 1.14(c) an irreducible \hat{d} -homogeneous factor of the polynomial \hat{b}_1 (this can be y, z or $\lambda y^k + \mu z^l$, where $\lambda, \mu \in \mathbb{C}^*$) restricts to S as a $\hat{\partial}_1$ -constant. If it were $\lambda y^k + \mu z^l$ then by 1.14(d) both $y|_S$ and $z|_S$ would be $\hat{\partial}_1$ -constants, whence $\hat{\partial}_1 = 0$, a contradiction.

Thus we may assume that, say, $y|_S \in \ker \hat{\partial}_1$. As $(x^m + y^k + z^l)|_S = 0$ we have $(x^m + z^l)|_S \in \ker \hat{\partial}_1$, and (since $m, l \geq 2$) again by 1.14(d) we obtain that $x|_S, z|_S \in \ker \hat{\partial}_1$, which is impossible.

This completes the proof of 3.19, 3.18 and 3.15. \square

Remark 3.20. Choosing a weight degree function likewise in 3.11 and repeating the proof of Proposition 5.3 in [12] it can be shown that under assumptions as in 3.6 (that is, as in 3.9 or 3.15) one has $\mathbb{C}[x] \subseteq \text{ML}(A')$, whence the Makar-Limanov invariant $\text{ML}(A')$ is non-trivial as well.

3.2. Simple affine modifications of \mathbb{C}^3 isomorphic to \mathbb{C}^3 . The following theorem is a central result of the paper. The expression ‘in appropriate (x, y) -coordinates’ below means ‘after performing an appropriate automorphism $\alpha \in \text{Aut } \mathbb{C}[x, y] \subset \text{Aut } \mathbb{C}[x, y, z, u]$ ’. (In fact, the new (x, y) -coordinates are chosen in a way to get $f_{\text{non-horiz}} \in \mathbb{C}[x]$; in particular, no coordinate change is necessary when $f_{\text{non-horiz}} = \text{const.}$)

Theorem 3.21. *For a hypersurface $X \subset \mathbb{C}^4$ given by an equation*

$$p = f(x, y)u + g(x, y, z) = 0 \quad (f \in \mathbb{C}^{[2]} \setminus \{0\}, g \in \mathbb{C}^{[3]}),$$

in appropriately chosen new (x, y) -coordinates the following conditions (i)-(vi) are equivalent.

- (i) $X \simeq \mathbb{C}^3$.
- (ii) *Every fiber $X_\mu = p^{-1}(\mu)$ ($\mu \in \mathbb{C}$) of the polynomial $p \in \mathbb{C}^{[4]}$ is isomorphic to \mathbb{C}^3 .*
- (iii) *Every fiber of the regular function $x|_X \in A' := \mathbb{C}[X]$ is reduced and isomorphic to \mathbb{C}^2 .*
- (iv) *Every fiber of the morphism $\rho : \mathbb{C}^4 \rightarrow \mathbb{C}^2$, $(x, y, z, u) \mapsto (x, p(x, y, z, u))$, is reduced and isomorphic to \mathbb{C}^2 .*
- (v) *The polynomial p is a residual x -variable ¹².*
- (vi) *With $Y := \mathbb{C}^3$ and $X \subseteq \mathbb{C}^4$ as above being smooth and irreducible, the conditions 2.11(α)-(δ) hold as well as the following one:*
- (ε) *The curves $C_{\text{non-horiz}}$ and $\Gamma_{\text{non-horiz}}$ are smooth.*

Proof. The implications (v) \Rightarrow (iv) \Rightarrow (iii) and (ii) \Rightarrow (i) are immediate; (i) \Rightarrow (vi) follows from 2.28 and 3.6. Thus it suffices to establish (vi) \Rightarrow (v) and (iii) \Rightarrow (ii).

Let us show (vi) \Rightarrow (v). If $f_{\text{non-horiz}} \neq \text{const}$ (that is, $h < n$) then in virtue of our assumptions 2.11(γ) and (vi)(ε), $\Gamma_{h+1} \simeq \mathbb{C}$. By the Abhyankar-Moh-Suzuki Theorem, after performing an appropriate automorphism of the plane $\mathbb{C}_{x,y}^2$ we may suppose that $f_{h+1} = x$. Hence by 2.11(γ) we have

$$f_{\text{non-horiz}} = c \prod_{i=h+1}^n (x - \lambda_i)^{a_i} \in \mathbb{C}[x].$$

Anyhow, $f_{\text{non-horiz}} = \text{const}$ or not, assuming that $f_{\text{horiz}} \neq \text{const}$ by 3.4 we may write

$$p_\lambda = \kappa f_{\text{horiz}, \lambda} u + b_{0, \lambda} + b_{1, \lambda} z + f_{\text{horiz}, \lambda}^{\text{red}} h_\lambda$$

with $f_{\text{horiz}, \lambda}, f_{\text{horiz}, \lambda}^{\text{red}}, b_{0, \lambda}, b_{1, \lambda} \in B := \mathbb{C}[y]$, $h_\lambda \in B[z]$ and $\kappa := f_{\text{non-horiz}}(\lambda) = c \prod_{i=h+1}^n (\lambda - \lambda_i)^{a_i} \in \mathbb{C}$. If $\lambda \in \mathbb{C} \setminus \{\lambda_{h+1}, \dots, \lambda_n\}$ (i.e., $\kappa \neq 0$) then in virtue of 3.4, $b_{1, \lambda} \in B$ is invertible mod $f_{\text{horiz}, \lambda}$, whereas $f_{\text{horiz}, \lambda}^{\text{red}}$ is nilpotent mod $f_{\text{horiz}, \lambda}$. Therefore by 1.29, $p_\lambda \in B[z, u]$ is a B -variable; in particular, $p_\lambda \in \mathbb{C}[y, z, u]$ is a variable (whence for every $\mu \in \mathbb{C}$, the surface $\{p_\lambda = \mu\} \subseteq \mathbb{C}_{y,z,u}^3$ is reduced and isomorphic to \mathbb{C}^2).

¹²That is (1.27) for any $\lambda \in \mathbb{C}$, the specialization $p_\lambda \in \mathbb{C}[y, z, u]$ of p is a variable.

For every $i = h+1, \dots, n$ we have $f_i = x - \lambda_i$ and in virtue of 2.11(γ) and (vi)(ε), $\Gamma_i \simeq \mathbb{C} \simeq C_i$. Moreover by 2.3(i) the polynomial $g_{\lambda_i} \in \mathbb{C}[y, z]$ is irreducible and $g_{\lambda_i}^{-1}(0) = C_i \simeq \mathbb{C}$, whence by the Abhyankar-Moh-Suzuki Theorem it is a variable of the polynomial ring $\mathbb{C}[y, z]$. Thus $p_{\lambda_i} = g_{\lambda_i} \in \mathbb{C}[y, z, u]$ is a variable as well. Now (v) follows. In the remaining case where $f_{\text{horiz}} = \text{const}$ the proof is similar (but simpler) and left to the reader.

Next we prove (iii) \Rightarrow (ii). We notice first of all that under the condition (iii), the hypersurface X is smooth and irreducible (indeed, for every $\lambda \in \mathbb{C}$ we have

$$\text{grad } p_\lambda = \text{pr}((\text{grad } p)_\lambda) \quad \text{with} \quad \text{pr} : (x, y, z, u) \mapsto (y, z, u).$$

Similarly, under the condition (iv) every fiber X_μ ($\mu \in \mathbb{C}$) is smooth and irreducible.

Actually we establish (iii) \Rightarrow (i), which at the same time shows (iv) \Rightarrow (ii). Together with the implication (i) \Rightarrow (iv) (which we already know) these yield (iii) \Rightarrow (ii), as needed.

To prove the implication (iii) \Rightarrow (i), in virtue of 1.7 it is enough to show that a smooth, irreducible 3-fold X satisfying (iii) is acyclic. Let $U = \mathbb{C} \setminus V$ be a Zariski open subset such that over U , the morphism $x|_X : X \rightarrow \mathbb{C}$ is a smooth fibration (with the fibers diffeomorphic to \mathbb{R}^4). In the notation of 1.9 we let $\hat{X} = X$, $\hat{Y} = Y = \mathbb{C}^3$, $\hat{E} = (x|_X)^{-1}(V) \subseteq \hat{X}$ and $\hat{D} = x^{-1}(V) \subseteq \hat{Y}$. It is easily seen that the induced homomorphisms

$$(x|_{\hat{X} \setminus \hat{E}})_* : H_*(\hat{X} \setminus \hat{E}) \rightarrow H_*(\mathbb{C} \setminus V),$$

$$(x|_{\hat{Y} \setminus \hat{D}})_* : H_*(\hat{Y} \setminus \hat{D}) \rightarrow H_*(\mathbb{C} \setminus V),$$

and hence also

$$(\sigma|_{\hat{X} \setminus \hat{E}})_* : H_*(\hat{X} \setminus \hat{E}) \rightarrow H_*(\hat{Y} \setminus \hat{D})$$

are isomorphisms, as well as

$$(\sigma|_{\hat{E}})_* : H_*(\hat{E}) \rightarrow H_*(\hat{D})$$

(indeed, $H_0(\hat{E}) \cong \mathbb{Z}^{\text{card } V} \cong H_0(\hat{D})$ and the senior homology are trivial). Thus the conditions (i) and (ii) of 1.9 are fulfilled, and so by 1.9

$$\sigma_* : H_*(\hat{X}) \rightarrow H_*(\hat{Y})$$

is an isomorphism. Therefore, the 3-fold $X = \hat{X}$ is acyclic. This completes the proof. \square

Remark 3.22. For another proof of the implication (iii) \Rightarrow (i) which do not use 1.7 see 4.23, 4.24 below.

3.23. If the polynomial $p \in \mathbb{C}^4$ is linear with respect to two (and not just one) variables we have a much stronger result (see 3.24 below). It is an analog in dimension 4 of Theorem 7.2 in [15] generalizing Sathaye's Theorem [30]. For a polynomial $\varphi \in \mathbb{C}[x, y]$ we let $\Gamma_\varphi := V(\varphi) \subseteq \mathbb{C}^2$. Also, we use below the variable v instead of z to emphasize the symmetry of the situation.

Theorem 3.24. *For a hypersurface $X \subset \mathbb{C}^4$ given by an equation*

$$(32) \quad p = \tilde{a}(x, y)u + \tilde{b}(x, y)v + c(x, y) = d(au + bv) + c = 0$$

with $a, b, c, d \in \mathbb{C}[x, y]$, $d \neq 0$, $(a, b) \neq (0, 0)$ and $\text{gcd}(a, b) = 1$, in appropriately chosen new (x, y) -coordinates the following conditions (i')-(v') are equivalent.

- (i') *The 3-fold X is irreducible, smooth and acyclic.*
- (ii') $X \simeq \mathbb{C}^3$.
- (iii') *The polynomial $p \in \mathbb{C}^4$ is a residual x -variable.*
- (iv') $p \in \mathbb{C}[x]^{[3]}$ *is an x -variable.*
- (v') $d \in \mathbb{C}[x]$, $\gcd(c, d) = 1$, $\Gamma_a \cap \Gamma_b \subseteq \Gamma_d$, *and for every root $x = x_i$ of d ($i = 1, \dots, \deg d$) we have*

$$(33) \quad c_{x_i}(y) = c(x_i, y) \in \mathbb{C}^* y + \mathbb{C}.$$

Proof. The strategy of the proof is as follows. In virtue of 3.21 the conditions (ii') and (iii') are equivalent to each other and to the other conditions of 3.21. The equivalence (iii') \iff (iv') will be established later on in 4.2(b). By 2.11 we have (i') \implies 2.11(α)-(δ). We show below that (i') also implies the condition (ε) of 3.21(vi). In virtue of 3.21 this yields the implications (i') \implies 3.21(vi) \iff (ii') \implies (i'), and so gives (i') \iff (ii'). Finally we show (iii') \iff (v'), which concludes the proof.

Interchanging (if necessary) the roles of u and v we may suppose in the sequel that $a \neq 0$.

(i') \implies (ε). We observe (see 3.3) that for p as in (32), $C_{\text{slant}} = \Gamma_{\text{slant}} = \emptyset$, whence $C_{\text{non-horiz}} = C_{\text{vert}}$ is smooth. And it was shown in the proof of 3.10 that if $\Gamma_{\text{non-horiz}}$ is singular then $\Gamma_{\text{non-horiz}} = \Gamma_{\text{slant}}$. Henceforth in our setting $\Gamma_{\text{non-horiz}}$ is also smooth, that shows (ε).

(iii') \implies (v'). As X is irreducible we have $\gcd(c, d) = 1$. Letting $q := p - c$ we fix a point $P = (x_0, y_0)$ such that

$$q_P = d(x_0, y_0)(a(x_0, y_0)u + b(x_0, y_0)v) = 0.$$

It is easily seen that $y - y_0$ divides $p_{x_0} - c(x_0, y_0) = q_{x_0} + (c_{x_0} - c(x_0, y_0))$. As $p_{x_0} \in \text{Var}_{\mathbb{C}}[y, u, v]$ we obtain that $q_{x_0} = 0$ and $p_{x_0} - c(x_0, y_0) = c_{x_0} - c(x_0, y_0) = \kappa(y - y_0)$ with $\kappa \in \mathbb{C}^*$, as needed in (33). As $\gcd(a, b) = 1$, from $q_{x_0} = 0$ we get $d_{x_0} = 0$. Thus for every root (x_0, y_0) of $d = 0$ we have $d_{x_0} = 0$, which means that $d \in \mathbb{C}[x]$. Moreover, for every root (x_0, y_0) of $a = b = 0$ we also have $d_{x_0} = 0$, which shows the inclusion $\Gamma_a \cap \Gamma_b \subseteq \Gamma_d$ and gives (v').

(v') \implies (iii'). In view of (33) for every root x_i of d ($i = h + 1, \dots, n$), $p_{x_i} = c_{x_i} \in \text{Var}_{\mathbb{C}}[y] \subseteq \text{Var}_{\mathbb{C}}[y, u, v]$. Let further $\lambda \in \mathbb{C}$ and $d(\lambda) \neq 0$. As by (v'), $\Gamma_a \cap \Gamma_b \subseteq \Gamma_d$ we have $\gcd(a_\lambda, b_\lambda) = 1$, whence $r_\lambda a_\lambda - s_\lambda b_\lambda = 1$ for certain polynomials $r_\lambda, s_\lambda \in \mathbb{C}[y]$. As the matrix

$$A_\lambda = \begin{pmatrix} a_\lambda & b_\lambda \\ s_\lambda & r_\lambda \end{pmatrix}$$

is invertible over the ring $\mathbb{C}[y]$, the mapping

$$\alpha : (u, v) \mapsto (u_1, v_1) := A_\lambda(u, v)$$

induces a $\mathbb{C}[y]$ -automorphism of $\mathbb{C}[y][u, v]$ with $\alpha(p_\lambda) = d(\lambda)u_1 + c_\lambda \in \text{Var}_{\mathbb{C}}[y, u, v]$. Thus (iii') follows. \square

4. SIMPLE BIRATIONAL EXTENSIONS OF $\mathbb{C}^{[3]}$ AS VARIABLES IN $\mathbb{C}^{[4]}$

4.1. Partial positive results. We recall the problem stated in the Introduction.

4.1. Problem. *Is it true that if a hypersurface $X \subseteq \mathbb{C}^4$ with equation of the form*

$$p := f(x, y)u + g(x, y, z) = 0 \quad (\text{where } f \in \mathbb{C}^{[2]} \setminus \{0\} \text{ and } g \in \mathbb{C}^{[3]})$$

is isomorphic to \mathbb{C}^3 then necessarily $p \in \text{Var}_{\mathbb{C}}\mathbb{C}^{[4]}$ and moreover, $p \in \text{Var}_B B^{[3]}$ (that is, p is an x -variable) provided in addition that $f_{\text{non-horiz}} \in B := \mathbb{C}[x]$?

The principal results of this subsection can be summarized as follows.

Theorem 4.2. *The answer to 4.1 is positive in each of the following cases:*

- (a) $f \in \mathbb{C}[x]$.
- (b) $\deg_z g \leq 1$.
- (c) $f_{\text{vert}} = 1$.
- (d) $f_{\text{horiz}} = f_{\text{horiz}}^{\text{red}}$.

The proof is given in 4.15 below. From 4.2(c),(d) we obtain the following corollary.

Corollary 4.3. *If in 4.1, $f = f_1^{a_1}$ with $f_1 \in \mathbb{C}[x, y]$ irreducible then p is a variable.*

4.4. By 3.21(v) we may suppose that p as in 4.1 is a residual x -variable (see 1.25, 1.27). Thus 4.1 would be answered in positive if the following conjecture¹³ were true.

Conjecture 4.5. *If $p \in \mathbb{C}^{[n]}$ is a residual x_1 -variable (that is,*

$$p_\lambda := p(\lambda, x_2, \dots, x_n) \in \text{Var}_{\mathbb{C}}\mathbb{C}^{[n-1]} \quad \forall \lambda \in \mathbb{C}$$

then p is an x_1 -variable: $p \in \text{Var}_{\mathbb{C}[x_1]}\mathbb{C}[x_1]^{[n-1]}$.

In 4.6-4.8 below we analyze the situation with the only assumption that $p = fu + g$ is a residual x -variable. First in 4.6 we include the case where $f = 0$, then in 4.7 we deal with the special case where $f \in \mathbb{C}[x] \setminus \{0\}$, whereas the general case is treated in 4.8.

Proposition 4.6. *Let X be a smooth affine surface and $\varphi : X \rightarrow \mathbb{C}$ be a morphism with $\varphi^*(\lambda) \simeq \mathbb{C} \quad \forall \lambda \in \mathbb{C}$. Then the following hold.*

- (a) $X \simeq \mathbb{C}^2$.
- (b) *Furthermore, if $X = \text{spec}(\mathbb{C}^3/(g))$ with $g \in \mathbb{C}^3 = \mathbb{C}[x, y, z]$ and $\varphi = x|_X$ then g is an x -variable.*
- (c) *Consequently, a residual x -variable $g \in \mathbb{C}[x, y, z]$ is an x -variable.*¹⁴

Proof. (a) The fact is well-known (e.g., cf. [26, Thm. 2.2.1]); nevertheless we indicate the proof. First extending $\varphi : X \rightarrow \mathbb{C}$ to a projective morphism $\bar{\varphi} : V \rightarrow \mathbb{P}^1$ of a smooth ruled surface V , we then pass to a smooth relatively minimal model of $(V, \bar{\varphi})$ by contracting successively the superfluous components of the reducible fibers of $\bar{\varphi}$ different from the closures of the original fibers of φ (this is always possible, see e.g. [10, Lemma 7], [38, Lemma 3.5] or [26, Lemma 1.4.1(6)]). Thus we obtain (as a completion of the original family) a Hirzebruch surface $\pi : \Sigma_n \rightarrow \mathbb{P}^1$ with a section $s : \mathbb{P}^1 \rightarrow \Sigma_n$ ‘at infinity’, so that $X = \Sigma_n \setminus (s(\mathbb{P}^1) \cup F_\infty)$. By means of elementary transformations over the point $\infty \in \mathbb{P}^1$ we replace Σ_n with $\Sigma_0 = \mathbb{P}^1 \times \mathbb{P}^1$ and $s(\mathbb{P}^1)$ with a constant section, say, C_0 . This yields the desired isomorphism $X = \Sigma_0 \setminus (C_0 \cup F_\infty) \simeq \mathbb{C}^2$.

¹³Proposed by the second author.

¹⁴Moreover, $g \in \mathbb{C}[x, y, z]$ is a residual x -variable of the ring $\mathbb{C}[x][y, z, u]$ iff it is an x -variable. Indeed, this follows from (c) and the cancellation for curves: $\Gamma \times \mathbb{C} \simeq \mathbb{C}^2 \implies \Gamma \simeq \mathbb{C}$.

(b) By (a) there is an isomorphism

$$\gamma : \mathbb{C}[x, y, z]/(g) \rightarrow \mathbb{C}^{[2]} = \mathbb{C}[x, y].$$

Letting $h := \gamma(x) \in \mathbb{C}^{[2]}$, by our assumption we obtain:

$$\mathbb{C}^{[2]}/(h) = \mathbb{C}[x, y, z]/(g, x) = \mathbb{C}[\varphi^*(0)] \cong \mathbb{C}^{[1]}.$$

Hence by the Abhyankar-Moh-Suzuki Theorem, up to an automorphism of $\mathbb{C}[x, y]$ we may suppose that $h = x$ i.e.,

$$\gamma : \mathbb{C}[x][y, z]/(g) \rightarrow \mathbb{C}[x]^{[1]}.$$

Now it follows from the generalized version of the Abhyankar-Moh-Suzuki Theorem [29, Thm. 2.6.2] that g is an x -variable, as stated.

(c) immediately follows from (a) and (b). \square

Proposition 4.7. *If $p = f(x)u + g(x, y, z) \in \mathbb{C}^{[4]}$ (with $f \in \mathbb{C}[x] \setminus \{0\}$) is a residual x -variable then actually it is an x -variable.*

Proof. As $u + g(x, y, z)$ is an x -variable, by 1.30(a) so is $p = fu + g$. \square

Proposition 4.8. *If $p = f(x, y)u + g(x, y, z) \in \mathbb{C}[x][y, z, u]$ with $f \notin \mathbb{C}[x]$ is a residual x -variable then p can be written as follows:*

$$(34) \quad p = q[r\tilde{f}u + \tilde{g}_0y^2 + \tilde{g}_1z + \tilde{f}^{\text{red}}\tilde{g}_2z^2] + a_0 + a_1y,$$

where

- $q, r, a_0, a_1 \in \mathbb{C}[x]$, $\tilde{f}, \tilde{g}_0, \tilde{g}_1 \in \mathbb{C}[x, y]$, $\tilde{g}_2 \in \mathbb{C}[x, y, z]$,
- $\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0)$ is a finite subset of $q^{-1}(0) \times \mathbb{C}_y$, and
- for every root x_0 of r , $q(x_0)\tilde{f}(x_0, y) \in \mathbb{C}$.

The proof is done in 4.11 below.

Corollary 4.9. *If $p = f(x, y)u + g(x, y, z) \in \mathbb{C}[x][y, z, u]$ is a residual x -variable then it is a $\mathbb{C}(x)$ -variable of $\mathbb{C}(x)^{[3]}$.*

Proof. As a $\mathbb{C}[x]$ -variable is also a $\mathbb{C}(x)$ -variable, by 4.6 and 4.7 we may suppose that $f \notin \mathbb{C}[x]$, and so 4.8 applies. Let p be presented as in (34). As $\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0)$ is finite, the polynomials $qr\tilde{f}$, $q\tilde{g}_1 \in \mathbb{C}[x, y]$ regarded as elements of $B := \mathbb{C}(x)[y]$ are coprime. Moreover the polynomial $\tilde{f}^{\text{red}}\tilde{g}_2z^2 \in B[z]$ is nilpotent mod $(qr\tilde{f})$, and so by 1.29(a), $p = fu + g \in \text{Var}_B B[z, u]$. \square

The proof of 4.8 relies on the following lemma (cf. [30, 15, 35]).

Lemma 4.10. *Let $p = f(y)u + g(y, z)$ be a variable of $\mathbb{C}[y, z, u]$.*

(a) *If $f \neq 0$ then p is an y -variable, and it can be written as follows:*

$$p = f(y)u + g_0(y) + g_1(y)z + f^{\text{red}}(y)\tilde{g}_2(y, z)z^2$$

with $\text{gcd}(f, g_1) = 1$.

(b) *If $f = 0$ then the coefficient of the highest order term in z of p (which a priori is a polynomial in y) is constant, unless $\deg_z g \leq 0$.*

Proof. (a) The statement is evidently true if $f = \text{const}$. Suppose that $f \notin \mathbb{C}$. By our assumption $\mathbb{C}^3/(p) \cong \mathbb{C}^2$. If $y_1 \in \mathbb{C}$ is such that $f(y_1) = 1$ then we have

$$\mathbb{C}[y, z, u]/(f(y)u + g(y, z), y - y_1) \cong \mathbb{C}[z, u]/(u + g_{y_1}(z)) \cong \mathbb{C}^1.$$

By the Abhyankar-Moh-Suzuki Theorem, the same is true for every value of y . In particular, for any root y_0 of f we have

$$\mathbb{C}[y, z, u]/(f(y)u + g(y, z), y - y_0) \cong \mathbb{C}[z, u]/(g_{y_0}(z)) \cong \mathbb{C}^1.$$

Therefore $\deg_z g_{y_0}(z) = 1$, $p = fu + g$ has the desired form, and by 1.29(a) (with $B := \mathbb{C}[y]$) it is an y -variable.

(b) In the course of the proof of 1.30 we have noticed that the polynomials of $\mathbb{C}[y, z]$ which are variables of $\mathbb{C}[y, z, u]$, are actually variables of $\mathbb{C}[y, z]$. Thus $g \in \text{Var}_{\mathbb{C}}\mathbb{C}[y, z]$, and so the statement (b) is well-known (e.g., see [1]). \square

4.11. Proof of Proposition 4.8. It is easily seen that the polynomial $p = fu + g \in \mathbb{C}[x][y, z, u]$ admits a presentation

$$(35) \quad p = q[\bar{f}u + \tilde{g}_0y^2 + \tilde{g}_1z + \tilde{g}_2z^2] + a_0 + a_1y$$

with $a_0, a_1 \in \mathbb{C}[x]$, $\bar{f}, \tilde{g}_0, \tilde{g}_1 \in \mathbb{C}[x, y]$, $\tilde{g}_2 \in \mathbb{C}[x, y, z]$ and with $q \in \mathbb{C}[x]$ being a monic polynomial of maximal degree such that (35) holds. Clearly,

$$q^{-1}(0) = L(p) := \{x_0 \in \mathbb{C} \mid p_{x_0} = f(x_0, y)u + g(x_0, y, z) \in \mathbb{C}y + \mathbb{C}\}$$

(possibly, this set is empty, and then we put $q := 1$). As by our assumption, $f \notin \mathbb{C}[x]$ the subset

$$K = K(f) := \{\lambda \in \mathbb{C} \mid f_\lambda(y) \in \mathbb{C}\} = q^{-1}(0) \cup \{\lambda \in \mathbb{C} \mid \bar{f}_\lambda(y) \in \mathbb{C}\} \subseteq \mathbb{C}$$

is finite. We let $x_0 \in \mathbb{C} \setminus K(f)$. The specialization p_{x_0} being a variable of $\mathbb{C}[y, z, u]$, by 4.10(a) we have $\gcd(\bar{f}(x_0, y), \tilde{g}_1(x_0, y)) = 1$, whence $\bar{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0) \subseteq K \times \mathbb{C}_y$.

Since by 4.10(a), $p_{x_0} \in \mathbb{C}[y, z, u]$ is a residual y -variable, the equality $\bar{f}(x_0, y_0) = 0$ implies that $\deg_z g_{x_0, y_0}(z) = 1$, whence also $\tilde{g}_2(x_0, y_0, z) = 0$. It follows that

$$\bar{f}^{-1}(0) \setminus (K \times \mathbb{C}_y) \subseteq \tilde{g}_2^{-1}(0).$$

It is easily seen that $\bar{f} \in \mathbb{C}[x, y]$ admits a factorization $\bar{f}(x, y) = r(x)\tilde{f}(x, y)$ such that $r^{-1}(0) \subseteq K$ and $\tilde{f}^{-1}(0) \cap (K \times \mathbb{C}_y)$ is finite. Then we have

$$\tilde{f}^{-1}(0) \setminus (K \times \mathbb{C}_y) \subseteq \tilde{g}_2^{-1}(0).$$

Passing to the Zariski closures we obtain

$$\tilde{f}^{-1}(0) \subseteq \tilde{g}_2^{-1}(0) \quad \text{i.e.,} \quad \tilde{g}_2 = \tilde{f}^{\text{red}}\tilde{g}_2.$$

Now (35) yields a desired presentation (34). As $r^{-1}(0) \subseteq K(f)$, by the definition of $K(f)$ for every root x_0 of r we have $q(x_0)\tilde{f}(x_0, y) \in \mathbb{C}$, and so it remains to check that $\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0)$ is a finite subset of $q^{-1}(0) \times \mathbb{C}_y$.

We already know that $\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0)$ is a finite subset of $K \times \mathbb{C}_y$. Suppose that there is a point

$$(x_0, y_0) \in [\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0)] \setminus q^{-1}(0).$$

Then $x_0 \in K(f) \setminus q^{-1}(0)$, whence $\bar{f}(x_0, y) \in \mathbb{C}$, and so

$$\bar{f}(x_0, y) = \tilde{f}(x_0, y_0) = r(x_0)\tilde{f}(x_0, y_0) = 0.$$

By our assumption,

$$p_{x_0} = q(x_0)[\tilde{g}_0(x_0, y)y^2 + \tilde{g}_1(x_0, y)z + \tilde{f}^{\text{red}}(x_0, y)\tilde{g}_2(x_0, y, z)z^2] + a_0(x_0) + a_1(x_0)y$$

is a variable of $\mathbb{C}[y, z, u]$, and thus also of $\mathbb{C}[y, z]$. Hence by 4.10(b), $\tilde{f}^{\text{red}}(x_0, y) = \tilde{f}^{\text{red}}(x_0, y_0) = 0$, and then again by 4.10(b), $\tilde{g}_1(x_0, y) = \tilde{g}_1(x_0, y_0) = 0$. Therefore, $p_{x_0} \in \mathbb{C}[y]$ is a variable, whence $q(x_0) = 0$, a contradiction. \square

Remark 4.12. In the notation of 4.8, if $p = fu + g = qrfu + g$ is a residual x -variable then also $qfu + g$ is so. Moreover by 1.30(a), if $qfu + g$ is an x -variable then $p = fu + g$ is so as well.

Lemma 4.13. *Let $p = fu + g \in \mathbb{C}[x][y, z, u]$ be a residual x -variable presented as in (34). If $\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0) = \emptyset$ then p is an x -variable.*

Proof. By 4.12 we may put in (34) $r = 1$. In virtue of 1.29(a), the polynomial

$$\tilde{p} := \tilde{f}(x, y)u + \tilde{g}_0(x, y)y^2 + \tilde{g}_1(x, y)z + \tilde{f}^{\text{red}}(x, y)\tilde{g}_2(x, y, z)z^2$$

is an (x, y) -variable i.e., there is an automorphism $\alpha \in \text{Aut}_B B[z, u]$ (with $B := \mathbb{C}[x, y]$) such that $\alpha(u) = \tilde{p}$ (indeed by our assumption, \tilde{g}_1 is invertible mod \tilde{f} in B). Furthermore by our assumption, for every root x_0 of q the specialization

$$p_{x_0} = f(x_0, y)u + g(x_0, y, z) = a_0(x_0) + a_1(x_0)y \in \mathbb{C}[y, z, u]$$

is a variable. Consequently, $a_1(x_0) \neq 0$, and so $\gcd(q, a_1) = 1$. It follows that there is an affine automorphism $\beta \in \text{Aut}_{\mathbb{C}[x]} \mathbb{C}[x][y, u]$ such that

$$\beta(y) = q(x)u + a_0(x) + a_1(x)y.$$

Now $\alpha\beta \in \text{Aut}_{\mathbb{C}[x]} \mathbb{C}[x][y, z, u]$ is such that

$$\alpha\beta(y) = q\tilde{p} + a_0(x) + a_1(x)y = fu + g = p,$$

as desired. \square

Lemma 4.14. *If $p = f(x, y)u + g(x, y, z)$ is a residual x -variable of degree at most 1 in z then it is an x -variable.*

Proof. It follows from 4.8 that if $f \notin \mathbb{C}[x]$ then in (34) $\tilde{g}_1 \neq 0$ (indeed, otherwise $\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0)$ cannot be a finite subset of $q^{-1}(0) \times \mathbb{C}_y$). Thus if $\deg_z g \leq 0$ then $f \in \mathbb{C}[x]$, and so the statement follows from 4.7.

If $\deg_z g = 1$ then by 4.8 $p = fu + g$ can be written as follows:

$$p = q(x)[r(x)\tilde{f}(x, y)u + \tilde{g}_0(x, y)y^2 + \tilde{g}_1(x, y)z] + a_0(x) + a_1(x)y$$

with $\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0)$ being a finite subset of $q^{-1}(0) \times \mathbb{C}_y$. By 4.12 we may assume that $r = 1$.

We proceed by induction on the intersection index $n := \tilde{f}^*(0) \cdot \tilde{g}_1^*(0)$. If $n = 0$ then the statement follows from 4.13. Suppose that $n \geq 1$.

Let $x_0 \in q^{-1}(0)$ be a coordinate of an intersection point $(x_0, y_0) \in \tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0)$. Up to a translation we may suppose that $x_0 = 0$. As

$$\min\{\deg_y \tilde{g}_1(0, y), \deg_y \tilde{f}(0, y)\} > 0$$

we may write

$$\tilde{g}_1(0, y)z + \tilde{f}(0, y)u = d(y)(a(y)z + b(y)u),$$

where $\gcd(a, b) = 1$. Hence there is a linear automorphism $\gamma \in \text{Aut}_B B[z, u]$ (with $B := \mathbb{C}[x, y]$) such that $\gamma(a(y)z + b(y)u) = z$. It is not difficult to verify that applying γ amounts to:

$$\text{Jac}(\gamma) \begin{bmatrix} \tilde{g}_1 \\ \tilde{f} \end{bmatrix} = \begin{bmatrix} \hat{g}_1 \\ x\hat{f} \end{bmatrix}$$

for certain $\hat{f}, \hat{g} \in B$ (where $\text{Jac}(\gamma)$ is the Jacobi matrix of γ), and hence

$$\gamma(p) = \gamma(fu + g) = q(x)[x\hat{f}(x, y)u + \tilde{g}_0(x, y)y^2 + \hat{g}_1(x, y)z] + a_0(x) + a_1(x)y.$$

In view of the equality of ideals:

$$(\hat{g}_1, x\hat{f}) = (\tilde{g}_1, \tilde{f}),$$

for the intersection indices we have:

$$\hat{g}_1^*(0) \cdot (x\hat{f})^*(0) = \tilde{g}_1^*(0) \cdot \tilde{f}^*(0) = n,$$

therefore

$$\hat{g}_1^*(0) \cdot \hat{f}^*(0) \leq n - 1.$$

Now in virtue of 4.12,

$$\hat{p} := q(x)[\hat{f}(x, y)u + \tilde{g}_0(x, y)y^2 + \hat{g}_1(x, y)z] + a_0(x) + a_1(x)y$$

is a residual x -variable. Hence by the inductive hypothesis, it is also an x -variable. Again by 4.12, so are $\gamma(p)$ and $p = fu + g$ as well. The induction step is done, so the proof is completed. \square

4.15. Proof of Theorem 4.2. By 3.21(v),(vi) we may suppose that $p = fu + g$ is a residual x -variable and $f_{\text{non-horiz}} \in \mathbb{C}[x]$ (cf. 4.4). Thus we must show that actually, under each of the assumptions (a)-(d) p is an x -variable.

(a) (resp., (b)) immediately follows from 4.7 (resp., 4.14).

(c) We write the polynomial $p = fu + g$ in the form (34). The assumption $f_{\text{vert}} = 1$ implies that $q^{-1}(0) = \emptyset$, and hence $\tilde{f}^{-1}(0) \cap \tilde{g}_1^{-1}(0) = \emptyset$. Now the conclusion follows from 4.13.

(d) As by our assumption, $f_{\text{horiz}} = f_{\text{horiz}}^{\text{red}}$ we have $\tilde{f} = \tilde{f}^{\text{red}}$. In view of 4.12 we may suppose that $r = 1$ in (34), and so p can be written as follows:

$$p = fu + g = q[\tilde{f}u + \tilde{g}_0y^2 + \tilde{g}_1z + \tilde{f}\tilde{g}_2z^2] + a_0 + a_1y.$$

The $\mathbb{C}[x, y, z]$ -automorphism $u \mapsto u - z^2\tilde{g}_2(x, y, z)$ transforms p into a residual x -variable of degree at most 1 in z . Now the conclusion follows from (b). \square

The following result extends our list of variables $p = fu + g$ beyond those provided in 4.2.

Proposition 4.16. *Let $p = fu + g \in \mathbb{C}^{[4]}$ as in (34) be a residual x -variable. If $\tilde{g}_1 \in \mathbb{C}[x]$ and $\tilde{g}_2z^2 = \hat{g}_2(x, y, \tilde{g}_1(x)z, u) \in \mathbb{C}[x][y, \tilde{g}_1z]$ then p is an x -variable.*

Proof. By 4.12 we may suppose that $r = 1$ in (34), and so $p = \bar{p}(x, y, \tilde{g}_1(x)z, u)$, where the polynomial

$$(36) \quad \bar{p} := q[\tilde{f}u + \tilde{g}_0y^2 + z + \tilde{f}^{\text{red}}\tilde{g}_2] + a_0 + a_1y$$

still is a residual x -variable. Indeed by our assumption for any $\lambda \in \mathbb{C} \setminus \tilde{g}_1^{-1}(0)$,

$$\bar{p}_\lambda = p_\lambda(y, z/\tilde{g}_1(\lambda), u) \in \text{Var}_{\mathbb{C}}\mathbb{C}^{[3]}.$$

For $x_0 \in g_1^{-1}(0)$, either $q(x_0) = 0$ and then by the assumption, $\bar{p}_{x_0} = a_0(x_0) + a_1(x_0)y = p_{x_0} \in \text{Var}_{\mathbb{C}}\mathbb{C}^{[3]}$, or $q(x_0) \neq 0$ and then again, in virtue of 1.29(a), $\bar{p}_{x_0} \in \text{Var}_B B[z, u]$ with $B := \mathbb{C}[y]$. It follows from 4.13 that the polynomial \bar{p} as in (36) is an x -variable. Hence by 1.30(a) (with z playing the role of u) so is $p = \bar{p}(x, y, \tilde{g}_1(x)z, u)$ as well. \square

Example 4.17. The polynomial

$$p := y + x(xz + y(yu + x^2z^2)) \in \mathbb{C}^{[4]}$$

is an x -variable. Indeed $p = \bar{p}(x, y, xz, u)$, where by 1.29(a), $\bar{p} := y + x(z + y(yu + z^2))$ is a residual x -variable (and hence by 4.13 is an x -variable). Therefore, p is a residual x -variable as well, and so 4.16 applies.

4.18. To conclude, we would like to rise the following question. It concerns a (probably simplest) example of a residual x -variable of $\mathbb{C}^{[4]}$ which does not fit any of the assumptions 4.2(a)-(d) or 4.16.

Question: *Is the polynomial $p = y + x(xz + y(yu + z^2)) \in \mathbb{C}^{[4]}$ a variable?*

4.2. Simple modifications of \mathbb{C}^3 rectifiable in \mathbb{C}^5 . The following definitions are inspired by [30].

4.19. Let B be a commutative ring. We say that $p \in B^{[n]}$ is a B -hyperplane if $B^{[n]}/(p) \cong B^{[n-1]}$. We say that p defines a B -hyperplane fibration if $p - \lambda$ is a B -hyperplane for every $\lambda \in B$.

If $B = \mathbb{C}$ or $B = \mathbb{C}[x]$ we simply say *hyperplane* resp., *x -hyperplane* instead of B -hyperplane.

Notice that an x -hyperplane $p(x, y_1, \dots, y_n) \in \mathbb{C}[x][y_1, \dots, y_n]$ becomes a hyperplane $p_{x_0}(y_1, \dots, y_n)$ for every fixed $x = x_0 \in \mathbb{C}$. A polynomial p with the latter property is called a *residual x -hyperplane*.

4.20. Observe [30, 16, 35] that a polynomial $p = f(x, y)u + g(x, y, z) \in \mathbb{C}^{[4]}$ is a residual x -plane if and only if it is a residual x -variable, and henceforth, if and only if it defines a residual x -plane fibration.

4.21. We say that two polynomials $p, q \in B[y_1, \dots, y_n]$ are *1-stably equivalent* if

$$\gamma((p, v)) = (q, v)$$

for an automorphism $\gamma \in \text{Aut}_B B[y_1, \dots, y_n, v]$.

4.22. For instance, with this terminology 1.31(b) says that $y + ap(y)$ with $a \in B$, $p \in B[y]$ is 1-stably equivalent to $y + p(ay)$, and moreover if $a^k | p(0)$ then $y + ap(y)$ is 1-stably equivalent to $y + p(a^{k+1}y)/a^k$.

Theorem 4.23. *Let $p = f(x, y)u + g(x, y, z) \in \mathbb{C}^{[4]}$ be a residual x -plane. Then it is 1-stably equivalent to an x -variable. Consequently, p defines an x -plane fibration, and each fiber $X_\lambda := \{p = \lambda\}$ ($\lambda \in \mathbb{C}$) of p is a 3-fold in \mathbb{C}^4 isomorphic to \mathbb{C}^3 which can be rectified in \mathbb{C}^5 .*

Remark 4.24. Notice that 4.23 provides yet another proof of the implication (iii) \Rightarrow (i) of 3.21 which does not need 1.7. Indeed, it proves (v) \Rightarrow (i) while (iii) \Rightarrow (v) is observed in 4.20 above.

In the proof of 4.23 (see 4.26 below) we use the following lemma.

Lemma 4.25. *If $p \in \mathbb{C}[x, y, z, u] \setminus \mathbb{C}[x, y]$ is a residual x -plane then it is 1-stably equivalent (under an automorphism $\gamma \in \text{Aut}_B B[y, v]$ with $B := \mathbb{C}[x, z, u]$) to a polynomial $\bar{p} \in \mathbb{C}[x][y, z, u]$ which is still a residual x -plane and such that for every $x_0 \in \mathbb{C}$, $\bar{p}_{x_0}(y, z, u) \notin \mathbb{C}^*y + \mathbb{C}$.*

Proof. As $p \notin \mathbb{C}[x, y]$, the subset

$$L(p) := \{x_0 \in \mathbb{C} \mid p_{x_0} \in \mathbb{C}^*y + \mathbb{C}\} =: \{x_1, \dots, x_n\} \subseteq \mathbb{C}$$

is finite. The proof proceeds by a descending induction on $n = \text{card } L(p)$.

If $\text{card } L(p) = 0$ there is nothing to prove. Suppose that $n > 0$, and let $x_n = 0$. Then there exists an affine automorphism $\tau \in \text{Aut}_{\mathbb{C}[x]} \mathbb{C}[x][y] \subseteq \text{Aut}_D D[y]$ (where $D := \mathbb{C}[x, z, u, v]$) such that $\tau(p) = y + x\hat{p}$, where $\hat{p} \in \mathbb{C}[x, y, z, u]$ and $\hat{p}(x, 0, 0, 0) = 0$. Indeed, up to an affine automorphism of $\mathbb{C}[y]$ we may assume that $p(0, y, z, u) = y$ i.e., $p = y + x\hat{p}$ with $\hat{p} \in \mathbb{C}^4$, and then we apply the shift $y \mapsto y - x\hat{p}(x, 0, 0, 0)$ to obtain $\tau(p) = y + x\hat{p}$ with $\hat{p} := \hat{p} - \hat{p}(x, 0, 0, 0)$.

The polynomial $\hat{p} \in \mathbb{C}^4$ admits a presentation:

$$\hat{p} = yq_1(x, y, z, u) + x^k(xq_2(x, z, u) + q_3(z, u)).$$

As by our assumption for every fixed $x \in \mathbb{C}$ the polynomial $\tau(p) = y + x\hat{p} \notin \mathbb{C}[x, y]$ is irreducible, we have $xq_2 + q_3 \neq 0$, and thereby $q_3 \neq 0$. Furthermore,

$$0 = \hat{p}(x, 0, 0, 0) = x^k(xq_2(x, 0, 0) + q_3(0, 0)) = q_3(0, 0),$$

whence $q_3 \notin \mathbb{C}$. Now by 1.31(b) (cf. 4.22 above) there exists an automorphism $\alpha \in \text{Aut}_B B[y, v]$ (with $B = \mathbb{C}[x, z, u]$) such that

$$\alpha((y + x\hat{p}, v)) = (y + \frac{\hat{p}(x, x^{k+1}y, z, u)}{x^k}, v).$$

Letting $\gamma_n := \alpha\tau$ we obtain:

$$\gamma_n((p, v)) = (p_n, v),$$

where $p_n := y + \hat{p}(x, x^{k+1}y, z, u)/x^k$. We have:

$$\begin{aligned} p_n &= y + \frac{x^{k+1}yq_1(x, x^{k+1}y, z, u) + x^k(xq_2(x, z, u) + q_3(z, u))}{x^k} \\ &= y + xyq_1(x, x^{k+1}y, z, u) + xq_2(x, z, u) + q_3(z, u). \end{aligned}$$

Hence $p_n(0, y, z, u) = y + q_3(z, u) \in \text{Var}_{\mathbb{C}} \mathbb{C}[y, z, u]$ with $q_3 \notin \mathbb{C}$, and so $p_n(0, y, z, u) \notin \mathbb{C}^*y + \mathbb{C}$. In other words,

$$0 = x_n \notin L(p_n) := \{\lambda \in \mathbb{C} \mid (p_n)_\lambda \in \mathbb{C}^*y + \mathbb{C}\} \subseteq \mathbb{C}.$$

Now the induction step and the proof of the lemma are completed by the following

Claim. (a) $L(p_n) = \{x_1, \dots, x_{n-1}\} = L(p) \setminus \{x_n\}$.

(b) Furthermore for any $x_0 \neq 0$, $(p_n)_{x_0} \in \mathbb{C}[y, z, u]$ is a plane.

Proof of the claim. ¹⁵ It is enough to show that for every $x_0 \neq 0$ there exists an affine automorphism $\beta_{x_0} \in \text{Aut}_{\mathbb{C}} \mathbb{C}[y] \subseteq \text{Aut}_{\mathbb{C}[z,u]} \mathbb{C}[z,u][y]$ such that

$$(37) \quad \beta_{x_0}(p_{x_0}) = x_0^{k+1}(p_n)_{x_0}.$$

Indeed, we have:

$$(38) \quad x_0^{k+1}(p_n)_{x_0} = x_0^{k+1}y + x_0 \hat{p}_{x_0}(x_0^{k+1}y, z, u).$$

Consider the linear automorphism $\sigma_{x_0} : y \mapsto x_0^{k+1}y$ extended to $\mathbb{C}[z,u][y]$ in a natural way. From (38) we get:

$$x_0^{k+1}(p_n)_{x_0} = \sigma_{x_0}(y + x_0 \hat{p}_{x_0}(y, z, u)) = \sigma_{x_0}((\tau(p))_{x_0}) = \sigma_{x_0}((\tau_{x_0}(p_{x_0})),$$

where τ_{x_0} denotes the specialization of τ at x_0 . Thus we obtain (37) with $\beta_{x_0} := \sigma_{x_0} \tau_{x_0} \in \text{Aut}_{\mathbb{C}} \mathbb{C}[y]$ extended to $\mathbb{C}[z,u][y]$. Hence $p_{x_0} \in \mathbb{C}^*y + \mathbb{C}$ if and only if so is $(p_n)_{x_0}$. \square

4.26. Proof of Theorem 4.23. If $f = 0$ then in virtue of 4.6(c) and the Abhyankar-Moh-Suzuki Theorem, p itself is an x -variable.

If $f \neq 0$ then by 4.25, $p = fu + g$ is 1-stably equivalent to a polynomial $\bar{p} \in \mathbb{C}^{[4]}$ such that

$$(39) \quad L(\bar{p}) := \{\lambda \in \mathbb{C} \mid \bar{p}\lambda \in \mathbb{C}^*y + \mathbb{C}\} = \emptyset.$$

It is easily seen that the polynomial \bar{p} as constructed in the proof of 4.25 still has the form $\bar{p} = \bar{f}u + \bar{g}$ with $\bar{f} \in \mathbb{C}[x,y]$ and $\bar{g} \in \mathbb{C}[x,y,z]$. It follows from (39) that $\bar{f}_{\text{vert}} = 1$. As \bar{p} is also a residual x -plane, by 4.2(c) it is an x -variable. Consequently,

$$\begin{aligned} \mathbb{C}[x][y,z,u]/(p) &\cong \mathbb{C}[x][y,z,u,v]/(p,v) \cong \mathbb{C}[x][y,z,u,v]/(\bar{p},v) \\ &\cong \mathbb{C}[x][y,z,u,v]/(u,v) \cong \mathbb{C}[x]^{[2]}. \end{aligned}$$

Hence p is an x -plane. In view of 4.20 the same arguments work for every ‘fiber’ $p - \lambda$ of $p = fu + g$ (with $\lambda \in \mathbb{C}[x]$). Therefore $p = fu + g$ defines an x -plane fibration. \square

4.3. On Sathaye-Wright’s Theorem. Recall that the Sathaye Theorem on linear planes [30] cited in the introduction was generalized by D. Wright [37] for the embeddings $\mathbb{C}^2 \hookrightarrow \mathbb{C}^3$ of the form $p = f(x,y)u^n + g(x,y) = 0$, and further generalized in [15] for the acyclic surfaces in \mathbb{C}^3 of this type. Here we prove the following theorem (based on the latter result).

Theorem 4.27. *If $p = f(x,y,z)u^n + g(x,y,z) \in \mathbb{C}^{[4]}$ (with $n \geq 2$) is a residual x -variable then it is an x -variable.*

Proof. By [15, Thm. 7.2], the polynomials f and g satisfy the following condition:

$$(40) \quad \forall \lambda \in \mathbb{C} \quad \exists \alpha_\lambda \in \text{Aut}_{\mathbb{C}} \mathbb{C}[y,z] \quad \text{and} \quad \exists \varphi_\lambda \in \mathbb{C}[y] \quad \text{such that}$$

$$f_\lambda = \alpha_\lambda(\varphi_\lambda) \quad \text{and} \quad g_\lambda = \alpha_\lambda(z).$$

Hence by 4.6(c), g is an x -variable of $\mathbb{C}[x,y,z]$, and so we may suppose that $g = z$ and $\alpha_\lambda \in \text{Aut}_{\mathbb{C}[z]} \mathbb{C}[z][y]$. Then (40) yields:

$$(41) \quad \forall \lambda \in \mathbb{C} \quad \exists q_\lambda \in \mathbb{C}[z] \quad \text{and} \quad \exists \varphi_\lambda \in \mathbb{C}[y] \quad \text{such that}$$

$$f_\lambda = \alpha_\lambda(\varphi_\lambda) = \varphi_\lambda(\alpha_\lambda(y)) = \varphi_\lambda(y + q_\lambda).$$

¹⁵Alternatively, (b) also follows from existence of a $\mathbb{C}[x]$ -isomorphism $\mathbb{C}[x]^{[3]}/(p) \cong \mathbb{C}[x]^{[3]}/(\bar{p})$, see 1.32(b).

Eventually, we prove below (by induction on $\deg_y f$) that for any polynomial $f \in \mathbb{C}^{[3]}$ which satisfies (41), we have:

$$(42) \quad f = \Phi(x, a(x)y + zQ(x, z))$$

with $a \in \mathbb{C}[x]$, $Q \in \mathbb{C}[x, z]$ and $\Phi \in \mathbb{C}[x, y]$. Therefore $p = P(x, a(x)y, z, u)$ is a residual x -variable with $P := \Phi(x, y')u + z \in \text{Var}_{\mathbb{C}[x]} \mathbb{C}[x]^{[3]}$ (where $y' := y + zQ(x, z)$). Thus by 1.30(a), p is an x -variable, as stated.

If $\deg_y f \leq 0$ then (41) implies that $f(x, y, z) = f(x, 0, 0) \in \mathbb{C}[x]$ and (42) follows. Assume further that $\deg_y f \geq 1$. By (41) we have:

$$\partial_y f_\lambda(y, z) = (\varphi_\lambda)'(y + q_\lambda(z)).$$

It follows that the polynomial $\partial_y f$ with $\deg_y \partial_y f = \deg_y f - 1$ also satisfies (41), whence by the inductive hypothesis,

$$\partial_y f = \tilde{\Phi}(x, \tilde{a}(x)y + z\tilde{Q}(x, z))$$

with $\tilde{a} \in \mathbb{C}[x]$, $\tilde{Q} \in \mathbb{C}[x, z]$ and $\tilde{\Phi} \in \mathbb{C}[x, y]$. Thereby,

$$\tilde{a}f = \Psi(x, \tilde{a}(x)y + z\tilde{Q}(x, z)) + zR(x, z) \quad \text{with } \Psi \in \mathbb{C}[x, y] \quad \text{and} \quad R \in \mathbb{C}[x, z],$$

where $\partial_y \Psi = \tilde{\Phi}$. If $\deg_y f = 1 = \deg_y \Psi$ then

$$\tilde{a}f = \Psi_0(x) + \Psi_1(x)[\tilde{a}(x)y + z\tilde{Q}(x, z)] + zR(x, z),$$

and so \tilde{a} divides $\Psi_0(x) + \Psi_1(x)z\tilde{Q}(x, z) + zR(x, z)$ i.e.,

$$\Psi_0(x) + \Psi_1(x)z\tilde{Q}(x, z) + zR(x, z) = \tilde{a}(zQ(x, z) + \Phi_0).$$

It follows that

$$f = \Psi_1(x)y + zQ(x, z) + \Phi_0$$

has the desired form (42).

Thus we may assume that $\deg_y f \geq 2$. By (41) for every $\lambda \in \mathbb{C}$ we obtain:

$$(43) \quad \tilde{a}(\lambda)f_\lambda = \Psi(\lambda, \tilde{a}(\lambda)y + z\tilde{Q}(\lambda, z)) + zR(\lambda, z) = \tilde{a}(\lambda)\varphi_\lambda(y + q_\lambda(z)),$$

and so

$$\Psi(\lambda, \tilde{a}(\lambda)y - \tilde{a}(\lambda)q_\lambda(z) + z\tilde{Q}(\lambda, z)) + zR(\lambda, z) = \tilde{a}(\lambda)\varphi_\lambda(y).$$

As $\deg_y \Psi = \deg_y f \geq 2$ it follows that for every $\lambda \in \mathbb{C}$ such that $\deg_y \Psi(\lambda, y) \geq 0$ we have $\deg_z(z\tilde{Q}(\lambda, z) - \tilde{a}(\lambda)q_\lambda(z)) \leq 0$, and so $\deg_z zR(\lambda, z) \leq 0$ as well i.e., $R(\lambda, z) = 0$ for almost every $\lambda \in \mathbb{C}$. Hence $R = 0$, and from (43)

we obtain:

$$(44) \quad \tilde{a}f = \Psi(x, \tilde{a}(x)y + z\tilde{Q}(x, z)).$$

If $\tilde{Q} = 0$ then $f = \Psi(x, \tilde{a}(x)y)/\tilde{a} \in \mathbb{C}[x, y]$, as desired. If $\tilde{Q} \neq 0$ then (up to factorizing $\tilde{a}(x)y + z\tilde{Q}(x, z)$ and changing appropriately $\Psi(x, y)$) we may assume that $\gcd(\tilde{a}, \tilde{Q}(x, z)) = 1$. Putting in (44) $y = 0$ gives:

$$\tilde{a}(x)f(x, 0, z) = \Psi(x, z\tilde{Q}(x, z)).$$

From this equality and the assumption that $\gcd(\tilde{a}, \tilde{Q}(x, z)) = 1$ it is not difficult to deduce that \tilde{a} divides Ψ in $\mathbb{C}[x, y]$. Thus from (44) with $\Phi := \Psi/\tilde{a}$ we get

$$f = \Phi(x, \tilde{a}(x)y + z\tilde{Q}(x, z)),$$

as needed. \square

REFERENCES

- [1] S.S. Abhyankar, T.T. Moh, *Embedding of the line in the plane*, J. Reine Angew. Math. **276** (1975), 148–166.
- [2] E. Bégle, *Duality theorems for generalized manifolds*, Amer. J. Math. **67** (1945), 59–70.
- [3] A.D.R. Choudary, A. Dimca, *Complex hypersurfaces diffeomorphic to affine spaces*, Kodai Math. J. **17** (1994), 171–178.
- [4] H. Derksen, *Constructive Invariant Theory and the Linearization Problem*, Ph.D. thesis, Basel, 1997.
- [5] A. Dold, *Lectures on algebraic topology*, Springer, Berlin e.a., 1974.
- [6] I.V. Dolgachev, B.Ju. Veisfeiler, *Unipotent group schemes over integral rings*, Math. USSR, Izv. **8** (1975), 761–800.
- [7] A. T. Fomenko and D. B. Fuks, *A course in homotopic topology* (Russian), “Nauka”, Moscow, 1989.
- [8] T. Fujita, *On the topology of non-complete algebraic surfaces*, J. Fac. Sci. Univ. Tokyo, Sect.IA, **29** (1982), 503–566.
- [9] D. B. Fuks, O. Ya. Viro, *Homology and cohomology*, in: Encyclopedia of Math. Sci. Vol.24, Topology-2, VINITI, Moscow, 1988, 123–245 (in Russian).
- [10] M. Gizatullin, *On affine surfaces that can be completed by a nonsingular rational curve*, Math. USSR, Izv. **4** (1970), 787–810.
- [11] S. Kaliman, *Exotic analytic structures and Eisenman intrinsic measures*, Israel Math. J. **88** (1994), 411–423.
- [12] S. Kaliman, *Polynomials with general \mathbb{C}^2 -fibers are variables*, preprint, 1999, 60p. (to appear in Pacific J. Math.).
- [13] S. Kaliman, L. Makar-Limanov, *On Russell–Koras contractible threefolds*, J. of Algebraic Geom. **6** (1997), 247–268.
- [14] S. Kaliman, L. Makar-Limanov, *Locally nilpotent derivations of Jacobian type*, Preprint, 1998, 16p.
- [15] S. Kaliman, M. Zaidenberg, *Affine modifications and affine varieties with a very transitive automorphism group*, Transformation Groups, **4:1** (1999), 53–95.
- [16] S. Kaliman, M. Zaidenberg, *Families of affine planes: the existence of a cylinder*, Preprint of the Max Planck Institut für Mathematik, Bonn, MPI-0075 (2000), 12p.; E-print math.AG/0007119 (to appear in Michigan Math. J.).
- [17] S. Kaliman, M. Zaidenberg, *Miyanishi’s characterization of the affine 3-space does not hold in higher dimensions*, Annales de l’Institut Fourier **50:6** (2000), 1649–1669.
- [18] V.Ya. Lin, M.G. Zaidenberg, *Finiteness theorems for holomorphic mappings*, Encyclopedia of Math. Sci. Vol.9. Several Complex Variables III. Berlin Heidelberg New York e.a.: Springer Verlag, 1989, 113–172.
- [19] L. Makar-Limanov, *On the hypersurface $x + x^2y + z^2 + t^3 = 0$ in \mathbb{C}^4 or a \mathbb{C}^3 -like threefold which is not \mathbb{C}^3* , Israel J. Math. **96** (1996), 419–429.
- [20] L. Makar-Limanov, *Again $x + x^2y + z^2 + t^3 = 0$* , Preprint, 1998, 3p.
- [21] L. Makar-Limanov, *Locally nilpotent derivations of affine domains*, Abstract in: Affine Algebraic Geometry 14.05-20.05.2000, Mathematisches Forschungsinstitut Oberwolfach, Tagungsbericht 21/2000, p. 11.
- [22] W.S. Massey, *Singular Homology Theory*, Springer, New York e.a., 1980.
- [23] C.R.F. Maunder, *Algebraic topology*, The New University Mathematics Series. Van Nostrand Reinhold Company Ltd., London etc., 1970.
- [24] M. Miyanishi, *An algebro-topological characterization of the affine space of dimension three*, Amer. J. Math. **106** (1984), 1469–1485.
- [25] M. Miyanishi, *Algebraic characterization of the affine 3-space*, Proc. Algebraic Geom. Seminar, Singapore, World Scientific, 1987, 53–67.
- [26] M. Miyanishi, *Open algebraic surfaces*, CRM Monograph Series **12**, AMS Providence, RI, 2001.
- [27] R. Rentschler, *Opérations du groupe additif sur le plane affine*, C.R. Acad. Sci. Paris, **267** (1968), 384–387.
- [28] P. Russell, *Simple birational extensions of two dimensional affine rational domains*, Compos. Math. **33** (1976), 197–208.

- [29] P. Russell, A. Sathaye, *On finding and cancelling variables in $k[X, Y, Z]$* , J. Algebra **57** (1979), 151–166.
- [30] A. Sathaye, *On linear planes*, Proc. Amer. Math. Soc. **56** (1976), 1–7.
- [31] A. Sathaye, *Polynomial ring in two variables over a D. V. R.: A criterion*, Invent. Math. **74** (1983), 159–168.
- [32] V. Shpilrain, Jie-Tai Yu, *Embeddings of hypersurfaces in affine spaces*, E-print math.AG/0010211, 2000, 11p.; J. Algebra (to appear).
- [33] V. Shpilrain, Jie-Tai Yu, *Embeddings of curves in the plane*, J. Algebra **217** (1999), 668–678.
- [34] M. Suzuki, *Propriétés topologiques des polynômes de deux variables complexes, et automorphismes algébriques de l'espace \mathbb{C}^2* , J. Math. Soc. Japan, **26** (1974), 241–257.
- [35] S. Vénéreau, *Quelques automorphismes et variables de $A[x, y]$* , preprint, Institut Fourier des mathématiques, **460** (1999), 10p.
- [36] R.L. Wilder, *Topology of manifolds*, Repr. of the 1963 ed. Amer. Mathem. Soc. Colloquium Publications, 32. Providence, Rhode Island, 1979.
- [37] D. Wright, *Cancellation of variables of the form $bT^n - a$* , J. Algebra **52** (1978), 94–100.
- [38] M. Zaidenberg, *Isotrivial families of curves on affine surfaces and characterization of the affine plane*, Math. USSR Izvestiya **30:3** (1988), 503–532. Additions and corrections, *ibid.*, **38:2** (1992), 435–437.
- [39] M. Zaidenberg, *On exotic algebraic structures on affine spaces*, Algebra and Analysis. St. Petersburg Mathem. J., **10:5** (1999), 3–73.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MIAMI, CORAL GABLES, FL 33124, U.S.A.
E-mail address: kaliman@math.miami.edu

UNIVERSITÉ GRENOBLE I, INSTITUT FOURIER, UMR 5582 CNRS-UJF, BP 74, 38402 ST.
MARTIN D'HÈRES CÉDEX, FRANCE
E-mail address: venereau@ujf-grenoble.fr

UNIVERSITÉ GRENOBLE I, INSTITUT FOURIER, UMR 5582 CNRS-UJF, BP 74, 38402 ST.
MARTIN D'HÈRES CÉDEX, FRANCE
E-mail address: zaidenbe@ujf-grenoble.fr