

On the Cohomology of Pseudoeffective Line

Jean-Pierre Demailly

*Dedicated to Professor Yum-Tong Siu on the occasion of his
70th birthday*

1 Introduction and Statement of the Main Results

Let X be a compact Kähler n -dimensional manifold, equipped with a Kähler metric, i.e. a positive definite Hermitian $(1, 1)$ -form $\omega = i \sum_{1 \leq j, k \leq n} \omega_{jk}(z) dz_j \wedge d\bar{z}_k$ such that $d\omega = 0$. By definition a holomorphic line bundle L on X is said to be *pseudoeffective* if there exists a singular hermitian metric h on L , given by $h(z) = e^{-\varphi(z)}$ with respect to a local trivialization $L|_U \simeq U \times \mathbb{C}$, such that the curvature form

$$i \Theta_{L,h} := i \partial \bar{\partial} \varphi \tag{1}$$

is (semi) positive in the sense of currents, i.e. φ is locally integrable and $i \Theta_{L,h} \geq 0$: in other words, the weight function φ is plurisubharmonic (psh) on the corresponding trivializing open set U . A basic concept is the notion of *multiplier ideal sheaf*, introduced in [50].

Definition 1 To any psh function φ on an open subset U of a complex manifold X , one associates the “multiplier ideal sheaf” $\mathcal{I}(\varphi) \subset \mathcal{O}_X|_U$ of germs of holomorphic functions $f \in \mathcal{O}_{X,x}$, $x \in U$, such that $|f|^2 e^{-\varphi}$ is integrable with respect to the Lebesgue measure in some local coordinates near x . We also define the global multiplier ideal sheaf $\mathcal{I}(h) \subset \mathcal{O}_X$ of a hermitian metric h on $L \in \text{Pic}(X)$ to be equal to $\mathcal{I}(\varphi)$ on any open subset U where $L|_U$ is trivial and $h = e^{-\varphi}$. In such a

J.-P. Demailly (✉)
Institut Fourier, Université de Grenoble I, Grenoble, France
e-mail: jean-pierre.demailly@ujf-grenoble.fr

definition, we may in fact assume $i\Theta_{L,h} \geq -C\omega$, i.e. locally $\varphi = \text{psh} + C^\infty$, we say in that case that φ is *quasi-psh*.

Let us observe that a multiplier ideal sheaf $\mathcal{I}(\varphi)$ is left unmodified by adding a smooth function to φ ; for such purposes, the additional C^∞ terms are irrelevant in quasi-psh functions. A crucial and well-known fact is that the ideal sheaves $\mathcal{I}(\varphi) \subset \mathcal{O}_X|_U$ and $\mathcal{I}(h) \subset \mathcal{O}_X$ are always *coherent analytic sheaves*; when $U \subset X$ is a coordinate open ball, this can be shown by observing that $\mathcal{I}(\varphi)$ coincides with the locally stationary limit $\mathcal{I} = \lim \uparrow_{N \rightarrow +\infty} \mathcal{I}_N$ of the increasing sequence of coherent ideals $\mathcal{I}_N = (g_j)_{0 \leq j < N}$ associated with a Hilbert basis $(g_j)_{j \in \mathbb{N}}$ of the Hilbert space of holomorphic functions $f \in \mathcal{O}_X(U)$ such that $\int_U |f|^2 e^{-\varphi} dV_\omega < +\infty$. The proof is a consequence of Hörmander's L^2 estimates applied to weights of the form

$$\psi(z) = \varphi(z) + (n+k) \log |z-x|^2.$$

This easily shows that $\mathcal{I}(\varphi)_x + \mathfrak{m}_x^k = \mathcal{I}_x + \mathfrak{m}_x^k$, and one then concludes that $\mathcal{I}(\varphi)_x = \mathcal{I}_x$ by the Krull lemma. When X is *projective algebraic*, Serre's GAGA theorem implies that $\mathcal{I}(h)$ is in fact a *coherent algebraic sheaf*, in spite of the fact that φ may have very "wild" analytic singularities – e.g. they might be everywhere dense in X in the Euclidean topology. Therefore, in some sense, the multiplier ideal sheaf is a powerful tool to extract algebraic (or at least analytic) data from arbitrary singularities of psh functions. In this context, assuming strict positivity of the curvature, one has the following well-known fundamental vanishing theorem.

Theorem 1 (Nadel Vanishing Theorem, [22, 50]) *Let (X, ω) be a compact Kähler n -dimensional manifold, and let L be a holomorphic line bundle over X equipped with a singular Hermitian metric h . Assume that $i\Theta_{L,h} \geq \varepsilon\omega$ for some $\varepsilon > 0$ on X . Then*

$$H^q(X, \mathcal{O}(K_X \otimes L) \otimes \mathcal{I}(h)) = 0 \quad \text{for all } q \geq 1,$$

where $K_X = \Omega_X^n = \Lambda^n T_X^*$ denotes the canonical line bundle.

The proof follows from an application of Hörmander's L^2 estimates with singular weights, themselves derived from the Bochner-Kodaira identity (see [19, 21, 40]). One should observe that the strict positivity assumption implies L to be big, hence X must be projective, since every compact manifold that is Kähler and Moishezon is also projective (cf. [48, 52, 53]). However, when relaxing the strict positivity assumption, one can enter the world of general compact Kähler manifolds, and their study is one of our main goals.

In many cases, one has to assume that the psh functions involved have milder singularities. We say that a psh or quasi-psh function φ has *analytic singularities* if locally on the domain of definition U of φ one can write

$$\varphi(z) = c \log \sum_{j=1}^N |g_j|^2 + O(1) \tag{2}$$

where the g_j 's are holomorphic functions, $c \in \mathbb{R}_+$ and $O(1)$ means a locally bounded remainder term. Assumption (2) implies that the set of poles $Z = \varphi^{-1}(-\infty)$ is an analytic set, locally defined as $Z = \bigcap g_j^{-1}(0)$, and that φ is locally bounded on $U \setminus Z$. We also refer to this situation by saying that φ has *logarithmic poles*. In general, one introduces the following comparison relations for psh or quasi-psh functions φ and hermitian metrics $h = e^{-\varphi}$; a more flexible comparison relation will be introduced in Sect. 5.

Definition 2 Let φ_1, φ_2 be psh functions on an open subset U of a complex manifold X . We say that

- (a) φ_1 has less singularities than φ_2 , and write $\varphi_1 \preceq \varphi_2$, if for every point $x \in U$, there exists a neighborhood V of x and a constant $C \geq 0$ such that $\varphi_1 \geq \varphi_2 - C$ on V .
- (b) φ_1 and φ_2 have equivalent singularities, and write $\varphi_1 \sim \varphi_2$, if locally near any point of U we have $\varphi_1 - C \leq \varphi_2 \leq \varphi_1 + C$.

Similarly, given a pair of hermitian metrics h_1, h_2 on a line bundle $L \rightarrow X$,

- (a') we say that h_1 is less singular than h_2 , and write $h_1 \preceq h_2$, if locally there exists a constant $C > 0$ such that $h_1 \leq Ch_2$.
- (b') we say that h_1, h_2 have equivalent singularities, and write $h_1 \sim h_2$, if locally there exists a constant $C > 0$ such that $C^{-1}h_2 \leq h_1 \leq Ch_2$.

(of course when h_1 and h_2 are defined on a compact manifold X , the constant C can be taken global on X in (a') and (b')).

Important features of psh singularities are the semi-continuity theorem (see [27]) and the strong openness property recently proved by Guan and Zhou [36–38]. Let U be an open set in a complex manifold X and φ a psh function on U . Following [27], we define the log canonical threshold of φ at a point $z_0 \in U$ by

$$c_{z_0}(\varphi) = \sup \{c > 0 : e^{-2c\varphi} \text{ is } L^1 \text{ on a neighborhood of } z_0\} \in]0, +\infty] \tag{3}$$

(Here L^1 integrability refers to the Lebesgue measure with respect to local coordinates). It is an important invariant of the singularity of φ at z_0 . We refer to [25–27, 29, 45, 50, 55, 61] for further information about properties of the log canonical threshold. In this setting, the semi-continuity theorem can be stated as follows.

Theorem 2 (cf. [27]) *For any given $z_0 \in U$, the map $\text{PSH}(U) \rightarrow]0, +\infty]$, $\varphi \mapsto c_{z_0}(\varphi)$ is upper semi-continuous with respect to the topology of weak convergence on the space of psh functions (the latter topology being actually the same as the topology of L^1_{loc} convergence).*

The original proof of [27] was rather involved and depended on uniform polynomial approximation, combined with a reduction to a semi-continuity theorem for algebraic singularities; the Ohsawa-Takegoshi L^2 extension theorem [51] was used in a crucial way. We will give here a simpler and more powerful derivation due to Hiep [54], still depending on the Ohsawa-Takegoshi theorem, that simultaneously yields effective versions of Berndtsson's result [3] on the openness conjecture, as well as Guan and Zhou's proof of the strong openness conjecture for multiplier ideal sheaves.

Theorem 3 ([36–38]) *Let $\varphi, \psi_j, j \in \mathbb{N}$, be psh functions on an open set U in a complex manifold X . Assume that $\psi_j \leq \varphi$ and that ψ_j converges to φ in L^1_{loc} topology as $j \rightarrow +\infty$. Then for every relatively compact subset $U' \Subset U$, the multiplier ideal sheaves $\mathcal{I}(\psi_j)$ coincide with $\mathcal{I}(\varphi)$ on U' for $j \geq j_0(U') \gg 1$.*

Before going further, notice that the family of multiplier ideals $\lambda \mapsto \mathcal{I}(\lambda\varphi)$ associated with a psh function φ is nonincreasing in $\lambda \in \mathbb{R}_+$. By the Noetherian property of ideal sheaves, they can jump only for a locally finite set of values λ in $[0, +\infty[$, and in particular, there exists a real value $\lambda_0 > 1$ such that

$$\mathcal{I}_+(\varphi) := \lim_{\varepsilon \rightarrow 0^+} \mathcal{I}((1 + \varepsilon)\varphi) = \mathcal{I}(\lambda\varphi), \quad \forall \lambda \in]1, \lambda_0]. \quad (4)$$

We will say that $\mathcal{I}_+(\varphi)$ is the upper semicontinuous regularization of the multiplier ideal sheaf. Berndtsson's result [3] states that the equality $\mathcal{I}(\varphi) = \mathcal{O}_X$ implies $\mathcal{I}_+(\varphi) = \mathcal{O}_X$. If we take $\psi_j = (1 + 1/j)\varphi$ and assume (without loss of generality) that $\varphi \leq 0$, Theorem 3 implies in fact

Corollary 1 *For every psh function φ , the upper semicontinuous regularization coincides with the multiplier ideal sheaf, i.e. $\mathcal{I}_+(\varphi) = \mathcal{I}(\varphi)$.*

Now, if L is a pseudoeffective line bundle, it was observed in [23] that there always exists a unique equivalence class h_{\min} of singular hermitian metrics with minimal singularities, such that $i\Theta_{L, h_{\min}} \geq 0$ (by this we mean that h_{\min} is unique up to equivalence of singularities). In fact, if h_∞ is a smooth metric on L , one can define the corresponding weight φ_{\min} of h_{\min} as an upper envelope

$$\varphi_{\min}(z) = \sup \{ \varphi(z); i\Theta_{L, h_\infty} + i\partial\bar{\partial}\varphi \geq 0, \varphi \leq 0 \text{ on } X \}, \quad (5)$$

and put $h_{\min} = h_\infty e^{-\varphi_{\min}}$. In general, h_{\min} need not have analytic singularities.

An important fact is that one can approximate arbitrary psh functions by psh functions with analytic singularities. The appropriate technique consists of using an asymptotic Bergman kernel procedure (cf. [21] and Sect. 2). If φ is a holomorphic function on a ball $B \subset \mathbb{C}^n$, one puts

$$\varphi_m(z) = \frac{1}{2m} \log \sum_{\ell \in \mathbb{N}} |g_{m,\ell}(z)|^2$$

where $(g_{m,\ell})_{\ell \in \mathbb{N}}$ is a Hilbert basis of the space $\mathcal{H}(B, m\varphi)$ of L^2 holomorphic functions on B such that $\int_B |f|^2 e^{-2m\varphi} dV < +\infty$. When $T = \alpha + dd^c \varphi$ is a closed $(1, 1)$ -current on X in the same cohomology class as a smooth $(1, 1)$ -form α and φ is a quasi-psh potential on X , a sequence of global approximations T_m can be produced by taking a finite covering of X by coordinate balls (B_j) . A partition of unity argument allows to glue the local approximations $\varphi_{m,j}$ of φ on B_j into a global potential φ_m , and one sets $T_m = \alpha + dd^c \varphi_m$. These currents T_m converge weakly to T , are smooth in the complement $X \setminus Z_m$ of an increasing family of analytic subsets $Z_m \subset X$, and their singularities approach those of T . More precisely, the Lelong numbers $\nu(T_m, z)$ converge uniformly to those of T , and whenever $T \geq 0$, it is possible to produce a current T_m that only suffers a small loss of positivity, namely $T_m \geq -\varepsilon_m \omega$ where $\lim_{m \rightarrow +\infty} \varepsilon_m = 0$. These considerations lead in a natural way to the concept of *numerical dimension* of a closed positive $(1, 1)$ -current T . We define

$$\text{nd}(T) = \max \{p = 0, 1, \dots, n; \limsup_{m \rightarrow +\infty} \int_{X \setminus Z_m} (T_m + \varepsilon_m \omega)^p \wedge \omega^{n-p} > 0\}. \quad (6)$$

One can easily show (see Sect. 5) that the right hand side of (6) does not depend on the sequence (T_m) , provided that the singularities approach those of T (we call this an “asymptotically equisingular approximation”).

These concepts are very useful to study cohomology groups with values in pseudoeffective line bundles (L, h) . Without assuming any strict positivity of the curvature, one can obtain at least a hard Lefschetz theorem with coefficients in L . The technique is based on a use of harmonic forms with respect to suitable “equisingular approximations” φ_m of the weight φ of h (in that case we demand that $\mathcal{I}(\varphi_m) = \mathcal{I}(\varphi)$ for all m); the main idea is to work with complete Kähler metrics in the open complements $X \setminus Z_m$ where φ_m is smooth, and to apply a variant of the Bochner formula on these sets. More details can be found in Sect. 4 and in [31].

Theorem 4 ([31]) *Let (L, h) be a pseudo-effective line bundle on a compact Kähler manifold (X, ω) of dimension n , let $\Theta_{L,h} \geq 0$ be its curvature current and $\mathcal{I}(h)$ the associated multiplier ideal sheaf. Then, the wedge multiplication operator $\omega^q \wedge \bullet$ induces a surjective morphism*

$$\Phi_{\omega,h}^q : H^0(X, \Omega_X^{n-q} \otimes L \otimes \mathcal{I}(h)) \longrightarrow H^q(X, \Omega_X^n \otimes L \otimes \mathcal{I}(h)).$$

The special case when L is nef is due to Takegoshi [62]. An even more special case is when L is semipositive, i.e. possesses a smooth metric with semipositive curvature. In that case the multiple ideal sheaf $\mathcal{I}(h)$ coincides with \mathcal{O}_X and we get the following consequence already observed by Enoki [33] and Mourougane [49].

Corollary 2 *Let (L, h) be a semipositive line bundle on a compact Kähler manifold (X, ω) of dimension n . Then, the wedge multiplication operator $\omega^q \wedge \bullet$ induces a surjective morphism*

$$\Phi_\omega^q : H^0(X, \Omega_X^{n-q} \otimes L) \longrightarrow H^q(X, \Omega_X^n \otimes L).$$

It should be observed that although all objects involved in Theorem 4 are algebraic when X is a projective manifold, there is no known algebraic proof of the statement; it is not even clear how to define algebraically $\mathcal{I}(h)$ for the case when $h = h_{\min}$ is a metric with minimal singularity. However, even in the special circumstance when L is nef, the multiplier ideal sheaf is crucially needed.

The next statement is taken from the PhD thesis of Junyan Cao [13]. The proof is a combination of our Bergman regularization techniques, together with an argument of Ch. Mourougane [49] relying on a use of the Calabi-Yau theorem for Monge-Ampère equations.

Theorem 5 ([13, 14]) *Let (L, h) be a pseudoeffective line bundle on a compact Kähler n -dimensional manifold X . Then*

$$H^q(X, K_X \otimes L \otimes \mathcal{I}(h)) = 0 \quad \text{for every } q \geq n - \text{nd}(L, h) + 1,$$

where $\text{nd}(L, h) := \text{nd}(i \mathcal{O}_{L, h})$.

Cao's technique of proof actually yields the result for the upper semicontinuous regularization

$$\mathcal{I}_+(h) = \lim_{\varepsilon \rightarrow 0} \mathcal{I}(h^{1+\varepsilon}) \tag{7}$$

instead of $\mathcal{I}(h)$, but we can apply Guan-Zhou's Theorem 3 to see that the equality $\mathcal{I}_+(h) = \mathcal{I}(h)$ always holds. As a final geometric application of this circle of ideas, we present the following result which was obtained in [16].

Theorem 6 ([16]) *Let X be a compact Kähler threefold that is "strongly simple" in the sense that it has no nontrivial analytic subvariety. Then the Albanese morphism $\alpha : X \rightarrow \text{Alb}(X)$ is a biholomorphism, and therefore X is biholomorphic to a 3-dimensional complex torus \mathbb{C}^3 / Λ .*

I would like to thank the referee wholeheartedly for numerous suggestions that led to substantial improvements of the exposition.

2 Approximation of psh Functions and of Closed (1,1)-Currents

We first recall here the basic result on the approximation of psh functions by psh functions with analytic singularities. The main idea is taken from [Dem92] and relies on the Ohsawa-Takegoshi extension theorem, For other applications to algebraic geometry, see [Dem93b] and Demailly-Kollár [DK01]. Let φ be a psh function on an open set $\Omega \subset \mathbb{C}^n$. Recall that the Lelong number of φ at a point $x_0 \in \Omega$ is defined to be

$$v(\varphi, x_0) = \liminf_{z \rightarrow x_0} \frac{\varphi(z)}{\log |z - x_0|} = \lim_{r \rightarrow 0^+} \frac{\sup_{B(x_0, r)} \varphi}{\log r}. \tag{8}$$

In particular, if $\varphi = \log |f|$ with $f \in \mathcal{O}(\Omega)$, then $v(\varphi, x_0)$ is equal to the vanishing order

$$\text{ord}_{x_0}(f) = \sup\{k \in \mathbb{N}; D^\alpha f(x_0) = 0, \forall |\alpha| < k\}.$$

Theorem 7 *Let φ be a plurisubharmonic function on a bounded pseudoconvex open set $\Omega \subset \mathbb{C}^n$. For every $m > 0$, let $\mathcal{H}_\Omega(m\varphi)$ be the Hilbert space of holomorphic functions f on Ω such that $\int_\Omega |f|^2 e^{-2m\varphi} dV_{2n} < +\infty$ and let $\varphi_m = \frac{1}{2m} \log \sum |g_{m,\ell}|^2$ where $(g_{m,\ell})$ is an orthonormal basis of $\mathcal{H}_\Omega(m\varphi)$. Then there are constants $C_1, C_2 > 0$ independent of m such that*

$$(a) \quad \varphi(z) - \frac{C_1}{m} \leq \varphi_m(z) \leq \sup_{|\zeta - z| < r} \varphi(\zeta) + \frac{1}{m} \log \frac{C_2}{r^n} \text{ for every } z \in \Omega \text{ and } r < d(z, \partial\Omega).$$

In particular, φ_m converges to φ pointwise and in L^1_{loc} topology on Ω when $m \rightarrow +\infty$ and

$$(b) \quad v(\varphi, z) - \frac{n}{m} \leq v(\varphi_m, z) \leq v(\varphi, z) \text{ for every } z \in \Omega.$$

Proof

(a) Note that $\sum |g_{m,\ell}(z)|^2$ is the square of the norm of the evaluation linear form $ev_z : f \mapsto f(z)$ on $\mathcal{H}_\Omega(m\varphi)$, since $g_{m,\ell}(z) = ev_z(g_{m,\ell})$ is the ℓ -th coordinate of ev_z in the orthonormal basis $(g_{m,\ell})$. In other words, we have

$$\sum |g_{m,\ell}(z)|^2 = \sup_{f \in B(1)} |f(z)|^2$$

where $B(1)$ is the unit ball of $\mathcal{H}_\Omega(m\varphi)$ (The sum is called the *Bergman kernel* associated with $\mathcal{H}_\Omega(m\varphi)$). As φ is locally bounded from above, the L^2 topology is actually stronger than the topology of uniform convergence on compact

subsets of Ω . It follows that the series $\sum |g_{m,\ell}|^2$ converges uniformly on Ω and that its sum is real analytic. Moreover, by what we just explained, we have

$$\varphi_m(z) = \sup_{f \in B(1)} \frac{1}{2m} \log |f(z)|^2 = \sup_{f \in B(1)} \frac{1}{m} \log |f(z)|.$$

For $z_0 \in \Omega$ and $r < d(z_0, \partial\Omega)$, the mean value inequality applied to the psh function $|f|^2$ implies

$$\begin{aligned} |f(z_0)|^2 &\leq \frac{1}{\pi^n r^{2n}/n!} \int_{|z-z_0|<r} |f(z)|^2 dV_{2n}(z) \\ &\leq \frac{1}{\pi^n r^{2n}/n!} \exp\left(2m \sup_{|z-z_0|<r} \varphi(z)\right) \int_{\Omega} |f|^2 e^{-2m\varphi} dV_{2n}. \end{aligned}$$

If we take the supremum over all $f \in B(1)$ we get

$$\varphi_m(z_0) \leq \sup_{|z-z_0|<r} \varphi(z) + \frac{1}{2m} \log \frac{1}{\pi^n r^{2n}/n!}$$

and the second inequality in (a) is proved – as we see, this is an easy consequence of the mean value inequality. Conversely, the Ohsawa-Takegoshi extension theorem ([51]) applied to the 0-dimensional subvariety $\{z_0\} \subset \Omega$ shows that for any $a \in \mathbb{C}$ there is a holomorphic function f on Ω such that $f(z_0) = a$ and

$$\int_{\Omega} |f|^2 e^{-2m\varphi} dV_{2n} \leq C_3 |a|^2 e^{-2m\varphi(z_0)},$$

where C_3 only depends on n and $\text{diam } \Omega$. We fix a such that the right hand side is 1. Then $\|f\| \leq 1$ and so we get

$$\varphi_m(z_0) \geq \frac{1}{m} \log |f(z_0)| = \frac{1}{m} \log |a| = \varphi(z) - \log \frac{C_3}{m}.$$

The inequalities given in (a) are thus proved. Taking $r = 1/m$, we find that $\lim_{m \rightarrow +\infty} \sup_{|\xi-z|<1/m} \varphi(\xi) = \varphi(z)$ by the upper semicontinuity of φ , and therefore $\lim \varphi_m(z) = \varphi(z)$, since $\lim \frac{1}{m} \log(C_2 m^n) = 0$.

(b) The above estimates imply

$$\sup_{|z-z_0|<r} \varphi(z) - \frac{C_1}{m} \leq \sup_{|z-z_0|<r} \varphi_m(z) \leq \sup_{|z-z_0|<2r} \varphi(z) + \frac{1}{m} \log \frac{C_2}{r^n}.$$

After dividing by $\log r < 0$ when $r \rightarrow 0$, we infer

$$\frac{\sup_{|z-z_0|<2r} \varphi(z) + \frac{1}{m} \log \frac{C_2}{r^n}}{\log r} \leq \frac{\sup_{|z-z_0|<r} \varphi_m(z)}{\log r} \leq \frac{\sup_{|z-z_0|<r} \varphi(z) - \frac{C_1}{m}}{\log r},$$

and from this and definition (8), it follows immediately that

$$v(\varphi, x) - \frac{n}{m} \leq v(\varphi_m, z) \leq v(\varphi, z).$$

□

Theorem 7 implies in a straightforward manner the deep result of [57] on the analyticity of the Lelong number upperlevel sets.

Corollary 3 ([57]) *Let φ be a plurisubharmonic function on a complex manifold X . Then, for every $c > 0$, the Lelong number upperlevel set*

$$E_c(\varphi) = \{z \in X; v(\varphi, z) \geq c\}$$

is an analytic subset of X .

Proof Since analyticity is a local property, it is enough to consider the case of a psh function φ on a pseudoconvex open set $\Omega \subset \mathbb{C}^n$. The inequalities obtained in Theorem 7 (b) imply that

$$E_c(\varphi) = \bigcap_{m \geq m_0} E_{c-n/m}(\varphi_m).$$

Now, it is clear that $E_c(\varphi_m)$ is the analytic set defined by the equations $g_{m,\ell}^{(\alpha)}(z) = 0$ for all multi-indices α such that $|\alpha| < mc$. Thus $E_c(\varphi)$ is analytic as a (countable) intersection of analytic sets. □

Remark 1 It has been observed by Dano Kim [44] that the functions φ_m produced by Theorem 7 do not in general satisfy $\varphi_{m+1} \succcurlyeq \varphi_m$, in other words their singularities may not always increase monotonically to those of φ . Thanks to the subadditivity result of [18], this is however the case for any subsequence φ_{m_k} such that m_k divides m_{k+1} , e.g. $m_k = 2^k$ or $m_k = k!$ (we will refer to such a sequence below as being a “multiplicative sequence”). In that case, a use of the Ohsawa-Takegoshi theorem on the diagonal of $\Omega \times \Omega$ shows that one can obtain $\varphi_{m_{k+1}} \leq \varphi_{m_k}$ (after possibly replacing φ_{m_k} by $\varphi_{m_k} + C/m_k$ with C large enough), see [18] and [31].

Our next goal is to study the regularization process more globally, i.e. on a compact complex manifold X . For this, we have to take care of cohomology class. It is convenient to introduce $d^c = \frac{i}{4\pi}(\bar{\partial} - \partial)$, so that $dd^c = \frac{i}{2\pi}\partial\bar{\partial}$. Let T be a closed $(1, 1)$ -current on X . We assume that T is *quasi-positive*, i.e. that there exists a $(1, 1)$ -form γ with continuous coefficients such that $T \geq \gamma$; observe that a function φ is

quasi-psh iff its complex Hessian is bounded below by a $(1, 1)$ -form with continuous or locally bounded coefficients, that is, if $dd^c\varphi$ is quasi-positive. The case of positive currents ($\gamma = 0$) is of course the most important.

Lemma 1 *There exists a smooth closed $(1, 1)$ -form α representing the same $\partial\bar{\partial}$ -cohomology class as T and an quasi-psh function φ on X such that $T = \alpha + dd^c\varphi$.*

Proof Select an open covering (B_j) of X by coordinate balls such that $T = dd^c\varphi_j$ over B_j , and construct a global function $\varphi = \sum \theta_j\varphi_j$ by means of a partition of unity $\{\theta_j\}$ subordinate to B_j . Now, we observe that $\varphi - \varphi_k$ is smooth on B_k because all differences $\varphi_j - \varphi_k$ are smooth in the intersections $B_j \cap B_k$ and we can write $\varphi - \varphi_k = \sum \theta_j(\varphi_j - \varphi_k)$. Therefore $\alpha := T - dd^c\varphi$ is smooth. \square

Thanks to Lemma 1, the problem of approximating a quasi-positive closed $(1, 1)$ -current is reduced to approximating a quasi-psh function. In this way, we get

Theorem 8 *Let $T = \alpha + dd^c\varphi$ be a quasi-positive closed $(1, 1)$ -current on a compact Hermitian manifold (X, ω) such that $T \geq \gamma$ for some continuous $(1, 1)$ -form γ . Then there exists a sequence of quasi-positive currents $T_m = \alpha + dd^c\varphi_m$ whose local potentials have the same singularities as $1/2m$ times a logarithm of a sum of squares of holomorphic functions and a decreasing sequence $\varepsilon_m > 0$ converging to 0, such that*

- (a) T_m converges weakly to T ,
- (b) $v(T, x) - \frac{n}{m} \leq v(T_m, x) \leq v(T, x)$ for every $x \in X$;
- (c) $T_m \geq \gamma - \varepsilon_m\omega$.

We say that our currents T_m are approximations of T with analytic singularities (possessing logarithmic poles). Moreover, for any multiplicative subsequence m_k , one can arrange that $T_{m_k} = \alpha + dd^c\varphi_{m_k}$ where (φ_{m_k}) is a non-increasing sequence of potentials.

Proof We just briefly sketch the idea – essentially a partition of unity argument – and refer to [21] for the details. Let us write $T = \alpha + dd^c\varphi$ with α smooth, according to Lemma 1. After replacing T with $T - \alpha$ and γ with $\gamma - \alpha$, we can assume without loss of generality that $\{T\} = 0$, i.e. that $T = dd^c\varphi$ with a quasi-psh function φ on X such that $dd^c\varphi \geq \gamma$. Now, for $\varepsilon > 0$ small, we select a finite covering $(B_j)_{1 \leq j \leq N(\varepsilon)}$ of X by coordinate balls on which there exists an ε -approximation of γ as

$$\sum_{1 \leq \ell \leq n} \lambda_{j,\ell} i dz_\ell^j \wedge d\bar{z}_\ell^j \leq \gamma|_{B_j} \leq \sum_{1 \leq \ell \leq n} (\lambda_{j,\ell} + \varepsilon) i dz_\ell^j \wedge d\bar{z}_\ell^j$$

in terms of holomorphic coordinates $(z_\ell^j)_{1 \leq \ell \leq n}$ on B_j (for this we just diagonalize $\gamma(a_j)$ at the center a_j of B_j , and take the radius of B_j small enough). By construction $\psi_{j,\varepsilon}(z) = \varphi(z) - \sum_{1 \leq \ell \leq n} \lambda_{j,\ell} |z_\ell^j|^2$ is psh on B_ℓ , and we can thus obtain

approximations $\psi_{j,\varepsilon,m}$ of ψ_j by the Bergman kernel process applied on each ball B_j . The idea is to define a global approximation of φ by putting

$$\varphi_{\varepsilon,m}(x) = \frac{1}{m} \log \left(\sum_{1 \leq j \leq N(\varepsilon)} \theta_{j,\varepsilon}(x) \exp \left(m \left(\psi_{j,\varepsilon,m}(x) + \sum_{1 \leq \ell \leq n} (\lambda_{j,\ell} - \varepsilon) |z_\ell^j|^2 \right) \right) \right)$$

where $(\theta_{j,\varepsilon})_{1 \leq j \leq N(\varepsilon)}$ is a partition of unity subordinate to the B_j 's. If we take $\varepsilon = \varepsilon_m$ and $\varphi_m = \varphi_{\varepsilon_m,m}$ where ε_m decays very slowly, then it is not hard to check that $T_m = dd^c \varphi_m$ satisfies the required estimates; it is essentially enough to observe that the derivatives of $\theta_{j,\varepsilon}$ are “killed” by the factor $\frac{1}{m}$ when $m \gg \frac{1}{\varepsilon}$. \square

We need a variant of Theorem 8 providing more “equisingularity” in the sense that the multiplier ideal sheaves are preserved. If one adds the requirement to obtain a non-increasing sequence of approximations of the potential, one can do this only at the expense of accepting “transcendental” singularities, which can no longer be guaranteed to be logarithmic poles.

Theorem 9 *Let $T = \alpha + dd^c \varphi$ be a closed $(1, 1)$ -current on a compact Hermitian manifold (X, ω) , where α is a smooth closed $(1, 1)$ -form and φ a quasi-psh function. Let γ be a continuous real $(1, 1)$ -form such that $T \geq \gamma$. Then one can write $\varphi = \lim_{m \rightarrow +\infty} \tilde{\varphi}_m$ where*

- (a) $\tilde{\varphi}_m$ is smooth in the complement $X \setminus Z_m$ of an analytic set $Z_m \subset X$;
- (b) $\{\tilde{\varphi}_m\}$ is a non-increasing sequence, and $Z_m \subset Z_{m+1}$ for all m ;
- (c) $\int_X (e^{-\varphi} - e^{-\tilde{\varphi}_m}) dV_\omega$ is finite for every m and converges to 0 as $m \rightarrow +\infty$;
- (d) (“equisingularity”) $\mathcal{I}(\tilde{\varphi}_m) = \mathcal{I}(\varphi)$ for all m ;
- (e) $T_m = \alpha + dd^c \tilde{\varphi}_m$ satisfies $T_m \geq \gamma - \varepsilon_m \omega$, where $\lim_{m \rightarrow +\infty} \varepsilon_m = 0$.

Proof (A substantial simplification of the original proof in [31].) As in the previous proof, we may assume that $\alpha = 0$ and $T = dd^c \varphi$, and after subtracting a constant to φ we can also achieve that $\varphi \leq -1$ everywhere on X . For every germ $f \in \mathcal{O}_{X,x}$, (c) implies $\int_U |f|^2 (e^{-\varphi} - e^{-\tilde{\varphi}_m}) dV_\omega < +\infty$ on some neighborhood U of x , hence the integrals $\int_U |f|^2 e^{-\varphi} dV_\omega$ and $\int_U |f|^2 e^{-\tilde{\varphi}_m} dV_\omega$ are simultaneously convergent or divergent, and (d) follows trivially. We define

$$\tilde{\varphi}_m(x) = \sup_{k \geq m} (1 + 2^{-k}) \varphi_{p_k}$$

where (p_k) is a multiplicative sequence that grows fast enough, with $\varphi_{p_{k+1}} \leq \varphi_{p_k} \leq 0$ for all k . Clearly $\tilde{\varphi}_m$ is a non-increasing sequence, and

$$\lim_{m \rightarrow +\infty} \tilde{\varphi}_m(x) = \lim_{k \rightarrow +\infty} \varphi_{p_k}(x) = \varphi(x)$$

at every point $x \in X$. If Z_m is the set of poles of φ_{p_m} , it is easy to see that

$$\tilde{\varphi}_m(x) = \lim_{\ell \rightarrow +\infty} \sup_{k \in [m, \ell]} (1 + 2^{-k})\varphi_{p_k}$$

converges uniformly on every compact subset of $X \setminus Z_m$, since any new term $(1 + 2^{-\ell})\varphi_{p_\ell}$ may contribute to the sup only in case

$$\varphi_{p_\ell} \geq \frac{1 + 2^{-p_m}}{1 + 2^{-p_\ell}} \varphi_{p_m} \quad (\geq 2\varphi_{p_m}),$$

and the difference of that term with respect to the previous term $(1 + 2^{-(\ell-1)})\varphi_{p_{\ell-1}} \geq (1 + 2^{-(\ell-1)})\varphi_{p_\ell}$ is less than $2^{-\ell}|\varphi_{p_\ell}| \leq 2^{1-\ell}|\varphi_{p_m}|$. Therefore $\tilde{\varphi}_m$ is continuous on $X \setminus Z_m$, and getting it to be smooth is only a matter of applying Richberg's approximation technique ([24, 56]). The only serious thing to prove is property (c). To achieve this, we observe that $\{\varphi < \tilde{\varphi}_m\}$ is contained in the union $\bigcup_{k \geq m} \{\varphi < (1 + 2^{-k})\varphi_{p_k}\}$, therefore

$$\int_X (e^{-\varphi} - e^{-\tilde{\varphi}_m}) dV_\omega \leq \sum_{k=m}^{+\infty} \int_X \mathbf{1}_{\varphi < (1+2^{-k})\varphi_{p_k}} e^{-\varphi} dV_\omega \quad (9)$$

and

$$\begin{aligned} \int_X \mathbf{1}_{\varphi < (1+2^{-k})\varphi_{p_k}} e^{-\varphi} dV_\omega &= \int_X \mathbf{1}_{\varphi < (1+2^{-k})\varphi_{p_k}} \exp(2^k \varphi - (2^k + 1)\varphi) dV_\omega \\ &\leq \int_X \mathbf{1}_{\varphi < (1+2^{-k})\varphi_{p_k}} \exp((2^k + 1)(\varphi_{p_k} - \varphi)) dV_\omega \\ &\leq \int_X \mathbf{1}_{\varphi < (1+2^{-k})\varphi_{p_k}} \exp(2p_k(\varphi_{p_k} - \varphi)) dV_\omega \end{aligned} \quad (10)$$

if we take $p_k > 2^{k-1}$ (notice that $\varphi_{p_k} - \varphi \geq 0$). Now, by Lemma 2 below, our integral (10) is finite. By Lebesgue's monotone convergence theorem, we have for k fixed

$$\lim_{p \rightarrow +\infty} \int_X \mathbf{1}_{\varphi < (1+2^{-k})\varphi_p} e^{-\varphi} dV_\omega = 0$$

as a decreasing limit, and we can take p_k so large that $\int_X \mathbf{1}_{\varphi < (1+2^{-k})\varphi_{p_k}} e^{-\varphi} dV_\omega \leq 2^{-k}$. This ensures that property (c) holds true by (9). \square

Lemma 2 *On a compact complex manifold, for any quasi-psh potential φ , the Bergman kernel procedure leads to quasi-psh potentials φ_m with analytic singularities such that*

$$\int_X e^{2m(\varphi_m - \varphi)} dV_\omega < +\infty.$$

Proof By definition of φ_m in Theorem 7, $\exp(2m(\varphi_m))$ is (up to the irrelevant partition of unity procedure) equal to the Bergman kernel $\sum_{\ell \in \mathbb{N}} |g_{m,\ell}|^2$. By local uniform convergence and the Noetherian property, it has the same local vanishing behavior as a finite sum $\sum_{\ell \leq N(m)} |g_{m,\ell}|^2$ with $N(m)$ sufficiently large. Since all terms $g_{m,\ell}$ have L^2 norm equal to 1 with respect to the weight $e^{-2m\varphi}$, our contention follows. \square

Remark 2 A very slight variation of the proof would yield the improved condition

$$(c') \quad \forall \lambda \in \mathbb{R}_+, \quad \int_X (e^{-\lambda\varphi} - e^{-\lambda\tilde{\varphi}_m}) dV_\omega \leq 2^{-m} \text{ for } m \geq m_0(\lambda),$$

and thus an equality $\mathcal{I}(\lambda\tilde{\varphi}_m) = \mathcal{I}(\lambda\varphi)$ for $m \geq m_0(\lambda)$. We just need to replace estimate (9) by

$$\int_X (e^{-m\varphi} - e^{-m\tilde{\varphi}_m}) dV_\omega \leq \sum_{k=m}^{+\infty} \int_X \mathbf{1}_{\varphi < (1+2^{-k})\varphi_{p_k}} e^{-k\varphi} dV_\omega$$

and take p_k so large that $2p_k \geq k(2^k + 1)$ and $\int_{\varphi < (1+2^{-k})\varphi_{p_k}} e^{-k\varphi} dV_\omega \leq 2^{-k-1}$. \square

We also quote the following very simple consequence of Lemma 2, which will be needed a bit later. Since φ_m is less singular than φ , we have of course an inclusion $\mathcal{I}(\lambda\varphi) \subset \mathcal{I}(\lambda\varphi_m)$ for all $\lambda \in \mathbb{R}_+$. Conversely :

Corollary 4 *For every pair of positive real numbers $\lambda' > \lambda > 0$, we have an inclusion of multiplier ideals*

$$\mathcal{I}(\lambda'\varphi_m) \subset \mathcal{I}(\lambda\varphi) \quad \text{as soon as } m \geq \left\lceil \frac{1}{2} \frac{\lambda\lambda'}{\lambda' - \lambda} \right\rceil.$$

Proof If $f \in \mathcal{O}_{X,x}$ and U is a sufficiently small neighborhood of x , the Hölder inequality for conjugate exponents $p, q > 1$ yields

$$\int_U |f|^2 e^{-\lambda\varphi} dV_\omega \leq \left(\int_U |f|^2 e^{-\lambda'\varphi_m} dV_\omega \right)^{1/p} \left(\int_U |f|^2 e^{\frac{q}{p}\lambda'\varphi_m - q\lambda\varphi} dV_\omega \right)^{1/q}.$$

Therefore, if $f \in \mathcal{I}(\lambda'\varphi_m)_x$, we infer that $f \in \mathcal{I}(\lambda\varphi)_x$ as soon as the integral $\int_X e^{\frac{q}{p}\lambda'\varphi_m - q\lambda\varphi} dV_\omega$ is convergent. If we select $p \in]1, \lambda'/\lambda]$, this is implied by the condition $\int_X e^{q\lambda(\varphi_m - \varphi)} dV_\omega < +\infty$. If we further take $q\lambda = 2m_0$ to be an even integer so large that

$$p = \frac{q}{q-1} = \frac{2m_0/\lambda}{2m_0/\lambda - 1} \leq \frac{\lambda'}{\lambda}, \quad \text{e.g. } m_0 = m_0(\lambda, \lambda') = \left\lceil \frac{1}{2} \frac{\lambda\lambda'}{\lambda' - \lambda} \right\rceil,$$

then we indeed have $\int_X e^{2m_0(\varphi_m - \varphi)} dV_\omega \leq \int_X e^{2m(\varphi_m - \varphi)} dV_\omega < +\infty$ for $m \geq m_0(\lambda, \lambda')$, thanks to Lemma 2. \square

Remark 3 Without the monotonicity requirement (b) for the sequence $(\widehat{\varphi}_m)$ in Theorem 9, the strong openness conjecture proved in the next section would directly provide an equisingular sequence, simply by taking

$$\widehat{\varphi}_m = \left(1 + \frac{1}{m}\right) \varphi_m$$

where φ_m is the Bergman approximation sequence. In fact all $\widehat{\varphi}_m$ have analytic singularities and Corollary 4 applied with $\lambda = 1$ and $\lambda' = 1 + 1/m$ shows that $\mathcal{I}(\widehat{\varphi}_m) \subset \mathcal{I}(\varphi)$. Since $\widehat{\varphi}_m \geq (1 + \frac{1}{m})\varphi$, the equality $\mathcal{I}(\widehat{\varphi}_m) = \mathcal{I}(\varphi)$ holds for m large by strong openness, and properties (a), (c), (d), (e) can be seen to hold. However, the sequence $(\widehat{\varphi}_m)$ is not monotone.

3 Semi-continuity of psh Singularities and Proof of the Strong Openness Conjecture

In this section, we present a proof of the strong openness conjecture for multiplier ideal sheaves. Let Ω be a domain in \mathbb{C}^n , $f \in \mathcal{O}(\Omega)$ a holomorphic function, and $\varphi \in \text{PSH}(\Omega)$ a psh function on Ω . For every holomorphic function f on Ω , we introduce the *weighted log canonical threshold* of φ with weight f at z_0

$$c_{f,z_0}(\varphi) = \sup \{c > 0 : |f|^2 e^{-2c\varphi} \text{ is } L^1 \text{ on a neighborhood of } z_0\} \in]0, +\infty].$$

The special case $f = 1$ yields the usual log canonical threshold $c_{z_0}(\varphi)$ that was defined in the introduction. The openness conjectures can be stated as follows.

Conjectures

- (a) (openness conjecture, [27])
The set $\{c > 0 : |f|^2 e^{-2c\varphi} \text{ is } L^1 \text{ on a neighborhood of } z_0\}$ equals the open interval $]0, c_{z_0}(\varphi)[$.
- (b) (strong openness conjecture, [23])
The set $\{c > 0 : |f|^2 e^{-2c\varphi} \text{ is } L^1 \text{ on a neighborhood of } z_0\}$ equals the open interval $]0, c_{f,z_0}(\varphi)[$.

The openness conjecture (a) was first established by Favre and Jonsson [34] in dimension 2 (see also [42, 43]), and 8 years later by Berndtsson [3] in arbitrary dimension. The strong form (b), which is equivalent to Corollary 1, was settled very recently by Guan and Zhou [36]. Their proof uses a sophisticated version of the L^2 -extension theorem of Ohsawa and Takegoshi in combination with the curve selection lemma. They have also obtained related semi-continuity statements in [37] and “effective versions” in [38]. A simplified proof along the same lines has been given by Lempert in [47].

Here, we follow Pham Hoang Hiep’s approach [54], which is more straightforward and avoids the curve selection lemma. It is based on the original version [51] of the L^2 -extension theorem, applied to members of a standard basis for a multiplier ideal sheaf of holomorphic functions associated with a plurisubharmonic function φ . In this way, by means of a simple induction on dimension, one can obtain the strong openness conjecture, and give simultaneously an effective version of the semicontinuity theorem for weighted log canonical thresholds. The main results are contained in the following theorem.

Theorem 10 ([54]) *Let f be a holomorphic function on an open set Ω in \mathbb{C}^n and let φ be a psh function on Ω .*

- (i) (“Semicontinuity theorem”) *Assume that $\int_{\Omega'} e^{-2c\varphi} dV_{2n} < +\infty$ on some open subset $\Omega' \subset \Omega$ and let $z_0 \in \Omega'$. Then there exists $\delta = \delta(c, \varphi, \Omega', z_0) > 0$ such that for every $\psi \in \text{PSH}(\Omega')$, $\|\psi - \varphi\|_{L^1(\Omega')} \leq \delta$ implies $c_{z_0}(\psi) > c$. Moreover, as ψ converges to φ in $L^1(\Omega')$, the function $e^{-2c\psi}$ converges to $e^{-2c\varphi}$ in L^1 on every relatively compact open subset $\Omega'' \Subset \Omega'$.*
- (ii) (“Strong effective openness”) *Assume that $\int_{\Omega'} |f|^2 e^{-2c\varphi} dV_{2n} < +\infty$ on some open subset $\Omega' \subset \Omega$. When $\psi \in \text{PSH}(\Omega')$ converges to φ in $L^1(\Omega')$ with $\psi \leq \varphi$, the function $|f|^2 e^{-2c\psi}$ converges to $|f|^2 e^{-2c\varphi}$ in L^1 norm on every relatively compact open subset $\Omega'' \Subset \Omega'$.*

Corollary 5 (“Strong openness”) *For any plurisubharmonic function φ on a neighborhood of a point $z_0 \in \mathbb{C}^n$, the set $\{c > 0 : |f|^2 e^{-2c\varphi}$ is L^1 on a neighborhood of $z_0\}$ is an open interval $(0, c_{f,z_0}(\varphi))$.*

Corollary 6 (“Convergence from below”) *If $\psi \leq \varphi$ converges to φ in a neighborhood of $z_0 \in \mathbb{C}^n$, then $c_{f,z_0}(\psi) \leq c_{f,z_0}(\varphi)$ converges to $c_{f,z_0}(\varphi)$.*

In fact, after subtracting a large constant to φ , we can assume $\varphi \leq 0$ in both corollaries. Then Corollary 5 is a consequence of assertion (ii) of the main theorem when we take Ω' small enough and $\psi = (1 + \delta)\varphi$ with $\delta \searrow 0$. In Corollary 6, we have by definition $c_{f,z_0}(\psi) \leq c_{f,z_0}(\varphi)$ for $\psi \leq \varphi$, but again (ii) shows that $c_{f,z_0}(\psi)$ becomes $\geq c$ for any given value $c \in (0, c_{f,z_0}(\varphi))$, whenever $\|\psi - \varphi\|_{L^1(\Omega')}$ is sufficiently small.

Remark 4 One cannot remove condition $\psi \leq \varphi$ in assertion (ii) of the main theorem. Indeed, choose $f(z) = z_1$, $\varphi(z) = \log |z_1|$ and $\varphi_j(z) = \log |z_1 + \frac{z_j}{j}|$, for $j \geq 1$. One has $\varphi_j \rightarrow \varphi$ in $L^1_{\text{loc}}(\mathbb{C}^n)$, however $c_{f,0}(\varphi_j) = 1 < c_{f,0}(\varphi) = 2$ for all $j \geq 1$. On the other hand, condition (i) of Theorem 10 does not require any given inequality between φ and ψ . Modulo Berndtsson’s solution of the openness conjecture, (i) follows from the effective semicontinuity result of [27], but (like Guan and Zhou) Hiep’s technique will reprove both by a direct and easier method.

A few preliminaries According to standard techniques in the theory of Gröbner bases, one equips the ring $\mathcal{O}_{\mathbb{C}^n,0}$ of germs of holomorphic functions at 0 with the homogeneous lexicographic order of monomials $z^\alpha = z_1^{\alpha_1} \dots z_n^{\alpha_n}$, that is, $z_1^{\alpha_1} \dots z_n^{\alpha_n} < z_1^{\beta_1} \dots z_n^{\beta_n}$ if and only if $|\alpha| = \alpha_1 + \dots + \alpha_n < |\beta| = \beta_1 + \dots + \beta_n$

or $|\alpha| = |\beta|$ and $\alpha_i < \beta_i$ for the first index i with $\alpha_i \neq \beta_i$. For each $f(z) = a_{\alpha^1} z^{\alpha^1} + a_{\alpha^2} z^{\alpha^2} + \dots$ with $a_{\alpha^j} \neq 0, j \geq 1$ and $z^{\alpha^1} < z^{\alpha^2} < \dots$, we define the *initial coefficient*, *initial monomial* and *initial term* of f to be respectively $\text{IC}(f) = a_{\alpha^1}$, $\text{IM}(f) = z^{\alpha^1}$, $\text{IT}(f) = a_{\alpha^1} z^{\alpha^1}$, and the support of f to be $\text{SUPP}(f) = \{z^{\alpha^1}, z^{\alpha^2}, \dots\}$. For any ideal \mathcal{I} of $\mathcal{O}_{\mathbb{C}^n, 0}$, we define $\text{IM}(\mathcal{I})$ to be the ideal generated by $\{\text{IM}(f)\}_{f \in \mathcal{I}}$. First, we recall the division theorem of Hironaka and the concept of standard basis of an ideal.

Theorem 11 (Division theorem of Hironaka, [1, 5, 6, 32, 35]) *Let $f, g_1, \dots, g_k \in \mathcal{O}_{\mathbb{C}^n, 0}$. Then there exist $h_1, \dots, h_k, s \in \mathcal{O}_{\mathbb{C}^n, 0}$ such that*

$$f = h_1 g_1 + \dots + h_k g_k + s,$$

and $\text{SUPP}(s) \cap \langle \text{IM}(g_1), \dots, \text{IM}(g_k) \rangle = \emptyset$, where $\langle \text{IM}(g_1), \dots, \text{IM}(g_k) \rangle$ denotes the ideal generated by the family $(\text{IM}(g_1), \dots, \text{IM}(g_k))$.

Standard basis of an ideal Let \mathcal{I} be an ideal of $\mathcal{O}_{\mathbb{C}^n, 0}$ and let $g_1, \dots, g_k \in \mathcal{I}$ be such that $\text{IM}(\mathcal{I}) = \langle \text{IM}(g_1), \dots, \text{IM}(g_k) \rangle$. Take $f \in \mathcal{I}$. By the division theorem of Hironaka, there exist $h_1, \dots, h_k, s \in \mathcal{O}_{\mathbb{C}^n, 0}$ such that

$$f = h_1 g_1 + \dots + h_k g_k + s,$$

and $\text{SUPP}(s) \cap \text{IM}(\mathcal{I}) = \emptyset$. On the other hand, since $s = f - h_1 g_1 - \dots - h_k g_k \in \mathcal{I}$, we have $\text{IM}(s) \in \text{IM}(\mathcal{I})$. Therefore $s = 0$ and the g_j 's are generators of \mathcal{I} . By permuting the g_j 's and performing ad hoc subtractions, we can always arrange that $\text{IM}(g_1) < \text{IM}(g_2) < \dots < \text{IM}(g_k)$, and we then say that (g_1, \dots, g_k) is a standard basis of \mathcal{I} .

Theorem 10 will be proved by induction on dimension n . All statements are trivial for $n = 0$. Assume that the theorem holds for dimension $n - 1$. Thanks to the L^2 -extension theorem of Ohsawa and Takegoshi ([51]), one obtains the following key lemma.

Lemma 3 *Let $\varphi \leq 0$ be a plurisubharmonic function and f be a holomorphic function on the polydisc Δ_R^n of center 0 and (poly)radius $R > 0$ in \mathbb{C}^n , such that for some $c > 0$*

$$\int_{\Delta_R^n} |f(z)|^2 e^{-2c\varphi(z)} dV_{2n}(z) < +\infty.$$

Let $\psi_j \leq 0, j \geq 1$, be a sequence of plurisubharmonic functions on Δ_R^n with $\psi_j \rightarrow \varphi$ in $L^1_{\text{loc}}(\Delta_R^n)$, and assume that either $f = 1$ identically or $\psi_j \leq \varphi$ for all $j \geq 1$. Then for every $r < R$ and $\varepsilon \in (0, \frac{1}{2}r]$, there exist a value $w_n \in \Delta_\varepsilon \setminus \{0\}$, an index j_0 , a constant $\tilde{c} > c$ and a sequence of holomorphic functions F_j on $\Delta_r^n, j \geq j_0$, such

that $\text{IM}(F_j) \leq \text{IM}(f)$, $F_j(z) = f(z) + (z_n - w_n) \sum a_{j,\alpha} z^\alpha$ with $|w_n| |a_{j,\alpha}| \leq r^{-|\alpha|} \varepsilon$ for all $\alpha \in \mathbb{N}^n$, and

$$\int_{\Delta_\varepsilon^n} |F_j(z)|^2 e^{-2\tilde{c} \psi_j(z)} dV_{2n}(z) \leq \frac{\varepsilon^2}{|w_n|^2} < +\infty, \quad \forall j \geq j_0.$$

Moreover, one can choose w_n in a set of positive measure in the punctured disc $\Delta_\varepsilon \setminus \{0\}$ (the index $j_0 = j_0(w_n)$ and the constant $\tilde{c} = \tilde{c}(w_n)$ may then possibly depend on w_n).

Proof By Fubini's theorem we have

$$\int_{\Delta_R} \left[\int_{\Delta_R^{n-1}} |f(z', z_n)|^2 e^{-2c \varphi(z', z_n)} dV_{2n-2}(z') \right] dV_2(z_n) < +\infty.$$

Since the integral extended to a small disc $z_n \in \Delta_\eta$ tends to 0 as $\eta \rightarrow 0$, it will become smaller than any preassigned value, say $\varepsilon_0^2 > 0$, for $\eta \leq \eta_0$ small enough. Therefore we can choose a set of positive measure of values $w_n \in \Delta_\eta \setminus \{0\}$ such that

$$\int_{\Delta_R^{n-1}} |f(z', w_n)|^2 e^{-2c \varphi(z', w_n)} dV_{2n-2}(z') \leq \frac{\varepsilon_0^2}{\pi \eta^2} < \frac{\varepsilon_0^2}{|w_n|^2}.$$

Since the main theorem is assumed to hold for $n - 1$, for any $\rho < R$ there exist $j_0 = j_0(w_n)$ and $\tilde{c} = \tilde{c}(w_n) > c$ such that

$$\int_{\Delta_\rho^{n-1}} |f(z', w_n)|^2 e^{-2\tilde{c} \psi_j(z', w_n)} dV_{2n-2}(z') < \frac{\varepsilon_0^2}{|w_n|^2}, \quad \forall j \geq j_0.$$

(For this, one applies part (i) in case $f = 1$, and part (ii) in case $\psi_j \leq \varphi$, using the fact that $\psi = \frac{\tilde{c}}{c} \psi_j$ converges to φ as $\tilde{c} \rightarrow c$ and $j \rightarrow +\infty$). Now, by the L^2 -extension theorem of Ohsawa and Takegoshi (see [51]), there exists a holomorphic function F_j on $\Delta_\rho^{n-1} \times \Delta_R$ such that $F_j(z', w_n) = f(z', w_n)$ for all $z' \in \Delta_\rho^{n-1}$, and

$$\begin{aligned} \int_{\Delta_\rho^{n-1} \times \Delta_R} |F_j(z)|^2 e^{-2\tilde{c} \psi_j(z)} dV_{2n}(z) &\leq C_n R^2 \int_{\Delta_\rho^{n-1}} |f(z', w_n)|^2 e^{-2\tilde{c} \psi_j(z', w_n)} dV_{2n-2}(z') \\ &\leq \frac{C_n R^2 \varepsilon_0^2}{|w_n|^2}, \end{aligned}$$

where C_n is a constant which only depends on n (the constant is universal for $R = 1$ and is rescaled by R^2 otherwise). By the mean value inequality for the

plurisubharmonic function $|F_j|^2$, we get

$$\begin{aligned} |F_j(z)|^2 &\leq \frac{1}{\pi^n (\rho - |z_1|)^2 \dots (\rho - |z_n|)^2} \int_{\Delta_{\rho-|z_1|}(z_1) \times \dots \times \Delta_{\rho-|z_n|}(z_n)} |F_j|^2 dV_{2n} \\ &\leq \frac{C_n R^2 \varepsilon_0^2}{\pi^n (\rho - |z_1|)^2 \dots (\rho - |z_n|)^2 |w_n|^2}, \end{aligned}$$

where $\Delta_\rho(z)$ is the disc of center z and radius ρ . Hence, for any $r < R$, by taking $\rho = \frac{1}{2}(r + R)$ we infer

$$\|F_j\|_{L^\infty(\Delta_r^n)} \leq \frac{2^n C_n^{\frac{1}{2}} R \varepsilon_0}{\pi^{\frac{n}{2}} (R - r)^n |w_n|}. \quad (11)$$

Since $F_j(z', w_n) - f(z', w_n) = 0$, $\forall z' \in \Delta_r^{n-1}$, we can write $F_j(z) = f(z) + (z_n - w_n)g_j(z)$ for some function $g_j(z) = \sum_{\alpha \in \mathbb{N}^n} a_{j,\alpha} z^\alpha$ on $\Delta_r^{n-1} \times \Delta_r$. By (11), we get

$$\begin{aligned} \|g_j\|_{\Delta_r^n} = \|g_j\|_{\Delta_r^{n-1} \times \partial \Delta_r} &\leq \frac{1}{r - |w_n|} \left(\|F_j\|_{L^\infty(\Delta_r^n)} + \|f\|_{L^\infty(\Delta_r^n)} \right) \\ &\leq \frac{1}{r - |w_n|} \left(\frac{2^n C_n^{\frac{1}{2}} R \varepsilon_0}{\pi^{\frac{n}{2}} (R - r)^n |w_n|} + \|f\|_{L^\infty(\Delta_r^n)} \right). \end{aligned}$$

Thanks to the Cauchy integral formula, we find

$$|a_{j,\alpha}| \leq \frac{\|g_j\|_{\Delta_r^n}}{r^{|\alpha|}} \leq \frac{1}{(r - |w_n|) r^{|\alpha|}} \left(\frac{2^n C_n^{\frac{1}{2}} R \varepsilon_0}{\pi^{\frac{n}{2}} (R - r)^n |w_n|} + \|f\|_{L^\infty(\Delta_r^n)} \right).$$

We take in any case $\eta \leq \varepsilon_0 \leq \varepsilon \leq \frac{1}{2}r$. As $|w_n| < \eta \leq \frac{1}{2}r$, this implies

$$|w_n| |a_{j,\alpha}| r^{|\alpha|} \leq \frac{2}{r} \left(\frac{2^n C_n^{\frac{1}{2}} R \varepsilon_0}{\pi^{\frac{n}{2}} (R - r)^n} + \|f\|_{L^\infty(\Delta_r^n)} |w_n| \right) \leq C' \varepsilon_0,$$

for some constant C' depending only on n, r, R and f . This yields the estimates of Lemma 3 for $\varepsilon_0 := C'' \varepsilon$ with C'' sufficiently small. Finally, we prove that $\text{IM}(F_j) \leq \text{IM}(f)$. Indeed, if $\text{IM}(g_j) \geq \text{IM}(f)$, since $|w_n| |a_{j,\alpha}| \leq r^{-|\alpha|} \varepsilon$, we can choose ε small enough such that $\text{IM}(F_j) = \text{IM}(f)$ and $\left| \frac{\text{IC}(F_j)}{\text{IC}(f)} \right| \in (\frac{1}{2}, 2)$. Otherwise, if $\text{IM}(g_j) < \text{IM}(f)$, we have $\text{IM}(F_j) = \text{IM}(g_j) < \text{IM}(f)$. \square

Proof of Theorem 10 By well-known properties of (pluri)potential theory, the L^1 convergence of ψ to φ implies that $\psi \rightarrow \varphi$ almost everywhere, and the assumptions guarantee that φ and ψ are uniformly bounded on every relatively compact subset of Ω' . In particular, after shrinking Ω' and subtracting constants, we can assume that

$\varphi \leq 0$ on Ω . Also, since the L^1 topology is metrizable, it is enough to work with a sequence $(\psi_j)_{j \geq 1}$ converging to φ in $L^1(\Omega')$. Again, we can assume that $\psi_j \leq 0$ and that $\psi_j \rightarrow \varphi$ almost everywhere on Ω' . By a trivial compactness argument, it is enough to show (i) and (ii) for some neighborhood Ω'' of a given point $z_0 \in \Omega'$. We assume here $z_0 = 0$ for simplicity of notation, and fix a polydisc Δ_R^n of center 0 with R so small that $\Delta_R^n \subset \Omega'$. Then $\psi_j(\bullet, z_n) \rightarrow \varphi(\bullet, z_n)$ in the topology of $L^1(\Delta_R^{n-1})$ for almost every $z_n \in \Delta_R$.

Proof of statement (i) in Theorem 10 We have here $\int_{\Delta_R^n} e^{-2c\varphi} dV_{2n} < +\infty$ for $R > 0$ small enough. By Lemma 3 with $f = 1$, for every $r < R$ and $\varepsilon > 0$, there exist $w_n \in \Delta_\varepsilon \setminus \{0\}$, an index j_0 , a number $\tilde{c} > c$ and a sequence of holomorphic functions F_j on $\Delta_r^n, j \geq j_0$, such that $F_j(z) = 1 + (z_n - w_n) \sum a_{j,\alpha} z^\alpha, |w_n| |a_{j,\alpha}| r^{-|\alpha|} \leq \varepsilon$ and

$$\int_{\Delta_r^n} |F_j(z)|^2 e^{-2\tilde{c}\psi_j(z)} dV_{2n}(z) \leq \frac{\varepsilon^2}{|w_n|^2}, \quad \forall j \geq j_0.$$

For $\varepsilon \leq \frac{1}{2}$, we conclude that $|F_j(0)| = |1 - w_n a_{j,0}| \geq \frac{1}{2}$ hence $c_0(\psi_j) \geq \tilde{c} > c$ and the first part of (i) is proved. In fact, after fixing such ε and w_n , we even obtain the existence of a neighborhood Ω'' of 0 on which $|F_j| \geq \frac{1}{4}$, and thus get a uniform bound $\int_{\Omega''} e^{-2\tilde{c}\psi_j(z)} dV_{2n}(z) \leq M < +\infty$. The second assertion of (i) then follows from the estimate

$$\begin{aligned} \int_{\Omega''} |e^{-2c\psi_j(z)} - e^{-2c\varphi(z)}| dV_{2n}(z) &\leq \int_{\Omega'' \cap \{|\psi_j| \leq A\}} |e^{-2c\psi_j(z)} - e^{-2c\varphi(z)}| dV_{2n}(z) \\ &\quad + \int_{\Omega'' \cap \{\psi_j < -A\}} e^{-2c\varphi(z)} dV_{2n}(z) \\ &\quad + e^{-2(\tilde{c}-c)A} \int_{\Omega'' \cap \{\psi_j < -A\}} e^{-2\tilde{c}\psi_j(z)} dV_{2n}(z). \end{aligned}$$

In fact the last two terms converge to 0 as $A \rightarrow +\infty$, and, for A fixed, the first integral in the right hand side converges to 0 by Lebesgue's bounded convergence theorem, since $\psi_j \rightarrow \varphi$ almost everywhere on Ω'' .

Proof of statement (ii) in Theorem 10 Take $f_1, \dots, f_k \in \mathcal{O}_{\mathbb{C}^n,0}$ such that (f_1, \dots, f_k) is a standard basis of $\mathcal{S}(c\varphi)_0$ with $\text{IM}(f_1) < \dots < \text{IM}(f_k)$, and Δ_R^n a polydisc so small that

$$\int_{\Delta_R^n} |f_l(z)|^2 e^{-2c\varphi(z)} dV_{2n}(z) < +\infty, \quad l = 1, \dots, k.$$

Since the germ of f at 0 belongs to the ideal (f_1, \dots, f_k) , we can essentially argue with the f_l 's instead of f . By Lemma 3, for every $r < R$ and $\varepsilon_l > 0$, there exist $w_{n,l} \in \Delta_{\varepsilon_l} \setminus \{0\}$, an index $j_0 = j_0(w_{n,l})$, a number $\tilde{c} = \tilde{c}(w_{n,l}) > c$ and a sequence of holomorphic functions $F_{j,l}$ on $\Delta_r^n, j \geq j_0$, such that $F_{j,l}(z) = 1 + (z_n - w_{n,l}) \sum a_{j,l,\alpha} z^\alpha$,

$|w_{n,l}||a_{j,l,\alpha}|r^{-|\alpha|} \leq \varepsilon_l$ and

$$\int_{\Delta_{\rho}^n} |F_{j,l}(z)|^2 e^{-2\tilde{c}\psi_j(z)} dV_{2n}(z) \leq \frac{\varepsilon_l^2}{|w_{n,l}|^2}, \quad \forall l = 1, \dots, k, \quad \forall j \geq j_0. \quad (12)$$

Since $\psi_j \leq \varphi$ and $\tilde{c} > c$, we get $F_{j,l} \in \mathcal{S}(\tilde{c}\psi_j)_0 \subset \mathcal{S}(c\varphi)_0$. The next step of the proof consists in modifying $(F_{j,l})_{1 \leq l \leq k}$ in order to obtain a standard basis of $\mathcal{S}(c\varphi)_0$. For this, we proceed by selecting successively $\varepsilon_1 \gg \varepsilon_2 \gg \dots \gg \varepsilon_k$ (and suitable $w_{n,l} \in \Delta_{\varepsilon_l} \setminus \{0\}$). We have $\text{IM}(F_{j,1}), \dots, \text{IM}(F_{j,k}) \in \text{IM}(\mathcal{S}(c\varphi)_0)$, in particular $\text{IM}(F_{j,1})$ is divisible by $\text{IM}(f_l)$ for some $l = 1, \dots, k$. Since $\text{IM}(F_{j,1}) \leq \text{IM}(f_1) < \dots < \text{IM}(f_k)$, we must have $\text{IM}(F_{j,1}) = \text{IM}(f_1)$ and thus $\text{IM}(g_{j,1}) \geq \text{IM}(f_1)$. As $|w_{n,1}||a_{j,1,\alpha}| \leq \varepsilon_1$, we will have $\left| \frac{\text{IC}(F_{j,1})}{\text{IC}(f_1)} \right| \in (\frac{1}{2}, 2)$ for ε_1 small enough. Now, possibly after changing ε_2 to a smaller value, we show that there exists a polynomial $P_{j,2,1}$ such that the degree and coefficients of $P_{j,2,1}$ are uniformly bounded, with $\text{IM}(F_{j,2} - P_{j,2,1}F_{j,1}) = \text{IM}(f_2)$ and $\left| \frac{\text{IC}(F_{j,2} - P_{j,2,1}F_{j,1})}{\text{IC}(f_2)} \right| \in (\frac{1}{2}, 2)$. We consider two cases:

Case 1: If $\text{IM}(g_{j,2}) \geq \text{IM}(f_2)$, since $|w_{n,2}||a_{j,2,\alpha}| \leq r^{-|\alpha|}\varepsilon_2$, we can choose ε_2 so small that $\text{IM}(F_{j,2}) = \text{IM}(f_2)$ and $\left| \frac{\text{IC}(F_{j,2})}{\text{IC}(f_2)} \right| \in (\frac{1}{2}, 2)$. We then take $P_{j,2,1} = 0$.

Case 2: If $\text{IM}(g_{j,2}) < \text{IM}(f_2)$, we have $\text{IM}(g_{j,2}) = \text{IM}(F_{j,2}) \in \text{IM}(\mathcal{S}(c\varphi)_0)$. Hence $\text{IM}(g_{j,2})$ is divisible by $\text{IM}(f_l)$ for some $l = 1, \dots, k$. However, since $\text{IM}(g_{j,2}) < \text{IM}(f_2) < \dots < \text{IM}(f_k)$, the only possibility is that $\text{IM}(g_{j,2})$ be divisible by $\text{IM}(f_1)$. Take $b \in \mathbb{C}$ and $\beta, \gamma \in \mathbb{N}^n$ such that $\text{IT}(g_{j,2}) := a_{j,2,\gamma}z^\gamma = bz^\beta \text{IT}(F_{j,1})$. We have $z^\beta \leq z^\gamma = \text{IM}(g_{j,2}) < \text{IM}(f_2)$ and

$$|w_{n,2}||b| = |w_{n,2}| \frac{|\text{IC}(g_{j,2})|}{|\text{IC}(F_{j,1})|} \leq \frac{2|w_{n,2}||a_{j,2,\gamma}|}{|\text{IC}(f_1)|} \leq \frac{2r^{-|\gamma|}\varepsilon_2}{|\text{IC}(f_1)|}$$

can be taken arbitrarily small. Set $\tilde{g}_{j,2}(z) = g_{j,2}(z) - bz^\beta F_{j,1}(z) = \sum \tilde{a}_{j,2,\alpha} z^\alpha$ and

$$\tilde{F}_{j,2}(z) = f_2(z) + (z_n - w_{n,2})\tilde{g}_{j,2}(z) = F_{j,2}(z) - b(z_n - w_{n,2})z^\beta F_{j,1}(z).$$

We have $\text{IM}(\tilde{g}_{j,2}) > \text{IM}(g_{j,2})$. Since $|w_{n,2}||b| = O(\varepsilon_2)$ and $|w_{n,2}||a_{j,2,\alpha}| = O(\varepsilon_2)$, we get $|w_{n,2}||\tilde{a}_{j,2,\alpha}| = O(\varepsilon_2)$ as well. Now, we consider two further cases. If $\text{IM}(\tilde{g}_{j,2}) \geq \text{IM}(f_2)$, we can again change ε_2 for a smaller value so that $\text{IM}(\tilde{F}_{j,2}) = \text{IM}(f_2)$ and $\left| \frac{\text{IC}(\tilde{F}_{j,2})}{\text{IC}(f_2)} \right| \in (\frac{1}{2}, 2)$. Otherwise, if $\text{IM}(\tilde{g}_{j,2}) < \text{IM}(f_2)$, we have $\text{IM}(F_{j,2}) = \text{IM}(g_{j,2}) < \text{IM}(\tilde{F}_{j,2}) = \text{IM}(\tilde{g}_{j,2}) < \text{IM}(f_2)$. Notice that $\{z^\gamma : z^\gamma < \text{IM}(f_2)\}$ is a finite set. By using similar arguments a finite number of times, we find ε_2 so small that $\text{IM}(F_{j,2} - P_{j,2,1}F_{j,1}) = \text{IM}(f_2)$ and $\left| \frac{\text{IC}(F_{j,2} - P_{j,2,1}F_{j,1})}{\text{IC}(f_2)} \right| \in (\frac{1}{2}, 2)$ for some polynomial $P_{j,2,1}$. Repeating the same arguments for $F_{j,3}, \dots, F_{j,k}$, we

select inductively $\varepsilon_l, l = 1, \dots, k$, and construct linear combinations

$$F'_{j,l} = F_{j,l} - \sum_{1 \leq m \leq l-1} P_{j,l,m} F'_{j,m}$$

with polynomials $P_{j,l,m}, 1 \leq m < l \leq k$, possessing uniformly bounded coefficients and degrees, such that $\text{IM}(F'_{j,l}) = \text{IM}(f_j)$ and $\frac{|\text{IC}(F'_{j,l})|}{|\text{IC}(f_j)|} \in (\frac{1}{2}, 2)$ for all $l = 1, \dots, k$ and $j \geq j_0$. This implies that $(F'_{j,1}, \dots, F'_{j,k})$ is also a standard basis of $\mathcal{S}(c\varphi)_0$. By Theorem 1.2.2 in [35], we can find $\rho, K > 0$ so small that there exist holomorphic functions $h_{j,1}, \dots, h_{j,k}$ on Δ_ρ^n with $\rho < r$, such that

$$f = h_{j,1}F'_{j,1} + h_{j,2}F'_{j,2} + \dots + h_{j,k}F'_{j,k} \text{ on } \Delta_\rho^n$$

and $\|h_{j,l}\|_{L^\infty(\Delta_\rho^n)} \leq K\|f\|_{L^\infty(\Delta_\rho^n)}$, for all $l = 1, \dots, k$ (ρ and K only depend on f_1, \dots, f_k). By (12), this implies a uniform bound

$$\int_{\Delta_\rho^n} |f(z)|^2 e^{-2\tilde{c}\psi_j(z)} dV_{2n}(z) \leq M < +\infty$$

for some $\tilde{c} > c$ and all $j \geq j_0$. Take $\Omega'' = \Delta_\rho^n$. We obtain the L^1 convergence of $|f|^2 e^{-2c\psi_j}$ to $|f|^2 e^{-2c\varphi}$ almost exactly as we argued for the second assertion of part (i), by using the estimate

$$\begin{aligned} & \int_{\Omega''} |f|^2 |e^{-2c\psi_j(z)} - e^{-2c\varphi(z)}| dV_{2n}(z) \\ & \leq \int_{\Omega'' \cap \{|\psi_j| \leq A\}} |f|^2 |e^{-2c\psi_j(z)} - e^{-2c\varphi(z)}| dV_{2n}(z) \\ & \quad + \int_{\Omega'' \cap \{\psi_j < -A\}} |f|^2 e^{-2c\varphi(z)} dV_{2n}(z) \\ & \quad + e^{-2(\tilde{c}-c)A} \int_{\Omega'' \cap \{\psi_j < -A\}} |f|^2 e^{-2\tilde{c}\psi_j(z)} dV_{2n}(z). \end{aligned}$$

4 Hard Lefschetz Theorem for Pseudoeffective Line Bundles

4.1 A Variant of the Bochner Formula

We first recall a variation of the Bochner formula that is required in the proof of the Hard Lefschetz Theorem with values in a positively curved (and therefore non flat) line bundle (L, h) . Here the base manifold is a Kähler (non necessarily compact) manifold (Y, ω) . We denote by $|\cdot| = |\cdot|_{\omega, h}$ the pointwise Hermitian norm on

$\Lambda^{p,q}T_Y^* \otimes L$ associated with ω and h , and by $\| \cdot \| = \| \cdot \|_{\omega,h}$ the global L^2 norm

$$\|u\|^2 = \int_Y |u|^2 dV_\omega \quad \text{where} \quad dV_\omega = \frac{\omega^n}{n!}$$

We consider the $\bar{\partial}$ operator acting on (p, q) -forms with values in L , its adjoint $\bar{\partial}_h^*$ with respect to h and the complex Laplace-Beltrami operator $\Delta_h'' = \bar{\partial}\bar{\partial}_h^* + \bar{\partial}_h^*\bar{\partial}$. Let v be a smooth $(n-q, 0)$ -form with compact support in Y . Then $u = \omega^q \wedge v$ satisfies

$$\|\bar{\partial}u\|^2 + \|\bar{\partial}_h^*u\|^2 = \|\bar{\partial}v\|^2 + \int_Y \sum_{I,J} \left(\sum_{j \in J} \lambda_j \right) |u_{IJ}|^2 \quad (13)$$

where $\lambda_1 \leq \dots \leq \lambda_n$ are the curvature eigenvalues of $\Theta_{L,h}$ expressed in an orthonormal frame $(\partial/\partial z_1, \dots, \partial/\partial z_n)$ (at some fixed point $x_0 \in Y$), in such a way that

$$\omega_{x_0} = i \sum_{1 \leq j \leq n} dz_j \wedge d\bar{z}_j, \quad (\Theta_{L,h})_{x_0} = dd^c \varphi_{x_0} = i \sum_{1 \leq j \leq n} \lambda_j dz_j \wedge d\bar{z}_j.$$

Formula (13) follows from the more or less straightforward identity

$$(\bar{\partial}_\varphi^* \bar{\partial} + \bar{\partial} \bar{\partial}_\varphi^*)(v \wedge \omega^q) - (\bar{\partial}_\varphi^* \bar{\partial} v) \wedge \omega^q = q i \bar{\partial} \bar{\partial} \varphi \wedge \omega^{q-1} \wedge v,$$

by taking the inner product with $u = \omega^q \wedge v$ and integrating by parts in the left hand side (we leave the easy details to the reader). Our formula is thus established when v is smooth and compactly supported. In general, we have:

Proposition 1 *Let (Y, ω) be a complete Kähler manifold and (L, h) a smooth Hermitian line bundle such that the curvature possesses a uniform lower bound $\Theta_{L,h} \geq -C\omega$. For every measurable $(n-q, 0)$ -form v with L^2 coefficients and values in L such that $u = \omega^q \wedge v$ has differentials $\bar{\partial}u, \bar{\partial}_h^*u$ also in L^2 , we have*

$$\|\bar{\partial}u\|^2 + \|\bar{\partial}_h^*u\|^2 = \|\bar{\partial}v\|^2 + \int_Y \sum_{I,J} \left(\sum_{j \in J} \lambda_j \right) |u_{IJ}|^2$$

(here, all differentials are computed in the sense of distributions).

Proof Since (Y, ω) is assumed to be complete, there exists a sequence of smooth forms v_ν with compact support in Y (obtained by truncating v and taking the convolution with a regularizing kernel) such that $v_\nu \rightarrow v$ in L^2 and such that $u_\nu = \omega^q \wedge v_\nu$ satisfies $u_\nu \rightarrow u, \bar{\partial}u_\nu \rightarrow \bar{\partial}u, \bar{\partial}_h^*u_\nu \rightarrow \bar{\partial}_h^*u$ in L^2 . By the curvature assumption, the final integral in the right hand side of (13) must be under control (i.e. the integrand becomes nonnegative if we add a term $C\|u\|^2$ on both sides, $C \gg 0$).

We thus get the equality by passing to the limit and using Lebesgue’s monotone convergence theorem. \square

4.2 Proof of Theorem 4

Here X denotes a compact Kähler manifold equipped with a Kähler metric ω , and (L, h) is a pseudoeffective line bundle on X . To fix the ideas, we first indicate the proof in the much simpler case when (L, h) has a smooth metric h (so that $\mathcal{S}(h) = \mathcal{O}_X$), and then treat the general case.

4.2.1 Special Case: (L, h) is Hermitian Semipositive (with a Smooth Metric)

Let $\{\beta\} \in H^q(X, \Omega_X^n \otimes L)$ be an arbitrary cohomology class. By standard L^2 Hodge theory, $\{\beta\}$ can be represented by a smooth harmonic $(0, q)$ -form β with values in $\Omega_X^n \otimes L$. We can also view β as a (n, q) -form with values in L . The pointwise Lefschetz isomorphism produces a unique $(n - q, 0)$ -form α such that $\beta = \omega^q \wedge \alpha$. Proposition 1 then yields

$$\|\bar{\partial}\alpha\|^2 + \int_Y \sum_{I, J} \left(\sum_{j \in J} \lambda_j \right) |\alpha_{IJ}|^2 = \|\bar{\partial}\beta\|^2 + \|\bar{\partial}_h^* \beta\|^2 = 0,$$

and the curvature eigenvalues λ_j are nonnegative by our assumption. Hence $\bar{\partial}\alpha = 0$ and $\{\alpha\} \in H^0(X, \Omega_X^{n-q} \otimes L)$ is mapped to $\{\beta\}$ by $\Phi_{\omega, h}^q = \omega^q \wedge \bullet$.

4.2.2 General Case

There are several difficulties. The first difficulty is that the metric h is no longer smooth and we cannot directly represent cohomology classes by harmonic forms. We circumvent this problem by smoothing the metric on an (analytic) Zariski open subset and by avoiding the remaining poles on the complement. However, some careful estimates have to be made in order to take the error terms into account.

Fix $\varepsilon = \varepsilon_\nu$ and let $h_\varepsilon = h_{\varepsilon_\nu}$ be an approximation of h , such that h_ε is smooth on $X \setminus Z_\varepsilon$ (Z_ε being an analytic subset of X), $\Theta_{L, h_\varepsilon} \geq -\varepsilon\omega$, $h_\varepsilon \leq h$ and $\mathcal{S}(h_\varepsilon) = \mathcal{S}(h)$. This is possible by Theorem 9. Now, we can find a family

$$\omega_{\varepsilon, \delta} = \omega + \delta(i \partial \bar{\partial} \psi_\varepsilon + \omega), \quad \delta > 0$$

of complete Kähler metrics on $X \setminus Z_\varepsilon$, where ψ_ε is a quasi-psh function on X with $\psi_\varepsilon = -\infty$ on Z_ε , ψ_ε smooth on $X \setminus Z_\varepsilon$ and $i \partial \bar{\partial} \psi_\varepsilon + \omega \geq 0$ (see e.g. [19], Théorème 1.5). By construction, $\omega_{\varepsilon, \delta} \geq \omega$ and $\lim_{\delta \rightarrow 0} \omega_{\varepsilon, \delta} = \omega$. We look at the L^2

Dolbeault complex $K_{\varepsilon,\delta}^\bullet$ of (n, \bullet) -forms on $X \setminus Z_\varepsilon$, where the L^2 norms are induced by $\omega_{\varepsilon,\delta}$ on differential forms and by h_ε on elements in L . Specifically

$$K_{\varepsilon,\delta}^q = \left\{ u: X \setminus Z_\varepsilon \rightarrow \Lambda^{n,q} T_X^* \otimes L; \int_{X \setminus Z_\varepsilon} (|u|_{\Lambda^{n,q}\omega_{\varepsilon,\delta} \otimes h_\varepsilon}^2 + |\bar{\partial}u|_{\Lambda^{n,q+1}\omega_{\varepsilon,\delta} \otimes h_\varepsilon}^2) dV_{\omega_{\varepsilon,\delta}} < \infty \right\}.$$

Let $\mathcal{K}_{\varepsilon,\delta}^q$ be the corresponding sheaf of germs of locally L^2 sections on X (the local L^2 condition should hold on X , not only on $X \setminus Z_\varepsilon$!). Then, for all $\varepsilon > 0$ and $\delta \geq 0$, $(\mathcal{K}_{\varepsilon,\delta}^q, \bar{\partial})$ is a resolution of the sheaf $\Omega_X^n \otimes L \otimes \mathcal{I}(h_\varepsilon) = \Omega_X^n \otimes L \otimes \mathcal{I}(h)$. This is because L^2 estimates hold locally on small Stein open sets, and the L^2 condition on $X \setminus Z_\varepsilon$ forces holomorphic sections to extend across Z_ε ([19], Lemma 6.9).

Let $\{\beta\} \in H^q(X, \Omega_X^n \otimes L \otimes \mathcal{I}(h))$ be a cohomology class represented by a smooth form with values in $\Omega_X^n \otimes L \otimes \mathcal{I}(h)$ (one can use a Čech cocycle and convert it to an element in the \mathcal{C}^∞ Dolbeault complex by means of a partition of unity, thanks to the usual De Rham-Weil isomorphism, see also the final proof in Sect. 6 for more details). Then

$$\|\beta\|_{\varepsilon,\delta}^2 \leq \|\beta\|^2 = \int_X |\beta|_{\Lambda^{n,q}\omega \otimes h}^2 dV_\omega < +\infty.$$

The reason is that $|\beta|_{\Lambda^{n,q}\omega \otimes h}^2 dV_\omega$ decreases as ω increases. This is just an easy calculation, shown by comparing two metrics ω, ω' which are expressed in diagonal form in suitable coordinates; the norm $|\beta|_{\Lambda^{n,q}\omega \otimes h}^2$ turns out to decrease faster than the volume dV_ω increases; see e.g. [19], Lemma 3.2; a special case is $q = 0$, then $|\beta|_{\Lambda^{n,q}\omega \otimes h}^2 dV_\omega = i^{n^2} \beta \wedge \bar{\beta}$ with the identification $L \otimes \bar{L} \simeq \mathbb{C}$ given by the metric h , hence the integrand is even independent of ω in that case.

By the proof of the De Rham-Weil isomorphism, the map $\alpha \mapsto \{\alpha\}$ from the cocycle space $Z^q(\mathcal{K}_{\varepsilon,\delta}^\bullet)$ equipped with its L^2 topology, into $H^q(X, \Omega_X^n \otimes L \otimes \mathcal{I}(h))$ equipped with its finite vector space topology, is continuous. Also, Banach's open mapping theorem implies that the coboundary space $B^q(\mathcal{K}_{\varepsilon,\delta}^\bullet)$ is closed in $Z^q(\mathcal{K}_{\varepsilon,\delta}^\bullet)$. This is true for all $\delta \geq 0$ (the limit case $\delta = 0$ yields the strongest L^2 topology in bidegree (n, q)). Now, β is a $\bar{\partial}$ -closed form in the Hilbert space defined by $\omega_{\varepsilon,\delta}$ on $X \setminus Z_\varepsilon$, so there is a $\omega_{\varepsilon,\delta}$ -harmonic form $u_{\varepsilon,\delta}$ in the same cohomology class as β , such that

$$\|u_{\varepsilon,\delta}\|_{\varepsilon,\delta} \leq \|\beta\|_{\varepsilon,\delta}. \quad (14)$$

Let $v_{\varepsilon,\delta}$ be the unique $(n - q, 0)$ -form such that $u_{\varepsilon,\delta} = v_{\varepsilon,\delta} \wedge \omega_{\varepsilon,\delta}^q$ ($v_{\varepsilon,\delta}$ exists by the pointwise Lefschetz isomorphism). Then

$$\|v_{\varepsilon,\delta}\|_{\varepsilon,\delta} = \|u_{\varepsilon,\delta}\|_{\varepsilon,\delta} \leq \|\beta\|_{\varepsilon,\delta} \leq \|\beta\|.$$

As $\sum_{j \in J} \lambda_j \geq -q\varepsilon$ by the assumption on $\Theta_{L, h_\varepsilon}$, the Bochner formula yields

$$\|\bar{\partial}v_{\varepsilon, \delta}\|_{\varepsilon, \delta}^2 \leq q\varepsilon \|u_{\varepsilon, \delta}\|_{\varepsilon, \delta}^2 \leq q\varepsilon \|\beta\|^2.$$

These uniform bounds imply that there are subsequences $u_{\varepsilon, \delta_v}$ and $v_{\varepsilon, \delta_v}$ with $\delta_v \rightarrow 0$, possessing weak- L^2 limits $u_\varepsilon = \lim_{v \rightarrow +\infty} u_{\varepsilon, \delta_v}$ and $v_\varepsilon = \lim_{v \rightarrow +\infty} v_{\varepsilon, \delta_v}$. The limit $v_\varepsilon = \lim_{v \rightarrow +\infty} v_{\varepsilon, \delta_v}$ is with respect to $L^2(\omega) = L^2(\omega_{\varepsilon, 0})$. To check this, notice that in bidegree $(n - q, 0)$, the space $L^2(\omega)$ has the weakest topology of all spaces $L^2(\omega_{\varepsilon, \delta})$; indeed, an easy calculation made in ([19], Lemma 3.2) yields

$$|f|_{\Lambda^{n-q, 0} \omega \otimes h}^2 dV_\omega \leq |f|_{\Lambda^{n-q, 0} \omega_{\varepsilon, \delta} \otimes h}^2 dV_{\omega_{\varepsilon, \delta}} \quad \text{if } f \text{ is of type } (n - q, 0).$$

On the other hand, the limit $u_\varepsilon = \lim_{v \rightarrow +\infty} u_{\varepsilon, \delta_v}$ takes place in all spaces $L^2(\omega_{\varepsilon, \delta})$, $\delta > 0$, since the topology gets stronger and stronger as $\delta \downarrow 0$ [possibly not in $L^2(\omega)$, though, because in bidegree (n, q) the topology of $L^2(\omega)$ might be strictly stronger than that of all spaces $L^2(\omega_{\varepsilon, \delta})$]. The above estimates yield

$$\begin{aligned} \|v_\varepsilon\|_{\varepsilon, 0}^2 &= \int_X |v_\varepsilon|_{\Lambda^{n-q, 0} \omega \otimes h_\varepsilon}^2 dV_\omega \leq \|\beta\|^2, \\ \|\bar{\partial}v_\varepsilon\|_{\varepsilon, 0}^2 &\leq q\varepsilon \|\beta\|_{\varepsilon, 0}^2, \\ u_\varepsilon &= \omega^q \wedge v_\varepsilon \equiv \beta \quad \text{in } H^q(X, \Omega_X^n \otimes L \otimes \mathcal{I}(h_\varepsilon)). \end{aligned}$$

Again, by arguing in a given Hilbert space $L^2(h_{\varepsilon_0})$, we find L^2 convergent subsequences $u_\varepsilon \rightarrow u$, $v_\varepsilon \rightarrow v$ as $\varepsilon \rightarrow 0$, and in this way get $\bar{\partial}v = 0$ and

$$\begin{aligned} \|v\|^2 &\leq \|\beta\|^2, \\ u &= \omega^q \wedge v \equiv \beta \quad \text{in } H^q(X, \Omega_X^n \otimes L \otimes \mathcal{I}(h)). \end{aligned}$$

Theorem 4 is proved. Notice that the equisingularity property $\mathcal{I}(h_\varepsilon) = \mathcal{I}(h)$ is crucial in the above proof, otherwise we could not infer that $u \equiv \beta$ from the fact that $u_\varepsilon \equiv \beta$. This is true only because all cohomology classes $\{u_\varepsilon\}$ lie in the same fixed cohomology group $H^q(X, \Omega_X^n \otimes L \otimes \mathcal{I}(h))$, whose topology is induced by the topology of $L^2(\omega)$ on $\bar{\partial}$ -closed forms (e.g. through the De Rham-Weil isomorphism). \square

Remark 5 In (14), the existence of a harmonic representative holds true only for $\omega_{\varepsilon, \delta}$, $\delta > 0$, because we need to have a complete Kähler metric on $X \setminus Z_\varepsilon$. The trick of employing $\omega_{\varepsilon, \delta}$ instead of a fixed metric ω , however, is not needed when Z_ε is (or can be taken to be) empty. This is the case if (L, h) is such that $\mathcal{I}(h) = \mathcal{O}_X$ and L is nef. Indeed, by definition, L is nef iff there exists a sequence of smooth metrics h_v such that $i\Theta_{L, h_v} \geq -\varepsilon_v \omega$, so we can take the φ_v 's to be everywhere smooth in Theorem 9. However, multiplier ideal sheaves are needed in the surjectivity statement even in case L is nef, as it may happen that $\mathcal{I}(h_{\min}) \neq \mathcal{O}_X$

even then, and $h := \lim h_\nu$ is anyway always more singular than h_{\min} . Let us recall a standard example (see [30, 31]). Let B be an elliptic curve and let V be the rank 2 vector bundle over B which is defined as the (unique) non split extension

$$0 \rightarrow \mathcal{O}_B \rightarrow V \rightarrow \mathcal{O}_B \rightarrow 0.$$

In particular, the bundle V is numerically flat, i.e. $c_1(V) = 0$, $c_2(V) = 0$. We consider the ruled surface $X = \mathbb{P}(V)$. On that surface there is a unique section $C = \mathbb{P}(\mathcal{O}_B) \subset X$ with $C^2 = 0$ and

$$\mathcal{O}_X(C) = \mathcal{O}_{\mathbb{P}(V)}(1)$$

is a nef line bundle. One can check that $L = \mathcal{O}_{\mathbb{P}(V)}(3)$ leads to a *zero* Lefschetz map

$$\omega \wedge \bullet : H^0(X, \Omega_X^1 \otimes L) \longrightarrow H^1(X, K_X \otimes L) \simeq \mathbb{C},$$

so this is a counterexample to Corollary 2 in the nef case. Incidentally, this also shows (in a somewhat sophisticated way) that $\mathcal{O}_{\mathbb{P}(V)}(1)$ is nef but not semipositive, a fact that was first observed in [30].

5 Numerical Dimension of Currents

A large part of this section borrows ideas from S. Boucksom's [7, 8] and Junyan Cao's [14] PhD theses. We try however to give here a slightly more formal exposition. The main difference with S. Boucksom's approach is that we insist on keeping track of singularities of currents and leaving them unchanged, instead of trying to minimize them in each cohomology class.

5.1 Monotone Asymptotically Equisingular Approximations

Let X be a compact complex n -dimensional manifold. We consider the closed convex cone of *pseudoeffective classes*, namely the set $\mathcal{E}(X)$ of cohomology classes $\{\alpha\} \in H^{1,1}(X, \mathbb{R})$ containing a closed positive $(1, 1)$ -current $T = \alpha + dd^c\varphi$ (in the non Kähler case one should use Bott-Chern cohomology groups here, but we will be mostly concerned with the Kähler case in the sequel). We also introduce the set $\mathcal{S}(X)$ of singularity equivalence classes of closed positive $(1, 1)$ -currents $T = \alpha + dd^c\varphi$ (i.e., α being fixed, up to equivalence of singularities of the potentials φ , cf. Definition 2). Clearly, there is a fibration

$$\pi : \mathcal{S}(X) \rightarrow \mathcal{E}(X), \quad T \mapsto \{\alpha\} \in \mathcal{E}(X) \subset H^{1,1}(X, \mathbb{R}). \quad (15)$$

We will denote by $\mathcal{S}_\alpha(X)$ the fiber $\pi^{-1}(\{\alpha\})$ of $\mathcal{S}(X)$ over a given cohomology class $\{\alpha\} \in \mathcal{E}(X)$. Observe that the base $\mathcal{E}(X)$ is a closed convex cone in a finite dimensional vector space, but in general the fiber $\mathcal{S}_\alpha(X)$ must be viewed as a very complicated infinite dimensional space : if we take e.g. $\{\alpha_1\} \in H^{1,1}(\mathbb{P}^n, \mathbb{R})$ to be the unit class $c_1(\mathcal{O}(1))$, then any current $T = \frac{1}{d}[H]$ where H_d is an irreducible hypersurface of degree d defines a point in $\mathcal{S}_{\alpha_1}(\mathbb{P}^n)$, and these points are all distinct. The set $\mathcal{S}(X)$ is nevertheless equipped in a natural way with an addition law $\mathcal{S}(X) \times \mathcal{S}(X) \rightarrow \mathcal{S}(X)$ that maps $\mathcal{S}_\alpha(X) + \mathcal{S}_\beta(X)$ into $\mathcal{S}_{\alpha+\beta}(X)$, a scalar multiplication $\mathbb{R}_+ \times \mathcal{S}(X) \rightarrow \mathcal{S}(X)$ that takes $\lambda \cdot \mathcal{S}_\alpha(X)$ to the fiber $\mathcal{S}_{\lambda\alpha}(X)$. In this way, $\mathcal{S}(X)$ should be viewed as some sort of infinite dimensional convex cone. The fibers $\mathcal{S}_\alpha(X)$ also possess a partial ordering \preceq (cf. Definition 2) such that $\forall j, S_j \preceq T_j \Rightarrow \sum S_j \preceq \sum T_j$, and a fiberwise “min” operation

$$\begin{aligned} \min : \mathcal{S}_\alpha(X) \times \mathcal{S}_\alpha(X) &\longrightarrow \mathcal{S}_\alpha(X), \\ (T_1, T_2) = (\alpha + dd^c \varphi_1, \alpha + dd^c \varphi_2) &\longmapsto T = \alpha + dd^c \max(\varphi_1, \varphi_2), \end{aligned} \quad (16)$$

with respect to which the addition is distributive, i.e.

$$\min(T_1 + S, T_2 + S) = \min(T_1, T_2) + S.$$

Notice that when $T_1 = \frac{1}{d}[H_1]$, $T_2 = \frac{1}{d}[H_2]$ are effective \mathbb{Q} -divisors, all these operations $+$, \cdot , $\min(\bullet)$ and the ordering \preceq coincide with the usual ones known for divisors. Following Junyan Cao [14] (with slightly more restrictive requirements that do not produce much change in practice), we introduce

Definition 3 Let $T = \alpha + dd^c \varphi$ be a closed positive $(1, 1)$ -current on X , where α is a smooth closed $(1, 1)$ -form and φ is a quasi-psh function on X . We say that the sequence of currents $T_k = \alpha + dd^c \psi_k$, $k \in \mathbb{N}$, is a “monotone asymptotically equisingular approximation of T by currents with analytic singularities” if the sequence of potentials (ψ_k) satisfies the following properties:

- (a) (*monotonicity*) The sequence (ψ_k) is non-increasing and converges to φ at every point of X .
- (b) The functions ψ_k have analytic singularities (and $\psi_k \preceq \psi_{k+1}$ by (a)).
- (c) (*lower bound of positivity*)

$$\alpha + dd^c \psi_k \geq -\varepsilon_k \cdot \omega \quad \text{with} \quad \lim_{k \rightarrow +\infty} \varepsilon_k = 0$$

for any given smooth positive hermitian $(1, 1)$ -form ω on X .

- (d) (*asymptotic equisingularity*) For every pair of positive numbers $\lambda' > \lambda > 0$, there exists an integer $k_0(\lambda, \lambda') \in \mathbb{N}$ such that

$$\mathcal{S}(\lambda' \psi_k) \subset \mathcal{S}(\lambda \varphi) \quad \text{for } k \geq k_0(\lambda, \lambda').$$

Remark 6 Without loss of generality, one can always assume that the quasi-psh potentials $\varphi_k = c_k \log |g_k|^2 + O(1)$ have rational coefficients $c_k \in \mathbb{Q}_+$; here again, g_k is a tuple of locally defined holomorphic functions. In fact, after subtracting constants, one can achieve that $\varphi \leq 0$ and $\psi_k \leq 0$ for all k . If the c_k are arbitrary nonnegative real numbers, one can always replace ψ_k by $\psi'_k = (1 - \delta_k)\psi_k$ with a decreasing sequence $\delta_k \in]0, 1[$ such that $\lim \delta_k = 0$ and $(1 - \delta_k)c_k \in \mathbb{Q}_+$. Then (a), (b), (d) are still valid, and (c) holds with $\varepsilon'_k = (1 - \delta_k)\varepsilon_k + C\delta_k$ and C a constant such that $\alpha \geq -C\omega$. \square

The fundamental observation is:

Theorem 12 *If $\psi_k := \varphi_{m_k}$ is the sequence of potentials obtained by the Bergman kernel approximation of $T = \alpha + dd^c\varphi$ given in the proof of Theorem 8 and (m_k) is a multiplicative sequence, then the ψ_k can be arranged to satisfy the positivity, monotonicity and asymptotic equisingularity properties of Definition 3. Moreover, if we start with currents $T \preceq T'$ in the same cohomology class $\{\alpha\}$, we obtain corresponding approximations that satisfy $\psi_k \preceq \psi'_k$.*

Proof By Corollary 4, the asymptotic equisingularity property (d) in Definition 3 is satisfied for $m_k \geq \lceil \frac{1}{2} \frac{\lambda\lambda'}{\lambda' - \lambda} \rceil$. The other properties are already known or obvious, especially the coefficients $c_k = \frac{1}{m_k}$ are just inverses of integers in that case. \square

The following proposition provides a precise comparison of analytic singularities of potentials when their multiplier ideal sheaves satisfy inclusion relations.

Proposition 2 *Let φ, ψ be quasi-psh functions with analytic singularities, let $c > 0$ be the constant such that φ can be expressed as $c \log \sum |g_j|^2 + O(1)$ with holomorphic functions g_j , and let $\lambda \in \mathbb{R}_+$. Denoting $t_+ := \max(t, 0)$, we have the implications*

- (a) $\forall f \in \mathcal{O}_{X,x}, \int_{B_x \ni x} |f|^2 e^{-\lambda\varphi} dV < +\infty \Rightarrow \log |f|^2 \geq \frac{1}{c}(\lambda c - n)_+ \varphi,$
 (b) $\mathcal{I}(\psi) \subset \mathcal{I}(\lambda\varphi) \Rightarrow \int e^{\psi - \lambda\varphi} dV < +\infty$ and $\psi \geq \frac{1}{c}(\lambda c - n)_+ \varphi$ (locally).

Proof Since everything is local, we may assume that φ, ψ are psh functions on a small ball $B \subset \mathbb{C}^n$, and $\varphi(z) = c \log |g|^2 = c \log \sum_{1 \leq j \leq N} |g_j(z)|^2$.

- (a) The convergence of the integral on a small ball B_x of center x implies

$$\int_{B_x} |f|^2 |g|^{-2\lambda c} dV \leq \text{Const} \int_{B_x} |f|^2 e^{-\lambda\varphi} dV < +\infty$$

By the openness of convergence exponents, one gets

$$\int_{B_x} |f|^2 |g|^{-2\lambda + \varepsilon} dV < +\infty$$

for $\varepsilon > 0$ small enough (this can be seen e.g. by using a log resolution of the ideal sheaf (f, g_j)). Now, if $\lambda c \geq n$, Skoda's division theorem [60] implies that each f can be written $f = \sum h_j g_j$ where h_j satisfies a similar estimate where the exponent of $|g|^{-2}$ is decreased by 1. An iteration of the Skoda division theorem for the h_j yields $f \in (g_j)^k$ where $k = (\lfloor \lambda c \rfloor - (n - 1))_+ \geq (\lambda c - n)_+$. Hence

$$\log |f|^2 \leq k \log |g|^2 + C \leq \frac{k}{c} \varphi + C'$$

and (a) is proved.

- (b) If $(f_\ell)_{\ell \in \mathbb{N}}$ is a Hilbert basis of the space of L^2 holomorphic functions f with $\int_B |f|^2 e^{-\psi} dV < +\infty$, the proof of Theorem 7 yields $\psi \leq C + \log \sum |f_\ell|^2$ (and locally the singularity is achieved by a finite sum of f_ℓ 's by the Noetherian property). After possibly shrinking B , the relations $f_\ell \in \mathcal{I}(\psi) \subset \mathcal{I}(\lambda\varphi)$ imply

$$\int_B |f_\ell|^2 e^{-\lambda\varphi} dV < +\infty,$$

hence $\int e^{\psi - \lambda\varphi} dV < +\infty$ locally by taking the sum over ℓ . The inequality proved in (a) for each $f = f_\ell$ also yields

$$\psi \leq \log \sum |f_\ell|^2 + C \leq \frac{1}{c} (\lambda c - n)_+ \varphi + C',$$

and our singularity comparison relation follows. □

Corollary 7 *If $T = \alpha + dd^c\varphi$ is a closed positive $(1, 1)$ -current and $(\psi_k), (\psi'_k)$ are two monotone asymptotically equisingular approximations of φ with analytic singularities, then for every k and every $\varepsilon > 0$, there exists ℓ such that $(1 - \varepsilon)\psi_k \leq \psi'_\ell$ (and vice versa by exchanging the roles of (ψ_k) and (ψ'_k)).*

Proof Let $c > 0$ be the constant occurring in the logarithmic poles of ψ_k (k being fixed). By condition (d) in Definition 3, for $\lambda' > \lambda \gg 1$ we have $\mathcal{I}(\lambda'\psi'_\ell) \subset \mathcal{I}(\lambda\varphi) \subset \mathcal{I}(\lambda\psi_k)$ for $\ell \geq \ell_0(\lambda, \lambda')$ large enough. Proposition 2 (b) implies the singularity estimate $\psi'_\ell \geq \frac{1}{c\lambda'}(c\lambda - n)_+\psi_k$, and the final constant in front of ψ_k can be taken arbitrary close to 1. □

Our next observation is that the $\min(\bullet)$ procedure defined above for currents is well behaved in terms of asymptotic equisingular approximations.

Proposition 3 *Let $T = \alpha + dd^c\varphi$ and $T' = \alpha + dd^c\varphi'$ be closed positive $(1, 1)$ -currents in the same cohomology class $\{\alpha\}$. Let (ψ_k) and (ψ'_k) be respective monotone asymptotically equisingular approximations with analytic singularities and rational coefficients. Then $\max(\psi_k, \psi'_k)$ provides a monotone asymptotically equisingular approximation of $\min(T, T') = \alpha + dd^c \max(\varphi, \varphi')$ with analytic singularities and rational coefficients.*

Proof If $\psi_k = c_k \log |g_k|^2 + O(1)$ and $\psi'_k = c'_k \log |g'_k|^2 + O(1)$, we can write $c_k = p_k/q_k$, $c'_k = p'_k/q'_k$ and

$$\max(\psi_k, \psi'_k) = \frac{1}{q_k q'_k} \log (|g_k|^{2p_k} + |g'_k|^{2p'_k}) + O(1),$$

hence $\max(\psi_k, \psi'_k)$ also has analytic singularities with rational coefficients (this would not be true with our definitions when the ratio c'_k/c_k is irrational, but of course we could just extend a little bit the definition of what we call analytic singularities, e.g. by allowing arbitrary positive real exponents, in order to avoid this extremely minor annoyance). It is well known that

$$\begin{aligned} \alpha + dd^c \psi_k &\geq -\varepsilon_k \omega, & \alpha + dd^c \psi'_k &\geq -\varepsilon'_k \omega \\ \Rightarrow \alpha + dd^c \max(\psi_k, \psi'_k) &\geq -\max(\varepsilon_k, \varepsilon'_k) \omega. \end{aligned}$$

Finally, if $\psi_{B,k}$ (resp. $\psi'_{B,k}$ and $\tilde{\psi}_{B,k}$) comes from the Bergman approximation of φ (resp. of φ' and $\tilde{\varphi} := \max(\varphi, \varphi')$), we have

$$\tilde{\varphi} \geq \varphi \Rightarrow \tilde{\psi}_{B,k} \geq \psi_{B,k}, \quad \tilde{\varphi} \geq \varphi' \Rightarrow \tilde{\psi}_{B,k} \geq \psi'_{B,k}$$

hence $\tilde{\psi}_{B,k} \geq \max(\psi_{B,k}, \psi'_{B,k})$ and so $\tilde{\psi}_{B,k} \leq \max(\psi_{B,k}, \psi'_{B,k})$. However, for every $\varepsilon > 0$, one has $(1-\varepsilon)\psi_{B,k} \leq \psi_\ell$ and $(1-\varepsilon)\psi'_{B,k} \leq \psi'_\ell$ for $\ell \geq \ell_0(k, \varepsilon)$ large, therefore $(1-\varepsilon)\tilde{\psi}_{B,k} \leq \max(\psi_\ell, \psi'_\ell)$. This shows that $\max(\psi_\ell, \psi'_\ell)$ has enough singularities (the ‘‘opposite’’ inequality $\max(\psi_\ell, \psi'_\ell) \geq \tilde{\varphi} = \max(\varphi, \varphi')$, i.e. $\max(\psi_\ell, \psi'_\ell) \leq \tilde{\varphi}$, holds trivially). \square

Following Junyan Cao [15], we now investigate the additivity properties of the Bergman approximation procedure.

Theorem 13 *Let $T = \alpha + dd^c \varphi$ and $T' = \beta + dd^c \varphi'$ be closed $(1, 1)$ -currents in cohomology classes $\{\alpha\}, \{\beta\} \in \mathcal{E}(X)$. Then for every multiplicative sequence (m_k) , the sum $\varphi_{m_k} + \varphi'_{m_k}$ of the Bergman approximations of φ, φ' gives a monotone asymptotically equisingular approximation of $\varphi + \varphi'$ and $T + T'$.*

Proof Let $\tilde{\varphi}_m$ be the Bergman kernel approximations of $\tilde{\varphi} = \varphi + \varphi'$. By the subadditivity property of ideal sheaves $\mathcal{I}(m\varphi + m\varphi') \subset \mathcal{I}(m\varphi)\mathcal{I}(m\varphi')$ ([18, Th. 2.6]), hence we have $\varphi_m + \varphi'_m \leq \tilde{\varphi}_m$. By Definition 3 (d), Theorem 12 and Corollary 7, to prove Theorem 13, it is sufficient to prove that for every $m \in \mathbb{N}$ fixed, there exists a positive sequence $\lim_{p \rightarrow +\infty} \varepsilon_p = 0$ such that

$$(1 - \varepsilon_p)\tilde{\varphi}_m \leq \varphi_p + \varphi'_p \quad \text{for every } p \gg 1. \quad (17)$$

For every $m \in \mathbb{N}$ fixed, there exists a bimeromorphic map $\pi : \widetilde{X} \rightarrow X$, such that

$$\tilde{\varphi}_m \circ \pi = \sum_i c_i \ln |s_i| + C^\infty \quad \text{for some } c_i > 0, \quad (18)$$

and the effective divisor $\sum_i \text{Div}(s_i)$ is normal crossing. By the construction of $\tilde{\varphi}_m$, we have $\tilde{\varphi}_m \leq \varphi + \varphi'$. Therefore

$$\tilde{\varphi}_m \circ \pi \leq (\varphi + \varphi') \circ \pi. \quad (19)$$

By Siu's decomposition formula for closed positive currents applied to $dd^c(\varphi \circ \pi)$, $dd^c(\varphi' \circ \pi)$ respectively, the divisorial parts add up to produce a divisor that is at least equal to the divisorial part in $dd^c(\tilde{\varphi}_m \circ \pi)$, thus (19) and (18) imply the existence of numbers $a_i, b_i \geq 0$ satisfying

- (i) $a_i + b_i = c_i$ for every i ,
- (ii) $\sum_i a_i \ln |s_i| \leq \varphi \circ \pi$ and $\sum_i b_i \ln |s_i| \leq \varphi' \circ \pi$.

Let $p \in \mathbb{N}$ be an integer, J be the Jacobian of π , $f \in \mathcal{S}(p\varphi)_x$ and $g \in \mathcal{S}(p\varphi')_x$ for some $x \in X$. The inequalities in (ii) and a change of variables $w = \pi(z)$ in the L^2 integrals yield

$$\int_{\pi^{-1}(U_x)} \frac{|f \circ \pi|^2 |J|^2}{\prod_i |s_i|^{2pa_i}} < +\infty \quad \text{and} \quad \int_{\pi^{-1}(U_x)} \frac{|g \circ \pi|^2 |J|^2}{\prod_i |s_i|^{2pb_i}} < +\infty \quad (20)$$

for some small open neighborhood U_x of x . Since $\sum_i \text{Div}(s_i)$ is normal crossing, (20) implies that

$$\sum_i (pa_i - 1) \ln |s_i| \leq \ln(|f \circ \pi|) + \ln |J| \quad \text{and} \quad \sum_i (pb_i - 1) \ln |s_i| \leq \ln(|g \circ \pi|) + \ln |J|.$$

Combining this with (i), we get

$$\sum_i (pc_i - 2) \ln |s_i| \leq \ln(|(f \cdot g) \circ \pi|) + 2 \ln |J|. \quad (21)$$

Note that J is independent of p , and $c_i > 0$. (21) implies thus that, when $p \rightarrow +\infty$, we can find a sequence $\varepsilon_p \rightarrow 0^+$, such that

$$\sum_i pc_i(1 - \varepsilon_p) \ln |s_i| \leq \ln |(f \cdot g) \circ \pi|. \quad (22)$$

Since f (respectively g) is an arbitrary element in $\mathcal{S}(p\varphi)$ (respectively $\mathcal{S}(p\varphi')$), by the construction of φ_p and φ'_p , (22) implies that

$$\sum_i c_i(1 - \varepsilon_p) \ln |s_i| \leq (\varphi_p + \varphi'_p) \circ \pi.$$

Combining this with the fact that $(1 - \varepsilon_p)\tilde{\varphi}_m \circ \pi \sim \sum_i c_i(1 - \varepsilon_p) \ln |s_i|$, we get

$$(1 - \varepsilon_p)\tilde{\varphi}_m \circ \pi \leq (\varphi_p + \varphi'_p) \circ \pi.$$

Therefore $(1 - \varepsilon_p)\tilde{\varphi}_m \leq \varphi_p + \varphi'_p$ and (17) is proved. \square

This motivates the following formal definition.

Definition 4 For each class $\{\alpha\} \in \mathcal{E}(X)$, we define $\widehat{\mathcal{S}}_\alpha(X)$ as a set of equivalence classes of sequences of quasi-positive currents $T_k = \alpha + dd^c \psi_k$ such that

- (a) $T_k = \alpha + dd^c \psi_k \geq -\varepsilon_k \cdot \omega$ with $\lim_{k \rightarrow +\infty} \varepsilon_k = 0$,
- (b) the functions ψ_k have analytic singularities and $\psi_k \leq \psi_{k+1}$ for all k .

We say that (T_k) is weakly less singular than (T'_k) in $\widehat{\mathcal{S}}_\alpha(X)$, and write $(T_k) \leq_W (T'_k)$, if for every $\varepsilon > 0$ and k , there exists ℓ such that $(1 - \varepsilon)T_k \leq T'_\ell$. Finally, we write $(T_k) \sim_W (T'_k)$ when we have $(T_k) \leq_W (T'_k)$ and $(T'_k) \leq_W (T_k)$, and define $\widehat{\mathcal{S}}_\alpha(X)$ to be the quotient space by this equivalence relation.

The set

$$\widehat{\mathcal{S}}(X) = \bigcup_{\{\alpha\} \in \mathcal{E}(X)} \widehat{\mathcal{S}}_\alpha(X) \quad (23)$$

is by construction a fiber space $\hat{\pi} : \widehat{\mathcal{S}}(X) \rightarrow \mathcal{E}(X)$, and, by fixing a multiplicative sequence such as $m_k = 2^k$, we find a natural ‘‘Bergman approximation functional’’

$$\mathbf{B} : \mathcal{S}(X) \rightarrow \widehat{\mathcal{S}}(X), \quad T = \alpha + dd^c \varphi \mapsto (T_{B,k}), \quad T_k = \alpha + dd^c \psi_{B,k} \quad (24)$$

where $\psi_{B,k} := \varphi_{m_k}$ is the corresponding subsequence of the sequence of Bergman approximations (φ_m) .

The set $\widehat{\mathcal{S}}(X)$ is equipped with a natural addition $(T_k) + (T'_k) = (T_k + T'_k)$, with a scalar multiplication $\lambda \cdot (T_k) = (\lambda T_k)$ for $\lambda \in \mathbb{R}_+$, as well as with the $\min(\bullet)$ operation $\min((T_k), (T'_k)) = (\min(T_k, T'_k))$ obtained by taking $\max(\psi_k, \psi'_k)$ of the corresponding potentials. By Proposition 3, \mathbf{B} is a morphism for the $\min(\bullet)$ operation, and by Theorem 13, \mathbf{B} is also a morphism for addition. Accordingly, it is natural to define a weak equivalence of singularities for closed positive currents by

$$T \leq_W T' \iff_{\text{def}} (T_{B,k}) \leq_W (T'_{B,k}), \quad (25)$$

$$T \sim_W T' \iff T \leq_W T' \text{ and } T' \leq_W T. \quad (26)$$

Related ideas are discussed in [4] (especially § 5), using the theory of valuations. One can summarize the above results in the following statement.

Theorem 14 *The Bergman approximation functional*

$$\mathbf{B} : \mathcal{S}(X) \rightarrow \widehat{\mathcal{S}}(X), \quad T = \alpha + dd^c \varphi \mapsto (T_{B,k})$$

is a morphism for addition and for the $\min(\bullet)$ operation on currents. Moreover \mathbf{B} induces an injection $\mathcal{S}(X)/\sim_W \rightarrow \widehat{\mathcal{S}}(X)$.

Remark 7 It is easy to see that the induced map $\mathcal{S}(X)/\sim_W \rightarrow \widehat{\mathcal{S}}(X)$ is an isomorphism when $\dim X = 1$. However, this map is not always surjective when $\dim X \geq 2$. In fact, [30, Example 1.7] exhibits a ruled surface over an elliptic curve Γ and a nef line bundle L over X , such that $\alpha = c_1(L)$ contains a unique closed positive current $T = [C]$, for some curve $C \subset X$ that is a section of $X \rightarrow \Gamma$. Then the Bergman approximation is (up to equivalence of singularities) the constant sequence $T_{B,k} = T$, while $\widehat{\mathcal{S}}_\alpha(X)$ also contains a sequence of smooth currents $T_k \geq -\varepsilon_k \omega$. This implies that $\mathcal{S}(X) \rightarrow \widehat{\mathcal{S}}(X)$ is not surjective in this situation. The following proposition shows however that the “formal elements” (T_k) from $\widehat{\mathcal{S}}(X)$ do not carry larger singularities than the closed positive current classes in $\mathcal{S}(X)$ (the latter being constrained by the singularities of the “limiting currents” T representing the class).

Proposition 4 *Let $T_k = \alpha + dd^c \psi_k$ be a sequence of closed $(1, 1)$ -currents representing an element in $\widehat{\mathcal{S}}_\alpha(X)$. Then there exists a closed positive current $T \in \alpha$ such that $(T_k) \preceq_W (T_{B,k})$.*

Proof We have $T_k \geq -\varepsilon_k \omega$ and $\psi_k \preceq \psi_{k+1}$ for some decreasing sequence $\varepsilon_k \downarrow 0$. We replace ψ_k by setting

$$\tilde{\psi}_k(x) = \sup \{ \tau(x) ; \sup_X \tau \leq 0, \alpha + dd^c \tau \geq -\varepsilon_k \omega, \text{ and } \exists C > 0, \tau \leq \psi_k + C \}.$$

Then $(\tilde{\psi}_k)$ is a decreasing sequence for the usual order relation \leq and $\tilde{\psi}_k \sim \psi_k$ (the argument to prove the equivalence of singularities is similar to the one already used in the proof of Theorem 13, clearly $\tilde{\psi}_k \geq \psi_k - M_k$ where $M_k = \sup_X \psi_k$, and the converse inequality $\tilde{\psi}_k \leq \psi_k + C_k$ is seen by using a blow-up to make the singularities of ψ_k divisorial). We take

$$\varphi = \lim_{k \rightarrow +\infty} \tilde{\psi}_k \quad \text{and} \quad T = \alpha + dd^c \varphi.$$

Since $\alpha + dd^c \tilde{\psi}_k \geq -\varepsilon_k \omega$, we get in the limit $T = \alpha + dd^c \varphi \geq 0$. Let (φ_m) be the Bergman approximation sequence of φ . Since $\varphi \leq \tilde{\psi}_\ell \leq \psi_\ell + C_\ell$, Proposition 2 (a) applied with $\lambda = 2m$ shows that $\varphi_m \geq \frac{1}{2mc_\ell} (2mc_\ell - n)_+ \psi_\ell$ where $c_\ell > 0$ is the coefficient of the log singularity of ψ_ℓ . Therefore, if we take $T_{B,k} = \alpha + dd^c \varphi_{m_k}$, we get in the limit $(T_{B,k}) \succeq_W (T_\ell)$. \square

Remark 8 When X is projective algebraic and $\{\alpha\}$ belongs to the Neron-Severi space

$$\text{NS}_{\mathbb{R}}(X) = (H^{1,1}(X, \mathbb{C}) \cap H^2(X, \mathbb{Z})/\text{torsion}) \otimes_{\mathbb{Z}} \mathbb{R},$$

the fiber $\widehat{\mathcal{S}}_{\alpha}(X)$ is essentially an algebraic object. In fact, we could define $\widehat{\mathcal{S}}_{\alpha}(X)$ as the set of suitable equivalence classes of “formal limits” $\lim_{c_1(D) \rightarrow \{\alpha\}} \lim_{k \rightarrow +\infty} \frac{1}{k} \mathfrak{a}_k$ associated with sequences of graded ideals $\mathfrak{a}_k \subset H^0(X, \mathcal{O}_X(kD))$ satisfying the subadditive property $\mathfrak{a}_{k+\ell} \subset \mathfrak{a}_k \mathfrak{a}_{\ell}$, where D are big \mathbb{Q} -divisors whose first Chern classes $c_1(D)$ approximate $\{\alpha\} \in \text{NS}_{\mathbb{R}}(X)$. Many related questions are discussed in the algebraic setting in Lazarfeld’s book [46]. It is nevertheless an interesting point, even in the projective case, that one can “extrapolate” these concepts to all transcendental classes, and get in this way a global space $\widehat{\mathcal{S}}(X)$ which looks well behaved, e.g. semicontinuous, under variation of the complex structure of X .

5.2 Intersection Theory on $\mathcal{S}(X)$ and $\widehat{\mathcal{S}}(X)$

Let X be a compact Kähler n -dimensional manifold equipped with a Kähler metric ω . We consider closed positive $(1, 1)$ -currents $T_j = \alpha_j + dd^c \varphi_j$, $1 \leq j \leq p$. Let us first assume that the functions φ_j have analytic singularities, and let $Z \subset X$ be an analytic set such that the φ_j ’s are locally bounded on $X \setminus Z$. The (p, p) -current

$$\Theta = \mathbf{1}_{X \setminus Z} T_1 \wedge \dots \wedge T_k$$

is well defined on $X \setminus Z$, thanks to Bedford and Taylor [10], and it is a closed positive current there. By [10] such a current does not carry mass on any analytic set, so we can enlarge Z without changing the total mass of Θ . In fact, Θ extends as a closed positive current on the whole of X . To see this, let us take a simultaneous *log resolution* of the T_j ’s, i.e. a modification

$$\mu : \widehat{X} \rightarrow X$$

such that if $\varphi_j = c_j \log \sum_{\ell} |g_{j,\ell}|^2 + O(1)$, then the pull-back of the ideals $(g_{j,\ell})_{\ell}$, namely $\mu^*(g_{j,\ell})_{\ell} = (g_{j,\ell} \circ \mu)_{\ell}$ is a purely divisorial ideal sheaf $\mathcal{O}_{\widehat{X}}(-D_j)$ on \widehat{X} . Let $u_j = 0$ be a local holomorphic equation of the divisor D_j on \widehat{X} . Since $\log \sum_{\ell} |g_{j,\ell}|^2 = \log |u_j|^2 + \log \sum_{\ell} |g_{j,\ell}/u_j|^2 = \log |u_j|^2 + v_j$, where $v_j \in C^{\infty}$ and $dd^c \log |u_j|^2 = [D_j]$ by the Lelong-Poincaré equation, we find

$$\mu^* T_j = \mu^* \alpha_j + dd^c(\varphi_j \circ \mu) = c_j [D_j] + \widehat{T}_j, \quad \text{where } \widehat{T}_j = \mu^* \alpha_j + dd^c \widehat{\varphi}_j \quad (27)$$

and $\widehat{\varphi}_j$ is a locally bounded potential on \widehat{X} such that $\widehat{T}_j \geq 0$. Now, if $E = \mu^{-1}(Z)$, we get

$$\mathbf{1}_{X \setminus Z} T_1 \wedge \dots \wedge T_p = \mu_*(\mathbf{1}_{\widehat{X} \setminus E} \widehat{T}_1 \wedge \dots \wedge \widehat{T}_p) = \mu_*(\widehat{T}_1 \wedge \dots \wedge \widehat{T}_p). \quad (28)$$

Hence the right-hand side defines the desired extension of $\mathbf{1}_{X \setminus Z} T_1 \wedge \dots \wedge T_p$ to X as the direct image of a closed positive current on \widehat{X} carrying no mass on E . An essential point is the following monotonicity lemma – the reader will find a more general version for non-pluripolar products in [2, Theorem 1.16].

Lemma 4 *Assume that we have closed positive $(1, 1)$ -currents with analytic singularities $T_j, T'_j \in \{\alpha_j\}$ with $T_j \preceq T'_j$, $1 \leq j \leq p$, and let $\gamma \geq 0$ be a closed positive smooth $(n - p, n - p)$ -form on X . If Z is an analytic set containing the poles of all T_j and T'_j , we have*

$$\int_X \mathbf{1}_{X \setminus Z} T_1 \wedge \dots \wedge T_p \wedge \gamma \geq \int_X \mathbf{1}_{X \setminus Z} T'_1 \wedge \dots \wedge T'_p \wedge \gamma.$$

Proof We take a log-resolution $\mu : \widehat{X} \rightarrow X$ that works for all T_j and T'_j simultaneously. By (27) and (28), we have $\mu^* T_j = c_j [D_j] + \widehat{T}_j$ where $\widehat{T}_j \geq 0$ has a locally bounded potential on \widehat{X} , and

$$\int_X \mathbf{1}_{X \setminus Z} T_1 \wedge \dots \wedge T_p \wedge \gamma = \int_{\widehat{X}} \widehat{T}_1 \wedge \dots \wedge \widehat{T}_p \wedge \mu^* \gamma.$$

There are of course similar formulas $\mu^* T'_j = c'_j [D'_j] + \widehat{T}'_j$ for the T'_j 's, and our assumption $T_j \preceq T'_j$ means that the corresponding divisors satisfy $c_j D_j \leq c'_j D'_j$, hence $\Delta_j := c'_j D'_j - c_j D_j \geq 0$. In terms of cohomology, we have

$$\mu^* \{\alpha_j\} = \{\mu^* T_j\} = \{\widehat{T}_j\} + \{c_j D_j\} = \{\mu^* T'_j\} = \{\widehat{T}'_j\} + \{c'_j D'_j\},$$

hence $\{\widehat{T}_j\} = \{\widehat{T}'_j\} + \{\Delta_j\}$ in $H^2(\widehat{X}, \mathbb{R})$. By Stokes' theorem, we conclude that

$$\begin{aligned} \int_{\widehat{X}} \widehat{T}_1 \wedge \widehat{T}_2 \wedge \dots \wedge \widehat{T}_p \wedge \mu^* \gamma &= \int_{\widehat{X}} (\widehat{T}'_1 + \{\Delta_1\}) \wedge \widehat{T}_2 \wedge \dots \wedge \widehat{T}_p \wedge \mu^* \gamma \\ &\geq \int_{\widehat{X}} \widehat{T}'_1 \wedge \widehat{T}_2 \wedge \dots \wedge \widehat{T}_p \wedge \mu^* \gamma \end{aligned}$$

thanks to the positivity of our currents $\widehat{T}_j, \widehat{T}'_j$ and the fact that the product of such currents with bounded potentials by the current of integration $[\Delta_j]$ is well defined

and positive ([10]). By replacing successively all terms $\{\widehat{T}_j\}$ by $\{\widehat{T}'_j\} + \{\Delta_j\}$ we infer

$$\int_{\widehat{X}} \widehat{T}_1 \wedge \dots \wedge \widehat{T}_p \wedge \mu^* \gamma \geq \int_{\widehat{X}} \widehat{T}'_1 \wedge \dots \wedge \widehat{T}'_p \wedge \mu^* \gamma. \quad \square$$

Now, assume that we have arbitrary closed positive $(1, 1)$ -currents T_1, \dots, T_p . For each of them, we take a sequence $T_{j,k} = \alpha_j + i\partial\bar{\partial}\psi_{j,k}$ of monotone asymptotically equisingular approximations by currents with analytic singularities, $T_{j,k} \geq -\varepsilon_{j,k}\omega$, $\lim_{k \rightarrow +\infty} \varepsilon_{j,k} = 0$. We have $T_{j,k} \leq T_{j,k+1}$, and we may also assume without loss of generality that $\varepsilon_{j,k} \geq \varepsilon_{j,k+1} > 0$ for all j, k . Let Z_k be an analytic containing all poles of the $T_{j,k}$, $1 \leq j \leq p$. It follows immediately from the above discussion and especially from Lemma 4 that the integrals

$$\int_X \mathbf{1}_{X \setminus Z_k} (T_{1,k} + \varepsilon_{1,k}\omega) \wedge \dots \wedge (T_{p,k} + \varepsilon_{p,k}\omega) \wedge \gamma \geq 0$$

are well defined and nonincreasing in k (the fact that $\varepsilon_{j,k}$ is non increasing even helps here). From this, we conclude

Theorem 15 *For every $p = 1, 2, \dots, n$, there is a well defined p -fold intersection product*

$$\widehat{\mathcal{S}}(X) \times \dots \times \widehat{\mathcal{S}}(X) \longrightarrow H_+^{p,p}(X, \mathbb{R})$$

which assigns to any p -tuple of equivalence classes of monotone sequences $(T_{j,k})$ in $\widehat{\mathcal{S}}(X)$, $1 \leq j \leq p$, the limit cohomology class

$$\lim_{k \rightarrow +\infty} \left\{ \mathbf{1}_{X \setminus Z_k} (T_{1,k} + \varepsilon_{1,k}\omega) \wedge \dots \wedge (T_{p,k} + \varepsilon_{p,k}\omega) \right\} \in H_+^{p,p}(X, \mathbb{R})$$

where $H_+^{p,p}(X, \mathbb{R}) \subset H^{p,p}(X, \mathbb{R})$ denotes the cone of cohomology classes of closed positive (p, p) -currents. This product is additive and homogeneous in each argument in the space $\widehat{\mathcal{S}}(X)$.

Corollary 8 *By combining the above formal intersection product with the Bergman approximation operator $\mathbf{B} : \mathcal{S}(X) \rightarrow \widehat{\mathcal{S}}(X)$, we get an intersection product*

$$\mathcal{S}(X) \times \dots \times \mathcal{S}(X) \longrightarrow H_+^{p,p}(X, \mathbb{R}) \quad \text{denoted} \quad (T_1, \dots, T_p) \longmapsto \langle T_1, \dots, T_p \rangle^+,$$

which is homogeneous and additive in each argument.

Proof (of Theorem 15) The existence of a limit in cohomology is seen by fixing a dual basis $(\{\gamma_j\})$ of $H^{n-p, n-p}(X)$, using the Serre duality pairing

$$H^{p,p}(X, \mathbb{R}) \times H^{n-p, n-p}(X) \rightarrow \mathbb{R}, \quad (\beta, \gamma) \mapsto \int_X \beta \wedge \gamma.$$

Since X is Kähler, we can take $\gamma_1 = \omega^{n-p}$ and replace if necessary γ_j by $\gamma_j + C\omega^{n-p}$, $C \gg 1$, to get $\gamma_j \geq 0$ for all $j \geq 2$. Then the integrals

$$\int_X \mathbf{1}_{X \setminus Z_k} (T_{1,k} + \varepsilon_{1,k}\omega) \wedge \dots \wedge (T_{p,k} + \varepsilon_{p,k}\omega) \wedge \gamma_j \geq 0$$

are nonincreasing in k , and the limit must therefore exist by monotonicity. \square

Remark 9 It is natural to ask how the above intersection product compares with the (cohomology class of the) “non-pluripolar product” $\langle T_1, \dots, T_p \rangle$ defined in [2, § 1]. In fact, the above product only neglects analytic parts of the currents involved. The simple example of a probability measure T without atoms supported on a polar set of a compact Riemann surface X yields e.g. $\langle T \rangle^+ = 1$, while the non-pluripolar part $\langle T \rangle$ vanishes.

5.3 Kähler Definition of the Numerical Dimension

Using the intersection product defined in Theorem 15, we can give a precise definition of the numerical dimension.

Definition 5 Let (X, ω) be a compact Kähler n -dimensional manifold. We define the numerical dimension $\text{nd}(T)$ of a closed positive $(1, 1)$ -current T on X to be the largest integer $p = 0, 1, \dots, n$ such that $\langle T^p \rangle^+ \neq 0$, i.e. $\int_X \langle T^p \rangle^+ \wedge \omega^{n-p} > 0$.

Accordingly, if (L, h) be a pseudoeffective line bundle on X , we define its numerical dimension to be

$$\text{nd}(L, h) = \text{nd}(i \Theta_{L,h}). \tag{29}$$

By the results of the preceding subsection, $\text{nd}(L, h)$ depends only on the weak equivalence class of singularities of the metric h .

Remark 10 H. Tsuji [63] has defined a notion of numerical dimension by a more algebraic method:

Definition 6 Let X be a projective variety and (L, h) a pseudo-effective line bundle. When V runs over all irreducible algebraic suvarieties of X , one defines

$$\nu_{\text{num}}(L, h) = \sup \left\{ p = \dim V ; \limsup_{m \rightarrow \infty} \frac{h^0(\widetilde{V}, \mu^*(L^{\otimes m}) \otimes \mathcal{I}(\mu^*h^m))}{m^p} > 0 \right\}$$

where $\mu : \widetilde{V} \rightarrow V \subset X$ is an embedded desingularization of V in X .

Junyan Cao [14] has shown that $\nu_{\text{num}}(L, h)$ coincides with $\text{nd}(L, h)$ as defined in (29). The idea is to make a reduction to the “big” case $\text{nd}(L, h) = \dim X$ and

to use holomorphic Morse inequalities [20] in combination with a regularization procedure. We omit the rather technical details here.

Remark 11 If L is pseudo-effective there is also a natural concept of numerical dimension $\text{nd}(L)$ that does not depend on the choice of a metric h on L . One can set e.g.

$$\text{nd}(L) = \max \left\{ p \in [0, n]; \exists c > 0, \forall \varepsilon > 0, \exists h_\varepsilon, \Theta_{L, h_\varepsilon} \geq -\varepsilon\omega, \text{ such that} \right. \\ \left. \int_{X \setminus Z_\varepsilon} (i \Theta_{L, h_\varepsilon} + \varepsilon\omega)^p \wedge \omega^{n-p} \geq c \right\},$$

where h_ε runs over all metrics with analytic singularities on L . It may happen in general that $\text{nd}(L, h_{\min}) < \text{nd}(L)$, even when L is nef; in that case the h_ε can be taken to be smooth in the definition of $\text{nd}(L)$, and therefore $\text{nd}(L)$ is the largest integer p such that $c_1(L)^p \neq 0$. In fact, for the line bundle L already mentioned in Remark 5, it is shown in [30] that there is unique positive current $T \in c_1(L)$, namely the current of integration $T = [C]$ on the negative curve $C \subset X$, hence $\text{nd}(L, h_{\min}) = \text{nd}([C]) = 0$, although we have $\text{nd}(L) = 1$ here.

6 Proof of Junyan Cao's Vanishing Theorem

This section is a brief account and a simplified exposition of Junyan Cao's proof, as detailed in his PhD thesis [13]. The key curvature and singularity estimates are contained in the following technical statement, which depends in a crucial way on Bergman regularization and on Yau's theorem [64] for solutions of Monge-Ampère equations.

Proposition 5 *Let (L, h) be a pseudoeffective line bundle on a compact Kähler manifold (X, ω) . Let us write $T = \frac{i}{2\pi} \Theta_{L, h} = \alpha + dd^c \varphi$ where α is smooth and φ is a quasi-psh potential. Let $p = \text{nd}(L, h)$ be the numerical dimension of (L, h) . Then, for every $\gamma \in]0, 1]$ and $\delta \in]0, 1]$, there exists a quasi-psh potential $\Phi_{\gamma, \delta}$ on X satisfying the following properties :*

- (a) $\Phi_{\gamma, \delta}$ is smooth in the complement $X \setminus Z_\delta$ of an analytic set $Z_\delta \subset X$.
- (b) $\alpha + \delta\omega + dd^c \Phi_{\gamma, \delta} \geq \frac{\delta}{2}(1 - \gamma)\omega$ on X .
- (c) $(\alpha + \delta\omega + dd^c \Phi_{\gamma, \delta})^n \geq a \gamma^n \delta^{n-p} \omega^n$ on $X \setminus Z_\delta$.
- (d) $\Phi_{\gamma, \delta} \leq (1 + b\delta)\psi_{B, k} + C_{\gamma, \delta}$ where $\psi_{B, k} \geq \varphi$ is a Bergman approximation of φ of sufficiently high index $k = k_0(\delta)$.
- (e) $\sup_X \Phi_{1, \delta} = 0$, and for all $\gamma \in]0, 1]$ there are estimates $\Phi_{\gamma, \delta} \leq A$ and

$$\exp(-\Phi_{\gamma, \delta}) \leq e^{-(1+b\delta)\varphi} \exp(A - \gamma\Phi_{1, \delta})$$

(f) For $\gamma_0, \delta_0 > 0$ small, $\gamma \in]0, \gamma_0]$, $\delta \in]0, \delta_0]$ and $k = k_0(\delta)$ large enough, we have

$$\mathcal{I}(\Phi_{\gamma,\delta}) = \mathcal{I}_+(\varphi) = \mathcal{I}(\varphi).$$

Here $a, b, A, \gamma_0, \delta_0, C_{\gamma,\delta} > 0$ are suitable constants ($C_{\gamma,\delta}$ being the only one that depends on γ, δ).

Proof Denote by $\psi_{B,k}$ the nonincreasing sequence of Bergman approximations of φ (obtained with denominators $m_k = 2^k$, say). We have $\psi_{B,k} \geq \varphi$ for all k , the $\psi_{B,k}$ have analytic singularities and $\alpha + dd^c \psi_{B,k} \geq -\varepsilon_k \omega$ with $\varepsilon_k \downarrow 0$. Then $\varepsilon_k \leq \frac{\delta}{4}$ for $k \geq k_0(\delta)$ large enough, and so

$$\begin{aligned} \alpha + \delta\omega + dd^c((1 + b\delta)\psi_{B,k}) &\geq \alpha + \delta\omega - (1 + b\delta)(\alpha + \varepsilon_k\omega) \\ &\geq \delta\omega - (1 + b\delta)\varepsilon_k\omega - b\delta\alpha \geq \frac{\delta}{2}\omega \end{aligned}$$

for $b > 0$ small enough (independent of δ and k). Let $\mu : \widehat{X} \rightarrow X$ be a log-resolution of $\psi_{B,k}$, so that

$$\mu^*(\alpha + \delta\omega + dd^c((1 + b\delta)\psi_{B,k})) = c_k[D_k] + \beta_k$$

where $\beta_k \geq \frac{\delta}{2}\mu^*\omega \geq 0$ is a smooth closed $(1, 1)$ -form on \widehat{X} that is > 0 in the complement $\widehat{X} \setminus E$ of the exceptional divisor, $c_k = \frac{1+b\delta}{m_k} > 0$, and D_k is a divisor that includes all components E_ℓ of E . The map μ can be obtained by Hironaka [39] as a composition of a sequence of blow-ups with smooth centers, and we can even achieve that D_k and E are normal crossing divisors. In this circumstance, it is well known that there exist arbitrary small numbers $\eta_\ell > 0$ such that $\beta_k - \sum \eta_\ell[E_\ell]$ is a Kähler class on \widehat{X} . Hence we can find a quasi-psh potential $\widehat{\theta}_k$ on \widehat{X} such that $\widehat{\beta}_k := \beta_k - \sum \eta_\ell[E_\ell] + dd^c\widehat{\theta}_k$ is a Kähler metric on \widehat{X} , and by taking the η_ℓ small enough, we may assume that $\int_{\widehat{X}}(\widehat{\beta}_k)^n \geq \frac{1}{2} \int_{\widehat{X}} \beta_k^n$. Now, we write

$$\begin{aligned} \alpha + \delta\omega + dd^c((1 + b\delta)\psi_{B,k}) &\geq \alpha + \varepsilon_k\omega + dd^c\psi_{B,k} + (\delta - \varepsilon_k)\omega - b\delta(\alpha + \varepsilon_k\omega) \\ &\geq (\alpha + \varepsilon_k\omega + dd^c\psi_{B,k}) + \frac{\delta}{2}\omega \end{aligned}$$

for $k \geq k_0(\delta)$ and $b > 0$ small (independent of δ and k). The assumption on the numerical dimension of $\frac{i}{2\pi}\Theta_{L,h} = \alpha + dd^c\varphi$ implies the existence of a constant $c > 0$ such that, with $Z = \mu(E) \subset X$, we have

$$\begin{aligned} \int_{\widehat{X}} \beta_k^n &= \int_X \mathbf{1}_{X \setminus Z} (\alpha + \delta\omega + dd^c((1 + b\delta)\psi_{B,k}))^n \\ &\geq \binom{n}{p} \left(\frac{\delta}{2}\right)^{n-p} \int_{X \setminus Z} (\alpha + \varepsilon_k\omega + dd^c\psi_{B,k})^p \wedge \omega^{n-p} \geq c \delta^{n-p} \int_X \omega^n \end{aligned}$$

for all $k \geq k_0(\delta)$. Therefore, we may assume

$$\int_{\widehat{X}} (\widehat{\beta}_k)^n \geq \frac{c}{2} \delta^{n-p} \int_X \omega^n.$$

By Yau's theorem [64], there exists a quasi-psh potential $\widehat{\tau}_k$ on \widehat{X} such that $\widehat{\beta}_k + dd^c \widehat{\tau}_k$ is a Kähler metric on \widehat{X} with a prescribed volume form $\widehat{f} > 0$ such that $\int_{\widehat{X}} \widehat{f} = \int_{\widehat{X}} \widehat{\beta}_k^n$. By the above discussion, we can take here $\widehat{f} > \frac{c}{3} \delta^{n-p} \mu^* \omega^n$ everywhere on \widehat{X} .

Now, we consider $\theta_k = \mu_* \widehat{\theta}_k$ and $\tau_k = \mu_* \widehat{\tau}_k \in L_{\text{loc}}^1(X)$. Since $\widehat{\theta}_k$ was defined in such a way that $dd^c \widehat{\theta}_k = \widehat{\beta}_k - \beta_k + \sum_{\ell} \eta_{\ell} [E_{\ell}]$, we get

$$\begin{aligned} \mu^*(\alpha + \delta\omega + dd^c((1 + b\delta)\psi_{B,k} + \gamma(\theta_k + \tau_k))) \\ = c_k[D_k] + (1 - \gamma)\beta_k + \gamma \left(\sum_{\ell} \eta_{\ell} [E_{\ell}] + \widehat{\beta}_k + dd^c \widehat{\tau}_k \right) \geq 0. \end{aligned}$$

This implies in particular that $\Phi_{\gamma,\delta} := (1 + b\delta)\psi_{B,k} + \gamma(\theta_k + \tau_k)$ is a quasi-psh potential on X and that

$$\mu^*(\alpha + \delta\omega + dd^c \Phi_{\gamma,\delta}) \geq (1 - \gamma)\beta_k \geq \frac{\delta}{2} (1 - \gamma) \mu^* \omega,$$

thus condition (b) is satisfied. Putting $Z_{\delta} = \mu(|D_k|) \supset \mu(E) = Z$, we also have

$$\mu^* \mathbf{1}_{X \setminus Z_{\delta}} (\alpha + \delta\omega + dd^c \Phi_{\gamma,\delta})^n \geq \gamma^n \widehat{\beta}_k^n \geq \frac{c}{3} \gamma^n \delta^{n-p} \mu^* \omega^n,$$

therefore condition (c) is satisfied as well with $a = c/3$. Property (a) is clear, and (d) holds since the quasi-psh function $\widehat{\theta}_k + \widehat{\tau}_k$ must be bounded from above on \widehat{X} . We will actually adjust constants in $\widehat{\theta}_k + \widehat{\tau}_k$ (as we may), so that $\sup_X \Phi_{1,\delta} = 0$. Since $\varphi \leq \psi_{B,k} \leq \psi_{B,0} \leq A_0 := \sup_X \psi_{B,0}$ and

$$\Phi_{\gamma,\delta} = (1 + b\delta)\psi_{B,k} + \gamma(\Phi_{1,\delta} - \psi_{B,k}) = (1 - \gamma + b\delta)\psi_{B,k} + \gamma\Phi_{1,\delta},$$

we have

$$(1 + b\delta)\varphi - \gamma(A_0 - \psi_{B,k}) \leq \Phi_{\gamma,\delta} \leq (1 - \gamma + b\delta)A_0$$

and the estimates in (e) follow with $A = (1 + b)A_0$. The only remaining property to be proved is (f). Condition (d) actually implies $\mathcal{S}(\Phi_{\gamma,\delta}) \subset \mathcal{S}((1 + b\delta)\psi_{B,k})$, and Corollary 4 also gives $\mathcal{S}((1 + b\delta)\psi_{B,k}) \subset \mathcal{S}((1 + b\delta/2)\varphi)$ if we take $k \geq k_0(\delta)$ large enough, hence $\mathcal{S}(\Phi_{\gamma,\delta}) \subset \mathcal{S}_+(\varphi)$ for $\delta \leq \delta_0$ small. In the opposite direction, we observe that $\Phi_{1,\gamma}$ satisfies $\alpha + \omega + dd^c \Phi_{1,\delta} \geq 0$ and $\sup_X \Phi_{1,\delta} = 0$, hence $\Phi_{1,\delta}$ belongs to a compact family of quasi-psh functions. A standard result of potential theory then shows the existence of a uniform small constant $c_0 > 0$

such that $\int_X \exp(-c_0 \Phi_{1,\delta}) dV_\omega < +\infty$ for all $\delta \in]0, 1]$. If $f \in \mathcal{O}_{X,x}$ is a germ of holomorphic function and U a small neighborhood of x , the Hölder inequality combined with estimate (e) implies

$$\int_U |f|^2 \exp(-\Phi_{\gamma,\delta}) dV_\omega \leq e^A \left(\int_U |f|^2 e^{-p(1+b\delta)\varphi} dV_\omega \right)^{\frac{1}{p}} \left(\int_U |f|^2 e^{-q\gamma\Phi_{1,\delta}} dV_\omega \right)^{\frac{1}{q}}.$$

We fix $\lambda_0 > 1$ so that $\mathcal{I}(\lambda_0\varphi) = \mathcal{I}_+(\varphi)$, $p \in]1, \lambda_0[$ (say $p = 1 + \lambda_0/2$), and take

$$\gamma \leq \gamma_0 := \frac{c_0}{q} = c_0 \frac{\lambda_0 - 1}{\lambda_0 + 1} \quad \text{and} \quad \delta \leq \delta_0 \in]0, 1] \text{ so small that } p(1 + b\delta_0) \leq \lambda_0.$$

Then clearly $f \in \mathcal{I}(\lambda_0\varphi)$ implies $f \in \mathcal{I}(\Phi_{\gamma,\delta})$, and (f) is proved. □

The rest of the arguments proceeds along the lines of [19, 49] and [28]. Let (L, h) be a pseuffective line bundle and $p = \text{nd}(L, h) = \text{nd}(i \mathcal{O}_{L,h})$. We equip L be the hermitian metric h_δ defined by the quasi-psh weight $\Phi_\delta = \Phi_{\gamma_0,\delta}$ obtained in Proposition 5, with $\delta \in]0, \delta_0]$. Since Φ_δ is smooth on $X \setminus Z_\delta$, the well-known Bochner-Kodaira identity shows that for every smooth (n, q) -form u with values in $K_X \otimes L$ that is compactly supported on $X \setminus Z_\delta$, one has

$$\|\bar{\partial}u\|_\delta^2 + \|\bar{\partial}^*u\|_\delta^2 \geq 2\pi \int_X (\lambda_{1,\delta} + \dots + \lambda_{q,\delta} - q\delta) |u|^2 e^{-\Phi_\delta} dV_\omega,$$

where $\|u\|_\delta^2 := \int_X |u|_{\omega, h_\delta}^2 dV_\omega = \int_X |u|^2 e^{-\Phi_\delta} dV_\omega$ and

$$0 < \lambda_{1,\delta}(x) \leq \dots \leq \lambda_{n,\delta}(x)$$

are, at each point $x \in X$, the eigenvalues of $\alpha + \delta\omega + dd^c \Phi_\delta$ with respect to the base Kähler metric ω . Notice that the $\lambda_{j,\delta}(x) - \delta$ are the actual eigenvalues of $\frac{i}{2\pi} \mathcal{O}_{L, h_\delta} = \alpha + dd^c \Phi_\delta$ with respect to ω and that the inequality $\lambda_{j,\delta}(x) \geq \frac{\delta}{2}(1 - \gamma) > 0$ is guaranteed by Proposition 5 (b). After dividing by $2\pi q$ (and neglecting that constant in the left hand side), we get

$$\|\bar{\partial}u\|_\delta^2 + \|\bar{\partial}^*u\|_\delta^2 + \delta \|u\|_\delta^2 \geq \int_X (\lambda_{1,\delta} + \dots + \lambda_{q,\delta}) |u|^2 e^{-\Phi_\delta} dV_\omega. \quad (30)$$

A standard Hahn-Banach argument in the L^2 -theory of the $\bar{\partial}$ -operator then yields the following conclusion.

Proposition 6 *For every L^2 section of $\Lambda^{n,q} T_X^* \otimes L$ such that $\|f\|_\delta < +\infty$ and $\bar{\partial}f = 0$ in the sense of distributions, there exists a L^2 section $v = v_\delta$ of $\Lambda^{n,q-1} T_X^* \otimes L$ and a*

L^2 section $w = w_\delta$ of $\Lambda^{n,q}T_X^* \otimes L$ such that $f = \bar{\partial}v + w$ with

$$\|v\|_\delta^2 + \frac{1}{\delta}\|w\|_\delta^2 \leq \int_X \frac{1}{\lambda_{1,\delta} + \dots + \lambda_{q,\delta}} |f|^2 e^{-\Phi_\delta} dV_\omega.$$

Because of the singularities of the weight on Z_δ , one should in fact argue first on $X \setminus Z_\delta$ and approximate the base Kähler metric ω by a metric $\widehat{\omega}_{\delta,\varepsilon} = \omega + \varepsilon\widehat{\omega}_\delta$ that is complete on $X \setminus Z_\delta$, exactly as explained in [19]; we omit the (by now standard) details here. A consequence of Proposition 6 is that the “error term” w satisfies the L^2 bound

$$\int_X |w|^2 e^{-\Phi_\delta} dV_\omega \leq \int_X \frac{\delta}{\lambda_{1,\delta} + \dots + \lambda_{q,\delta}} |f|^2 e^{-\Phi_\delta} dV_\omega. \quad (31)$$

The idea for the next estimate is taken from Mourougane’s PhD thesis [49].

Lemma 5 *The ratio $\rho_\delta(x) := \delta/(\lambda_{1,\delta}(x) + \dots + \lambda_{q,\delta}(x))$ is uniformly bounded on X (independently of δ), and, as soon as $q \geq n - \text{nd}(L, h) + 1$, there exists a subsequence (ρ_{δ_ℓ}) , $\delta_\ell \rightarrow 0$, that tends almost everywhere to 0 on X .*

Proof By estimates (b,c) in Proposition 5, we have $\lambda_{j,\delta}(x) \geq \frac{\delta}{2}(1 - \gamma_0)$ and

$$\lambda_{1,\delta}(x) \dots \lambda_{n,\delta}(x) \geq a\gamma_0^n \delta^{n-p} \quad \text{where } p = \text{nd}(L, h). \quad (32)$$

Therefore we already find $\rho_\delta(x) \leq 2/q(1 - \gamma_0)$. Now, we have

$$\int_{X \setminus Z_\delta} \lambda_{n,\delta}(x) dV_\omega \leq \int_X (\alpha + \delta\omega + dd^c\Phi_\delta) \wedge \omega^{n-1} = \int_X (\alpha + \delta\omega) \wedge \omega^{n-1} \leq \text{Const},$$

therefore the “bad set” $S_\varepsilon \subset X \setminus Z_\delta$ of points x where $\lambda_{n,\delta}(x) > \delta^{-\varepsilon}$ has a volume $\text{Vol}(S_\varepsilon) \leq C\delta^\varepsilon$ converging to 0 as $\delta \rightarrow 0$ (with a slightly more elaborate argument we could similarly control any elementary symmetric function in the $\lambda_{j,\delta}$ ’s, but this is not needed here). Outside of S_ε , the inequality (32) yields

$$\lambda_{q,\delta}(x)^q \delta^{-\varepsilon(n-q)} \geq \lambda_{q,\delta}(x)^q \lambda_{n,\delta}(x)^{n-q} \geq a\gamma_0^n \delta^{n-p}$$

hence

$$\lambda_{q,\delta}(x) \geq c\delta^{\frac{n-p+(n-q)\varepsilon}{q}} \quad \text{and} \quad \rho_\delta(x) \leq C\delta^{1 - \frac{n-p+(n-q)\varepsilon}{q}}.$$

If we take $q \geq n - p + 1$ and $\varepsilon > 0$ small enough, the exponent of δ in the final estimate is positive, and Lemma 5 follows. \square

Proof (of Junyan Cao’s Theorem 5) Let $\{f\}$ be a cohomology class in the group $H^q(X, K_X \otimes L \otimes \mathcal{I}_+(h))$, $q \geq n - \text{nd}(L, h) + 1$. Consider a finite Stein open covering $\mathcal{U} = (U_\alpha)_{\alpha=1,\dots,N}$ by coordinate balls U_α . There is an isomorphism between Čech

cohomology $\check{H}^q(\mathcal{U}, \mathcal{F})$ with values in the sheaf $\mathcal{F} = \mathcal{O}(K_X \otimes L) \otimes \mathcal{I}_+(h)$ and the cohomology of the complex $(K_\delta^\bullet, \bar{\partial})$ of (n, q) -forms u such that both u and $\bar{\partial}u$ are L^2 with respect to the weight Φ_δ , i.e. $\int_X |u|^2 \exp(-\Phi_\delta) dV_\omega < +\infty$ and $\int_X |\bar{\partial}u|^2 \exp(-\Phi_\delta) dV_\omega < +\infty$. The isomorphism comes from Leray’s theorem and from the fact that the sheafed complex $(\mathcal{K}_\delta^\bullet, \bar{\partial})$ is a complex of \mathcal{C}^∞ -modules that provides a resolution of the sheaf \mathcal{F} : the main point here is that $\mathcal{I}(\Phi_\delta) = \mathcal{I}_+(\varphi) = \mathcal{I}_+(h)$, as asserted by Proposition 5 (f), and that we can locally solve $\bar{\partial}$ -equations by means of Hörmander’s estimates [40].

Let (ψ_α) be a partition of unity subordinate to \mathcal{U} . The explicit isomorphism between Čech cohomology and L^2 cohomology yields a smooth L^2 representative $f = \sum_{|\alpha|=q} f_\alpha(z) dz_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_I$ which is a combination

$$f = \sum_{\alpha_0} \psi_{\alpha_0} c_{\alpha_0 \alpha_1 \dots \alpha_q} \bar{\partial} \omega_{\alpha_1} \wedge \dots \wedge \bar{\partial} \psi_{\alpha_q}$$

of the components of the corresponding Čech cocycle

$$c_{\alpha_0 \alpha_1 \dots \alpha_q} \in \Gamma(U_{\alpha_0} \cap U_{\alpha_1} \cap \dots \cap U_{\alpha_q}, \mathcal{O}(\mathcal{F})).$$

Estimate (e) in Proposition 5 implies the Hölder inequality

$$\int_X \rho_\delta |f|^2 \exp(-\Phi_\delta) dV_\omega \leq e^A \left(\int_X \rho_\delta^p |f|^2 e^{-p(1+b\delta)\varphi} dV_\omega \right)^{\frac{1}{p}} \left(\int_X |f|^2 e^{-q\gamma_0 \Phi_{1,\delta}} dV_\omega \right)^{\frac{1}{q}}.$$

Our choice of $\delta \leq \delta_0$, γ_0 and p, q shows that the integrals in the right hand side are convergent, and especially $\int_X |f|^2 e^{-p(1+b\delta)\varphi} dV_\omega < +\infty$. Lebesgue’s dominated convergence theorem combined with Lemma 5 implies that the L^p -part goes to 0 as $\delta = \delta_\ell \rightarrow 0$, hence the “error term” w converges to 0 in L^2 norm by estimate (31). If we express the corresponding class $\{w\}$ in Čech cohomology and use Hörmander’s estimates on the intersections $U_\alpha = \bigcap U_{\alpha_j}$, we see that $\{w\}$ will be given by a Čech cocycle (\tilde{w}_α) such that $\int_{U_\alpha} |\tilde{w}_\alpha|^2 e^{-\Phi_\delta} dV_\omega \rightarrow 0$ as $\delta = \delta_\ell \rightarrow 0$ (we may lose here some fixed constants since Φ_δ is just quasi-psh on our balls, but this is irrelevant thanks to the uniform lower bounds for the Hessian). The inequality $\Phi_\delta \leq A$ in Proposition 5 (e) shows that we have as well an unweighted L^2 estimate $\int_{U_\alpha} |\tilde{w}_\alpha|^2 dV \rightarrow 0$. However it is well-known that when one takes unweighted L^2 norms on spaces of Čech cocycles (or uniform convergence on compact subsets, for that purpose), the resulting topology on the finite dimensional space $\check{H}^q(\mathcal{U}, \mathcal{F})$ is Hausdorff, so the subspace of coboundaries is closed in the space of cocycles. Hence we conclude from the above that f is a coboundary, as desired. \square

Remark 12 In this proof, it is remarkable that one can control the error term w , but a priori completely loses control on the element v such that $\bar{\partial}v \approx f$ when $\delta \rightarrow 0$!

7 Compact Kähler Threefolds Without Nontrivial Subvarieties

The bimeromorphic classification of compact Kähler manifolds leads to considering those, termed as “simple” by Campana, that have as little internal structure as possible, and are somehow the elementary bricks needed to reconstruct all others through meromorphic fibrations (cf. [11, 12]).

Definition 7 A compact Kähler manifold X is said to be simple if there does not exist any irreducible analytic subvariety Z with $0 < \dim Z < \dim X$ through a very general point $x \in X$, namely a point x in the complement $X \setminus \bigcup S_j$ of a countable union of analytic sets $S_j \subsetneq X$.

Of course, every one dimensional manifold X is simple, but in higher dimensions $n > 1$, one can show that a very general torus $X = \mathbb{C}^n / \Lambda$ has no nontrivial analytic subvariety Z at all (i.e. none beyond finite sets and X itself), in any dimension n . In even dimension, a very general Hyperkähler manifold can be shown to be simple as well. It has been known since Kodaira that there are no other simple Kähler surfaces (namely only very general 2-dimensional tori and K3 surfaces). Therefore, the next dimension to be investigated is dimension 3. In this case, Campana, Höring and Peternell have shown in [17] that X is bimeromorphically a quotient of a torus by a finite group (see Theorem 18 at the end). Following [16], we give here a short self-contained proof for “strongly simple” Kähler threefolds, namely threefolds that do not possess any proper analytic subvariety.

The simplicity assumption implies that the algebraic dimension is $a(X) = 0$, in particular X cannot be projective, and cannot either be uniruled (i.e. covered by rational curves). By the Kodaira embedding theorem, we also infer that $H^0(X, \Omega_X^2) \neq 0$, otherwise X would be projective. One of the most crucial arguments is the following strong and difficult theorem of Brunella [9].

Theorem 16 ([9]) *Let X be a compact Kähler manifold with a 1-dimensional holomorphic foliation F given by a nonzero morphism of vector bundle $L \rightarrow T_X$, where L is a line bundle on X , and T_X is its holomorphic tangent bundle. If L^{-1} is not pseudoeffective, the closures of the leaves of F are rational curves, and X is thus uniruled.*

We use this result in the form of the following corollary, which has been observed in [41], Proposition 4.2.

Corollary 9 *If X is a non uniruled n -dimensional compact Kähler manifold with $H^0(X, \Omega_X^{n-1}) \neq 0$, then K_X is pseudoeffective.*

Proof Ω_X^{n-1} is canonically isomorphic to $K_X \otimes T_X$. Any nonzero section of Ω_X^{n-1} thus provides a nonzero map $K_X^{-1} \rightarrow T_X$, and an associated foliation. \square

It follows from the above that the canonical line bundle K_X of our simple threefold X must be pseudoeffective. We then use the following simple observation.

Proposition 7 *Assume that X is a strongly simple compact complex manifold. Then every pseudoeffective line bundle (L, h) is nef, and all multiplier sheaves $\mathcal{I}(h^m)$ are trivial, i.e. $\mathcal{I}(h^m) = \mathcal{O}_X$. Moreover, we have $c_1(L)^n = 0$.*

Proof Since there are not positive dimensional analytic subvarieties, the zero varieties of the ideal sheaves $\mathcal{I}(h^m)$ must be finite sets of points, hence, by Skoda [60], the Lelong numbers $\nu(i \Theta_{L,h}, x)$ are zero except on a countable set $S \subset X$. By [21], this implies that L is nef and $c_1(L)^n \geq \sum_{x \in S} \nu(i \Theta_{L,h}, x)^n$. However, by the Grauert-Riemenschneider conjecture solved in [58, 59] and [20], the positivity of $c_1(L)^n$ would imply that $a(X) = n$ (i.e. X Moishezon, a contradiction). Therefore $c_1(L)^n = 0$ and $S = \emptyset$. \square

Proposition 8 *Let X be a compact Kähler manifold of dimension $n > 1$ without any non-trivial subvariety, and with K_X pseudoeffective. Then*

$$h^j(X, K_X^{\otimes m}) \leq h^0(X, \Omega_X^j \otimes K_X^{\otimes m}) \leq \binom{n}{j} \text{ for every } j \geq 0,$$

and the Hilbert polynomial $P(m) := \chi(X, K_X^{\otimes m})$ is constant, equal to $\chi(X, \mathcal{O}_X)$.

Proof The inequality $h^j(X, K_X^{\otimes m}) \leq h^0(X, \Omega_X^j \otimes K_X^{\otimes m})$ follows from the Hard Lefschetz Theorem 4 applied with $L = K_X$ and the corresponding trivial multiplier ideal sheaf. Also, for any holomorphic vector bundle E on X , we have $h^0(X, E) \leq \text{rank}(E)$, otherwise, some ratios of determinants of sections would produce a nonconstant meromorphic function, and thus $a(X) > 0$, contradiction; here we take $E = \Omega_X^j \otimes K_X^{\otimes m}$ and get $\text{rank } E = \binom{n}{j}$. The final claim is clear because a polynomial function $P(m)$ which remains bounded as $m \rightarrow +\infty$ is necessarily constant. \square

Corollary 10 *Let X be a strongly simple Kähler threefold. Let $h^{i,j} = \dim H^{i,j}(X, \mathbb{C})$ be the Hodge numbers. We have*

$$c_1(X)^3 = c_1(X) \cdot c_2(X) = 0, \quad \chi(X, \mathcal{O}_X) = 0 \quad \text{and} \quad q := h^{1,0} > 0.$$

Proof The intersection number $K_X^3 = -c_1(X)^3$ vanishes because it is the leading term of $P(m)$, up to the factor $3!$. The Riemann-Roch formula then gives

$$P(m) = \frac{(1 - 12m)}{24} c_1(X) \cdot c_2(X).$$

The boundedness of $P(m)$ implies $\chi(X, \mathcal{O}_X) = \frac{1}{24} c_1(X) \cdot c_2(X) = 0$. Now, we write

$$0 = \chi(X, \mathcal{O}_X) = 1 - h^{1,0} + h^{2,0} - h^{3,0}.$$

By Kodaira's theorem, $h^{2,0} > 0$ since X is not projective, and $h^{3,0} \leq 1$ since $a(X) = 0$. Thus $0 = 1 - h^{1,0} + h^{2,0} - h^{3,0} \geq 1 - q + 1 - 1 = 1 - q$, and $q > 0$. \square

Everything is now in place for the final conclusion.

Theorem 17 *Let X be a strongly simple Kähler threefold. Then the Albanese map $\alpha : X \rightarrow \text{Alb}(X)$ is a biholomorphism of 3-dimensional tori.*

Proof Since $q = h^{1,0} > 0$, the Albanese map α is non constant. By simplicity, X cannot possess any fibration with positive dimensional fibers, so we must have $\dim \alpha(X) = \dim X = 3$, and as $q = h^{1,0} = h^0(X, \Omega_X^1) \leq 3$ (Proposition 8 with $j = 1, m = 0$) the Albanese map α must be surjective. The function $\det(d\alpha)$ cannot vanish, otherwise we would get a non trivial divisor, so α is étale. Therefore X is a 3-dimensional torus, as a finite étale cover of the 3-dimensional torus $\text{Alb}(X)$, and α must be an isomorphism. \square

In [17], the following stronger result is established as a consequence of the existence of good minimal models for Kähler threefolds:

Theorem 18 *Let X be smooth compact Kähler threefold. If X is simple, there exists a bimeromorphic morphism $X \rightarrow T/G$ where T is a torus and G a finite group acting on T .*

References

1. Bayer, D.A.: The division algorithm and the Hilbert scheme. ProQuest LLC, Ann Arbor (1982). Thesis (Ph.D.)—Harvard University
2. Boucksom, S., Eyssidieux, P., Guedj, V., Zeriahi, A.: Monge-Ampère equations in big cohomology classes. *Acta Math.* **205**(2), 199–262 (2010)
3. Berndtsson, B.: The openness conjecture for plurisubharmonic functions (2013). ArXiv e-prints, math.CV/1305.5781
4. Boucksom, S., Favre, C., Jonsson, M.: Valuations and plurisubharmonic singularities. *Publ. Res. Inst. Math. Sci.* **44**(2), 449–494 (2008)
5. Bierstone, E., Milman, P.D.: Relations among analytic functions. I. *Ann. Inst. Fourier (Grenoble)* **37**(1), 187–239 (1987)
6. Bierstone, E., Milman, P.D.: Uniformization of analytic spaces. *J. Am. Math. Soc.* **2**(4), 801–836 (1989)
7. Boucksom, S.: Cônes positifs des variétés complexes compactes. Ph.D. thesis, Grenoble (2002). <https://tel.archives-ouvertes.fr/tel-00002268/>
8. Boucksom, S.: Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup. (4)* **37**(1), 45–76 (2004)
9. Brunella, M.: Uniformisation of foliations by curves. In: *Holomorphic Dynamical Systems. Volume 1998 of Lecture Notes in Mathematics*, pp. 105–163. Springer, Berlin (2010). Also available from arXiv e-prints, math.CV/0802.4432
10. Bedford, E., Taylor, B.A.: The Dirichlet problem for a complex Monge-Ampère equation. *Invent. Math.* **37**(1), 1–44 (1976)
11. Campana, F.: Algèbricité et compacité dans l'espace des cycles d'un espace analytique complexe. *Math. Ann.* **251**(1), 7–18 (1980)

12. Campana, F.: Réduction d'Albanese d'un morphisme propre et faiblement kählérien. II. Groupes d'automorphismes relatifs. *Compos. Math.* **54**(3), 399–416 (1985)
13. Cao, J.: Théorèmes d'annulation et théorèmes de structure sur les variétés kähleriennes compactes. Ph.D. thesis, Grenoble (2013)
14. Cao, J.: Numerical dimension and a Kawamata-Viehweg-Nadel type vanishing theorem on compact Kähler manifolds. *Compos. Math.* (2014). (to appear, cf. arXiv e-prints: math.AG/1210.5692)
15. Cao, J.: Additivity of the approximation functional of currents induced by Bergman kernels. *C. R. Math. Acad. Sci. Paris* (2015). (to appear, cf. doi:10.1016/j.crma.2014.11.004, arXiv e-prints: math.AG/1410.8288)
16. Campana, F., Demailly, J.-P., Verbitsky, M.: Compact Kähler 3-manifolds without non-trivial subvarieties (2013). ArXiv e-prints, math.CV/1305.5781
17. Campana, F., Höring, A., Peternell, T.: Abundance for Kähler threefolds (2014). ArXiv e-prints, math.AG/1403.3175
18. Demailly, J.-P., Ein, L., Lazarsfeld, R.: A subadditivity property of multiplier ideals. *Michigan Math. J.* **48**, 137–156 (2000). Dedicated to William Fulton on the occasion of his 60th birthday.
19. Demailly, J.-P.: Estimations L^2 pour l'opérateur $\bar{\partial}$ d'un fibré vectoriel holomorphe semi-positif au-dessus d'une variété kählérienne complète. *Ann. Sci. École Norm. Sup. (4)*, **15**(3), 457–511 (1982)
20. Demailly, J.-P.: Champs magnétiques et inégalités de Morse pour la d'' -cohomologie. *Ann. Inst. Fourier (Grenoble)* **35**(4), 189–229 (1985)
21. Demailly, J.-P.: Regularization of closed positive currents and intersection theory. *J. Algebra. Geom.* **1**(3), 361–409 (1992)
22. Demailly, J.-P.: A numerical criterion for very ample line bundles. *J. Differ. Geom.* **37**(2), 323–374 (1993)
23. Demailly, J.-P.: Multiplier ideal sheaves and analytic methods in algebraic geometry. In: *School on Vanishing Theorems and Effective Results in Algebraic Geometry* (Trieste, 2000). Volume 6 of ICTP Lecture Notes, pp 1–148. Abdus Salam International Centre for Theoretical Physics, Trieste (2001)
24. Demailly, J.-P.: Complex analytic and differential geometry, online book at: <http://www-fourier.ujf-grenoble.fr/~demailly/documents.html>. Institut Fourier (2012)
25. de Fernex, T., Ein, L., Mustașă, M.: Bounds for log canonical thresholds with applications to birational rigidity. *Math. Res. Lett.* **10**(2–3), 219–236 (2003)
26. de Fernex, T., Ein, L., Mustașă, M.: Shokurov's ACC conjecture for log canonical thresholds on smooth varieties. *Duke Math. J.* **152**(1), 93–114 (2010)
27. Demailly, J.-P., Kollár, J.: Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds. *Ann. Sci. École Norm. Sup. (4)* **34**(4), 525–556 (2001)
28. Demailly, J.-P., Peternell, T.: A Kawamata-Viehweg vanishing theorem on compact Kähler manifolds. In: *Surveys in Differential Geometry* (Boston, 2002). *Surveys in Differential Geometry*, vol. VIII, pp. 139–169. International Press, Somerville (2003)
29. Demailly, J.-P., Pham, H.H.: A sharp lower bound for the log canonical threshold. *Acta Math.* **212**(1), 1–9 (2014)
30. Demailly, J.-P., Peternell, T., Schneider, M.: Compact complex manifolds with numerically effective tangent bundles. *J. Algebra. Geom.* **3**(2), 295–345 (1994)
31. Demailly, J.-P., Peternell, T., Schneider, M.: Pseudo-effective line bundles on compact Kähler manifolds. *Int. J. Math.* **12**(6), 689–741 (2001)
32. Eisenbud, D.: *Commutative Algebra with a View Toward Algebraic Geometry*. Volume 150 of Graduate Texts in Mathematics. Springer, New York (1995)
33. Enoki, I.: Strong-Lefschetz-type theorem for semi-positive line bundles over compact Kähler manifolds. In: *Geometry and Global Analysis* (Sendai, 1993), pp. 211–212. Tohoku University, Sendai (1993)
34. Favre, C., Jonsson, M.: Valuations and multiplier ideals. *J. Am. Math. Soc.* **18**(3), 655–684 (electronic) (2005)

35. Galligo, A.: Théorème de division et stabilité en géométrie analytique locale. *Ann. Inst. Fourier (Grenoble)* **29**(2), vii, 107–184 (1979)
36. Guan, Q., Zhou, X.: Strong openness conjecture for plurisubharmonic functions (2013). ArXiv e-prints
37. Guan, Q., Zhou, X.: Strong openness conjecture and related problems for plurisubharmonic functions. ArXiv e-prints, math.CV/1401.7158
38. Guan, Q., Zhou, X.: Effectiveness of demailly's strong openness conjecture and related problems (2014). ArXiv e-prints, math.CV/1403.7247
39. Hironaka, H.: Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II. *Ann. Math. (2)* **79**, 109–203; *ibid.* (2), **79**, 205–326 (1964)
40. Hörmander, L.: An Introduction to Complex Analysis in Several Variables. Volume 7 of North-Holland Mathematical Library, 3rd edn. North-Holland, Amsterdam (1990); First edition 1966
41. Höring, A., Peternell, T., Radloff, I.: Uniformisation in dimension four: towards a conjecture of itaka (2011). arXiv e-prints, math.AG/1103.5392, to appear in *Math. Zeitschrift*
42. Jonsson, M., Mustață, M.: Valuations and asymptotic invariants for sequences of ideals. *Ann. Inst. Fourier (Grenoble)* **62**(6):2145–2209 (2012)
43. Jonsson, M., Mustață, M.: An algebraic approach to the openness conjecture of Demailly and Kollár. *J. Inst. Math. Jussieu* **13**(1), 119–144 (2014)
44. Kim, D.: A remark on the approximation of plurisubharmonic functions. *C. R. Math. Acad. Sci. Paris* **352**(5), 387–389 (2014)
45. Kiselman, C.O.: Attenuating the singularities of plurisubharmonic functions. *Ann. Polon. Math.* **60**(2), 173–197 (1994)
46. Lazarsfeld, R.: Positivity in Algebraic Geometry. I, II. Volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer, Berlin (2004)
47. Lempert, L.: Modules of square integrable holomorphic germs (2014). ArXiv e-prints, math.CV/1404.0407
48. Moišezon, B.G.: On n -dimensional compact complex manifolds having n algebraically independent meromorphic functions. I, II, III. *Izv. Akad. Nauk SSSR Ser. Mat.* **30**, 133–174, 345–386, 621–656 (1966)
49. Mourougane, C.: Versions kählériennes du théorème d'annulation de Bogomolov-Sommese. *C. R. Acad. Sci. Paris Sér. I Math.* **321**(11), 1459–1462 (1995)
50. Nadel, A.M.: Multiplier ideal sheaves and Kähler-Einstein metrics of positive scalar curvature. *Ann. Math. (2)*, **132**(3), 549–596 (1990)
51. Ohsawa, T., Takegoshi, K.: On the extension of L^2 holomorphic functions. *Math. Z.* **195**(2), 197–204 (1987)
52. Peternell, T.: Algebraicity criteria for compact complex manifolds. *Math. Ann.* **275**(4), 653–672 (1986)
53. Peternell, T.: Moishezon manifolds and rigidity theorems. *Bayreuth. Math. Schr.* **54**, 1–108 (1998)
54. Phạm, H.H.: The weighted log canonical threshold. *C. R. Math. Acad. Sci. Paris* **352**(4), 283–288 (2014)
55. Phong, D.H., Sturm, J.: On a conjecture of Demailly and Kollár. *Asian J. Math.* **4**(1), 221–226 (2000). Kodaira's issue
56. Richberg, R.: Stetige streng pseudokonvexe Funktionen. *Math. Ann.* **175**, 257–286 (1968)
57. Siu, Y.T.: Analyticity of sets associated to Lelong numbers and the extension of closed positive currents. *Invent. Math.* **27**, 53–156 (1974)
58. Siu, Y.T.: A vanishing theorem for semipositive line bundles over non-Kähler manifolds. *J. Differ. Geom.* **19**(2), 431–452 (1984)
59. Siu, Y.T.: Some recent results in complex manifold theory related to vanishing theorems for the semipositive case. In: *Workshop Bonn 1984 (Bonn, 1984)*. Volume 1111 of *Lecture Notes in Mathematics*, pp. 169–192. Springer, Berlin (1985)
60. Skoda, H.: Application des techniques L^2 à la théorie des idéaux d'une algèbre de fonctions holomorphes avec poids. *Ann. Sci. École Norm. Sup. (4)*, **5**, 545–579 (1972)

61. Skoda, H.: Sous-ensembles analytiques d'ordre fini ou infini dans \mathbb{C}^n . *Bull. Soc. Math. Fr.* **100**, 353–408 (1972)
62. Takegoshi, K.: On cohomology groups of nef line bundles tensorized with multiplier ideal sheaves on compact Kähler manifolds. *Osaka J. Math.* **34**(4), 783–802 (1997)
63. Tsuji, H.: Extension of log pluricanonical forms from subvarieties (2007). ArXiv e-prints
64. Yau, S.T.: On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I. *Commun. Pure Appl. Math.* **31**(3), 339–411 (1978)