

A SIMPLE PROOF OF THE KOBAYASHI CONJECTURE ON THE HYPERBOLICITY OF GENERAL ALGEBRAIC HYPERSURFACES

JEAN-PIERRE DEMAILLY

CONTENTS

0. Introduction	1
1. Hyperbolicity concepts	2
2. Semple tower associated to a directed manifold	5
3. Jet differentials and Green-Griffiths bundles	10
4. Existence of hyperbolic hypersurfaces of low degree	18
5. Proof of the Kobayashi conjecture on the hyperbolicity of general hypersurfaces.....	22
References	33

0. INTRODUCTION

For a compact complex space X , a well known result of Brody [Bro78] asserts that the hyperbolicity property introduced by Kobayashi [Kob67a, Kob67b] is equivalent to the nonexistence of nonconstant entire holomorphic curves $f : \mathbb{C} \rightarrow X$. The aim of this chapter is to describe some geometric techniques that are useful to investigate the existence or nonexistence of such curves. A central conjecture due to Green-Griffiths [GrGr80] and Lang [Lang86] stipulates that for every projective variety X of general type over \mathbb{C} , there exists a proper algebraic subvariety Y of X containing all nonconstant entire curves.

According to Green-Griffiths [GrGr80], jet bundles can be used to give sufficient conditions for Kobayashi hyperbolicity. As in [Dem95], we introduce the formalism of directed varieties and Semple towers [Sem54] to express these conditions in terms of intrinsic algebraic differential equations that entire curves must satisfy; see the “fundamental vanishing theorem” 3.23 below. An important application is a confirmation of an old-standing conjecture of Kobayashi (cf. [Kob70]): a general hypersurface X of complex projective space \mathbb{P}^{n+1} of degree $d \geq d_n$ large enough is Kobayashi hyperbolic. The main arguments are based on techniques introduced in 2016 by Damian Brotbek [Brot17]; they make use of Wronskian differential operators and their associated multiplier ideals. Shortly afterwards, Ya Deng [Deng16] found how to make the method effective, and produced in this way an explicit value of d_n . We describe here a proof based on a simplification of their ideas, producing a very similar bound, namely $d_n = \lfloor \frac{1}{3}(en)^{2n+2} \rfloor$ (cf. [Dem18]). This extends in particular earlier results of Demailly-El Goul [DeEG97], McQuillan [McQ99], Păun [Pau08], Diverio-Merker-Rousseau [DMR10], Diverio-Trapani [DT10] and [Siu15]. According to work of Clemens[86], Zaidenberg [Zai87], Ein [Ein88, Ein91], Voisin [Voi96] and Pacienza [Pac04], every subvariety of a general algebraic hypersurface X of \mathbb{P}^{n+1} is of general type for degrees $d \geq \delta_n$, with an optimal lower bound given by $\delta_n = 2n + 1$ for $2 \leq n \leq 4$ and $\delta_n = 2n$ for $n \geq 5$ – that the same bound $d_n = \delta_n$ holds for Kobayashi hyperbolicity would then be a consequence of the Green-Griffiths-Lang conjecture.

In the same vein, we present a construction of hyperbolic hypersurfaces of \mathbb{P}^{n+1} for all degrees $d \geq 4n^2$. The main idea is inspired from the method of Shiffman-Zaidenberg [ShZa02]; by using again Wronskians, it is possible to give a direct and self-contained argument.

I wish to thank Damian Brotbek, Ya Deng, Simone Diverio, Gianluca Pacienza, Erwan Rousseau, Mihai Păun and Mikhail Zaidenberg for very stimulating discussions on these questions. These notes owe a lot to their work.

1. HYPERBOLICITY CONCEPTS

1.A. KOBAYASHI PSEUDODISTANCE AND PSEUDOMETRIC

We first recall a few basic facts concerning the concept of hyperbolicity, according to S. Kobayashi [Kob67a, Kob67b, Kob70, Kob76]. Let X be a complex space. Given two points $p, q \in X$, let us consider a *chain of analytic disks* from p to q , that is a sequence of holomorphic maps $f_0, f_1, \dots, f_k : \mathbb{D} \rightarrow X$ from the unit disk $\mathbb{D} = D(0, 1) \subset \mathbb{C}$ to X , together with pairs of points $a_0, b_0, \dots, a_k, b_k$ of \mathbb{D} such that

$$p = f_0(a_0), \quad q = f_k(b_k), \quad f_i(b_i) = f_{i+1}(a_{i+1}), \quad i = 0, \dots, k-1.$$

Denoting this chain by α , we define its length $\ell(\alpha)$ to be

$$(1.1') \quad \ell(\alpha) = d_P(a_1, b_1) + \dots + d_P(a_k, b_k),$$

where d_P is the Poincaré distance on \mathbb{D} , and the *Kobayashi pseudodistance* d_X^K on X to be

$$(1.1'') \quad d_X^K(p, q) = \inf_{\alpha} \ell(\alpha).$$

A *Finsler metric* (resp. *pseudometric*) on a vector bundle E is a homogeneous positive (resp. nonnegative) function N on the total space E , that is,

$$N(\lambda\xi) = |\lambda| N(\xi) \quad \text{for all } \lambda \in \mathbb{C} \text{ and } \xi \in E,$$

but in general N is not assumed to be subadditive (i.e. convex) on the fibers of E . A Finsler (pseudo-)metric on E is thus nothing but a hermitian (semi-)norm on the tautological line bundle $\mathcal{O}_{P(E)}(-1)$ of lines of E over the projectivized bundle $Y = P(E)$. The *Kobayashi-Royden infinitesimal pseudometric* on X is the Finsler pseudometric on the tangent bundle T_X defined by

$$(1.2) \quad \mathbf{k}_X(\xi) = \inf \{ \lambda > 0; \exists f : \mathbb{D} \rightarrow X, f(0) = x, \lambda f'(0) = \xi \}, \quad x \in X, \xi \in T_{X,x}.$$

If $\Phi : X \rightarrow Y$ is a morphism of complex spaces, by considering the compositions $\Phi \circ f : \mathbb{D} \rightarrow Y$, this definition immediately implies the monotonicity property $\Phi^* \mathbf{k}_Y \leq \mathbf{k}_X$, i.e.

$$(1.3) \quad \mathbf{k}_Y(\Phi_* \xi) \leq \mathbf{k}_X(\xi) \quad \text{for all } x \in X \text{ and } \xi \in T_{X,x}.$$

When X is a manifold, it follows from the work of H.L. Royden ([Roy71], [Roy74]) that d_X^K is the integrated pseudodistance associated with the pseudometric, i.e.

$$(1.4) \quad d_X^K(p, q) = \inf_{\gamma} \int_{\gamma} \mathbf{k}_X(\gamma'(t)) dt,$$

where the infimum is taken over all piecewise smooth curves joining p to q ; in the case of complex spaces, a similar formula holds, involving jets of analytic curves of arbitrary order, cf. S. Venturini [Ven96]. When X is a non-singular projective variety, it has been shown in [DeLS94] that the Kobayashi pseudodistance and the Kobayashi-Royden infinitesimal pseudometric can be computed by looking only at analytic disks that are contained in algebraic curves.

1.5. Definition. *A complex space X is said to be hyperbolic (in the sense of Kobayashi) if d_X^K is actually a distance, namely if $d_X^K(p, q) > 0$ for all pairs of distinct points (p, q) in X .*

1.B. BRODY CRITERION

In the above context, we have the following well-known result of Brody [Bro78]. Its main interest is to relate hyperbolicity to the non-existence of entire curves.

1.6. Brody reparametrization lemma. *Let ω be a hermitian metric on X and let $f : \mathbb{D} \rightarrow X$ be a holomorphic map. For every $\varepsilon > 0$, there exists a radius $R \geq (1 - \varepsilon)\|f'(0)\|_\omega$ and a homographic transformation ψ of the disk $D(0, R)$ onto $(1 - \varepsilon)\mathbb{D}$ such that*

$$\|(f \circ \psi)'(0)\|_\omega = 1, \quad \|(f \circ \psi)'(t)\|_\omega \leq \frac{1}{1 - |t|^2/R^2} \quad \text{for every } t \in D(0, R).$$

Proof. Select $t_0 \in \mathbb{D}$ such that $(1 - |t|^2)\|f'((1 - \varepsilon)t)\|_\omega$ reaches its maximum for $t = t_0$. The reason for this choice is that $(1 - |t|^2)\|f'((1 - \varepsilon)t)\|_\omega$ is the norm of the differential $f'((1 - \varepsilon)t) : T_{\mathbb{D}} \rightarrow T_X$ with respect to the Poincaré metric $|dt|^2/(1 - |t|^2)^2$ on $T_{\mathbb{D}}$, which is conformally invariant under $\text{Aut}(\mathbb{D})$. One then adjusts R and ψ so that $\psi(0) = (1 - \varepsilon)t_0$ and $|\psi'(0)|\|f'(\psi(0))\|_\omega = 1$. As $|\psi'(0)| = \frac{1-\varepsilon}{R}(1 - |t_0|^2)$, the only possible choice for R is

$$R = (1 - \varepsilon)(1 - |t_0|^2)\|f'(\psi(0))\|_\omega \geq (1 - \varepsilon)\|f'(0)\|_\omega.$$

The inequality for $(f \circ \psi)'$ follows from the fact that the Poincaré norm is maximum at the origin, where it is equal to 1 by the choice of R . Using the Ascoli-Arzelà theorem we obtain immediately:

1.7. Corollary (Brody). *Let (X, ω) be a compact complex hermitian manifold. Given a sequence of holomorphic mappings $f_\nu : \mathbb{D} \rightarrow X$ such that $\lim \|f'_\nu(0)\|_\omega = +\infty$, one can find a sequence of homographic transformations $\psi_\nu : D(0, R_\nu) \rightarrow (1 - 1/\nu)\mathbb{D}$ with $\lim R_\nu = +\infty$, such that, after passing possibly to a subsequence, $(f_\nu \circ \psi_\nu)$ converges uniformly on every compact subset of \mathbb{C} towards a nonconstant holomorphic map $g : \mathbb{C} \rightarrow X$ with $\|g'(0)\|_\omega = 1$ and $\sup_{t \in \mathbb{C}} \|g'(t)\|_\omega \leq 1$.*

An entire curve $g : \mathbb{C} \rightarrow X$ such that $\sup_{\mathbb{C}} \|g'\|_\omega = M < +\infty$ is called a *Brody curve*; this concept does not depend on the choice of ω when X is compact, and one can always assume $M = 1$ by rescaling the parameter t .

1.8. Brody criterion. *Let X be a compact complex manifold. The following properties are equivalent.*

- (a) X is hyperbolic.
- (b) X does not possess any entire curve $f : \mathbb{C} \rightarrow X$.
- (c) X does not possess any Brody curve $g : \mathbb{C} \rightarrow X$.
- (d) The Kobayashi infinitesimal metric \mathbf{k}_X is uniformly bounded below, namely

$$\mathbf{k}_X(\xi) \geq c\|\xi\|_\omega, \quad c > 0,$$

for any hermitian metric ω on X .

When property (b) holds, X is said to be Brody hyperbolic.

Proof. (a) \Rightarrow (b) If X possesses an entire curve $f : \mathbb{C} \rightarrow X$, then by looking at arbitrary large analytic disks $f : D(t_0, R) \subset \mathbb{C}$ and rescaling them on \mathbb{D} as $t \mapsto f(t_0 + Rt)$, it is easy to see that the Kobayashi distance of any two points in $f(\mathbb{C})$ is zero, so X is not hyperbolic.

(b) \Rightarrow (c) is trivial.

(c) \Rightarrow (d) If (d) does not hold, there exists a sequence of tangent vectors $\xi_\nu \in T_{X, x_\nu}$ with $\|\xi_\nu\|_\omega = 1$ and $\mathbf{k}_X(\xi_\nu) \rightarrow 0$. By definition, this means that there exists an analytic curve $f_\nu : \mathbb{D} \rightarrow X$ with $f_\nu(0) = x_\nu$ and $\|f'_\nu(0)\|_\omega \geq (1 - \frac{1}{\nu})/\mathbf{k}_X(\xi_\nu) \rightarrow +\infty$. One can then produce a Brody curve $g : \mathbb{C} \rightarrow X$ by Corollary 1.7, contradicting (c).

(d) \Rightarrow (a). In fact (d) implies after integrating that $d_X^K(p, q) \geq c d_\omega(p, q)$ where d_ω is the geodesic distance associated with ω , so d_X^K must be non degenerate. \square

As a consequence, any projective variety containing a rational curve C (i.e. a curve normalized by $\bar{C} \simeq \mathbb{P}_{\mathbb{C}}^1 \simeq \mathbb{C} \cup \{\infty\}$) or an elliptic curve (i.e. a curve normalized by a nonsingular elliptic curve $\mathbb{C}/(\mathbb{Z} \oplus \mathbb{Z}\tau)$) is non-hyperbolic. An immediate consequence of the Brody criterion is the openness property of hyperbolicity for the metric topology:

1.9. Proposition. *Let $\pi : \mathcal{X} \rightarrow S$ be a holomorphic family of compact complex manifolds. Then the set of $s \in S$ such that the fiber $X_s = \pi^{-1}(s)$ is hyperbolic is open in the metric topology.*

Proof. Let ω be an arbitrary hermitian metric on \mathcal{X} , $(X_{s_\nu})_{s_\nu \in S}$ a sequence of non-hyperbolic fibers, and $s = \lim s_\nu$. By the Brody criterion, one obtains a sequence of entire maps $f_\nu : \mathbb{C} \rightarrow X_{s_\nu}$ such that $\|f'_\nu(0)\|_\omega = 1$ and $\|f'_\nu\|_\omega \leq 1$. Ascoli's theorem shows that there is a subsequence of f_ν converging uniformly to a limit $f : \mathbb{C} \rightarrow X_s$, with $\|f'(0)\|_\omega = 1$. Hence X_s is not hyperbolic and the collection of non-hyperbolic fibers is closed in S . \square

1.C. RELATIONSHIP OF HYPERBOLICITY WITH ALGEBRAIC PROPERTIES

In the case of projective algebraic varieties, Kobayashi hyperbolicity is expected to be an algebraic property. In fact, the following classical conjectures would give a necessary and sufficient algebraic characterization. Recall that a projective variety X of dimension $n = \dim_{\mathbb{C}} X$ is said to be of *general type* if the canonical bundle $K_{\tilde{X}} = \Lambda^n T_{\tilde{X}}^*$ of some desingularization \tilde{X} of X is big. When $n = \dim_{\mathbb{C}} X = 1$, this is equivalent to say that \tilde{X} is not rational or elliptic.

1.10. Some classical conjectures. *Let X be a projective variety.*

- (i) (Green-Griffiths-Lang conjecture) *If X is of general type, there should exist a proper algebraic variety $Y \subsetneq X$ (possibly empty) containing all nonconstant entire curves $f : \mathbb{C} \rightarrow X$.*
- (ii) *Conversely, if X is Kobayashi hyperbolic and nonsingular, it is expected that K_X should be ample. More generally, if X is singular, any desingularization \tilde{X} should be of general type.*
- (iii) (Conjectural algebraic characterization of Kobayashi hyperbolicity). *A projective variety X is Kobayashi hyperbolic if and only if every positive dimensional algebraic subvariety $Y \subset X$ (including X itself) is of general type.*

In fact, since every analytic subspace of Kobayashi hyperbolic space is again hyperbolic by definition, it is not difficult to see by induction on dimension that 1.10 (iii) would follow formally from 1.10 (i) and (ii) [the “if” part is a consequence of 1.10 (i), and the “only if” part follows from 1.10 (ii)]. Thanks to fundamental work of Clemens [Cle86], Ein [Ein88, Ein91] and Voisin [Voi96], it is known that every subvariety Y of a generic algebraic hypersurface $X \subset \mathbb{P}^{n+1}$ of degree $d \geq 2n + 1$ is of general type for $n \geq 2$; Pacienza [Pac04] has even shown that this holds for $d \geq 2n$ when $n \geq 5$. The Green-Griffiths-Lang conjecture would then imply that these hypersurfaces are Kobayashi hyperbolic.

1.11. Definition. *Let X be a projective algebraic manifold, and A a very ample line bundle on X . We say that X is algebraically hyperbolic if there exists $\varepsilon > 0$ such that every closed irreducible curve $C \subset X$ has a normalization \bar{C} such that its Euler characteristic satisfies*

$$-\chi(\bar{C}) = 2g(\bar{C}) - 2 \geq \varepsilon \deg_A(C),$$

where $g(\bar{C})$ is the genus and $\deg_A(C) = C \cdot A = \int_C c_1(A)$.

1.12. Theorem. *Every Kobayashi hyperbolic projective variety is algebraically hyperbolic. More generally, if X is a hyperbolic compact complex manifold equipped with a hermitian metric ω , there exists $\varepsilon > 0$ such that every closed irreducible curve $C \subset X$ satisfies*

$$2g(\bar{C}) - 2 \geq \varepsilon \deg_\omega(C) \quad \text{where } \deg_\omega(C) = \int_C \omega.$$

Proof ([Dem95]). When Γ is a nonsingular compact curve of genus at least 2, the uniformization theorem implies that the universal cover $\rho : \widehat{\Gamma} \rightarrow \Gamma$ is isomorphic to the unit disk \mathbb{D} , and one then sees that the Kobayashi metric \mathbf{k}_Γ is induced by the Kobayashi metric of the disk, i.e.

$$\mathbf{k}_\mathbb{D}^2 = \frac{idz \wedge d\bar{z}}{(1 - |z|^2)^2}.$$

These metrics have constant negative curvature $-\frac{i}{2\pi}\partial\bar{\partial}\log \mathbf{k}_\Gamma^2 = -\frac{1}{\pi}\mathbf{k}_\Gamma^2$, hence

$$\frac{1}{\pi} \int_\Gamma \mathbf{k}_\Gamma^2 = -\chi(\Gamma) = 2g(\Gamma) - 2$$

by the Gauss-Bonnet formula. Now, if X is hyperbolic and $C \subset X$ is a closed analytic curve, the monotonicity formula (1.3) applied to the normalization map $\nu : \overline{C} \rightarrow X$ implies $\mathbf{k}_{\overline{C}} \geq \nu^*\mathbf{k}_X$, and we also have $\mathbf{k}_X^2 \geq c^2\omega$ for some $c > 0$ by 1.8 (d). Therefore

$$2g(\overline{C}) - 2 = \frac{1}{\pi} \int_{\overline{C}} \mathbf{k}_{\overline{C}}^2 \geq \frac{1}{\pi} \int_{\overline{C}} \nu^*\mathbf{k}_X^2 = \frac{1}{\pi} \int_C \mathbf{k}_X^2 \geq \frac{c^2}{\pi} \int_C \omega = \frac{c^2}{\pi} \deg_\omega(C).$$

It is not very difficult to check that the proof can be extended to the case of singular hyperbolic compact complex spaces (a smooth hermitian metric on X being a metric that has extensions with respect to local embeddings of X in open sets $U \subset \mathbb{C}^N$). \square

1.13. Proposition. *Let $\mathcal{X} \rightarrow S$ be an algebraic family of projective algebraic manifolds, given by a projective morphism $\mathcal{X} \rightarrow S$. Then the set of $t \in S$ such that the fiber X_t is algebraically hyperbolic is open with respect to the countable Zariski topology*

Proof. After replacing S by a Zariski open subset, we may assume that the total space \mathcal{X} itself is quasi-projective. Let ω be the Kähler metric on \mathcal{X} obtained by pulling back the Fubini-Study metric via an embedding in a projective space. If integers $d > 0$, $g \geq 0$ are fixed, the set $A_{d,g}$ of $t \in S$ such that X_t contains an algebraic 1-cycle $C = \sum m_j C_j$ with $\deg_\omega(C) = d$ and $g(\overline{C}) = \sum m_j g(\overline{C}_j) \leq g$ is a closed algebraic subset of S (this follows from the existence of a relative cycle space of curves of given degree, and from the fact that the geometric genus is Zariski lower semicontinuous). Now, the set of non algebraically hyperbolic fibers is by definition

$$\bigcap_{k>0} \bigcup_{2g-2<d/k} A_{d,g}.$$

This concludes the proof. \square

It is expected that the concepts of Kobayashi hyperbolicity and algebraic hyperbolicity coincide for projective varieties. This would of course imply that Kobayashi hyperbolicity is an open property with respect to the countable Zariski topology. Combined with the existence of hyperbolic low degree hypersurfaces, such a result would also lead to much improved bounds for the Kobayashi conjecture on generic hyperbolicity.

2. SIMPLE TOWER ASSOCIATED TO A DIRECTED MANIFOLD

2.A. CATEGORY OF DIRECTED VARIETIES

Let us consider a pair (X, V) consisting of a n -dimensional complex manifold X equipped with a linear subspace $V \subset T_X$: assuming X connected, this is by definition an irreducible closed analytic subspace of the total space of T_X such that each fiber $V_x = V \cap T_{X,x}$ is a vector subspace of $T_{X,x}$; the rank $x \mapsto \dim_{\mathbb{C}} V_x$ is Zariski lower semicontinuous, and it may a priori jump.

2.1. Definition. *We will refer to such a pair (X, V) where $V \subset T_X$ is a linear subspace as being a (complex) directed manifold. A morphism $\Phi : (X, V) \rightarrow (Y, W)$ in the category of (complex) directed manifolds is a holomorphic map such that $\Phi_*(V) \subset W$.*

The rank $r \in \{0, 1, \dots, n\}$ of V is by definition the dimension of V_x at a generic point. The dimension may be larger at non generic points; this happens e.g. on $X = \mathbb{C}^n$ for the rank 1 linear space V generated by the Euler vector field: $V_z = \mathbb{C} \sum_{1 \leq j \leq n} z_j \frac{\partial}{\partial z_j}$ for $z \neq 0$, and $V_0 = \mathbb{C}^n$. The absolute situation is the case $V = T_X$ and the relative situation is the case when $V = T_{X/S}$ is the relative tangent space to a smooth holomorphic map $X \rightarrow S$. In general, we can associate to V a sheaf $\mathcal{V} = \mathcal{O}(V) \subset \mathcal{O}(T_X)$ of holomorphic sections. These sections need not generate the fibers of V at singular points, as one sees already in the case of the Euler vector field when $n \geq 2$. However, \mathcal{V} is a *saturated* subsheaf of $\mathcal{O}(T_X)$, i.e. $\mathcal{O}(T_X)/\mathcal{V}$ has no torsion: in fact, if the components of a section have a common divisorial component, one can always simplify this divisor and produce a new section without any such common divisorial component. Instead of defining directed manifolds by picking a linear space V , one could equivalently define them by considering saturated coherent subsheaves $\mathcal{V} \subset \mathcal{O}(T_X)$. One could also take the dual viewpoint, looking at arbitrary quotient morphisms $\Omega_X^1 \rightarrow \mathcal{W} = \mathcal{V}^*$ (and recovering $\mathcal{V} = \mathcal{W}^* = \text{Hom}_{\mathcal{O}}(\mathcal{W}, \mathcal{O})$, as $\mathcal{V} = \mathcal{V}^{**}$ is reflexive). We want to stress here that no assumption need be made on the Lie bracket tensor $[\bullet, \bullet] : \mathcal{V} \times \mathcal{V} \rightarrow \mathcal{O}(T_X)/\mathcal{V}$, i.e. we do not assume any kind of integrability for \mathcal{V} or \mathcal{W} . Even though we will not consider such situations here, one can even generalize the concept of directed structure to the case when X is a singular (say reduced) complex space X . In fact $V|_{X'}$ should then be a holomorphic vector subbundle of $T_{X'}$ on some analytic Zariski open set $X' \subset X_{\text{reg}}$, and if $U \hookrightarrow Z$ is an embedding of an open neighborhood $U \subset X$ of a point $x_0 \in X$ into an open set $Z \subset \mathbb{C}^N$, we demand that the directed structure $V|_U$ be a (closed and analytic) subspace of T_Z , obtained as the closure of $V|_{X' \cap U}$ in T_Z via the obvious “inclusion morphism” $(X' \cap U, V|_{X' \cap U}) \hookrightarrow (Z, T_Z)$. A morphism $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ in the category of directed varieties is the same as a holomorphic curve $t \mapsto f(t)$ that is *tangent to V* , i.e. $f'(t) \in V_{f(t)}$ for all t . The concept of Kobayashi hyperbolicity can be extended to directed varieties as follows.

2.2. Definition. *Let (X, V) be a complex directed manifold. The Kobayashi-Royden infinitesimal metric of (X, V) is the Finsler metric on V defined for any $x \in X$ and $\xi \in V_x$ by*

$$\mathbf{k}_{(X,V)}(\xi) = \inf \{ \lambda > 0; \exists f : (\mathbb{D}, T_{\mathbb{D}}) \rightarrow (X, V), f(0) = x, \lambda f'(0) = \xi \}.$$

We say that (X, V) is *infinitesimally hyperbolic* if $\mathbf{k}_{(X,V)}$ is positive definite on every fiber V_x and satisfies a uniform lower bound $\mathbf{k}_{(X,V)}(\xi) \geq \varepsilon \|\xi\|_{\omega}$ in terms of any smooth hermitian metric ω on X , when x runs over a compact subset of X . When X is compact, the Brody criterion shows that this is equivalent to the nonexistence of nonconstant entire curves $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$, or even to the nonexistence of entire curves $g : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ with $\sup \|g'(t)\|_{\omega} = \|g'(0)\|_{\omega} = 1$. In this context we have the

2.3. Generalized Green-Griffiths-Lang conjecture. *Let (X, V) be a projective directed manifold where $V \subset T_X$ is nonsingular (i.e. a subbundle of T_X). Assume that (X, V) is of “general type” in the sense that $K_V := \det V^*$ is a big line bundle. Then there should exist a proper algebraic subvariety $Y \subsetneq X$ containing the images $f(\mathbb{C})$ of all entire curves $f : \mathbb{C} \rightarrow X$ tangent to V .*

A similar statement can be made when V is singular, but then K_V has to be replaced by a certain (nonnecessarily invertible) rank 1 sheaf of “locally bounded” forms of $\mathcal{O}(\det V^*)$, with respect to a smooth hermitian form ω on T_X . The reader will find a more precise definition in [Dem18].

2.B. THE 1-JET FONCTOR

The basic idea is to introduce a functorial process which produces a new complex directed manifold (\tilde{X}, \tilde{V}) from a given one (X, V) . The new structure (\tilde{X}, \tilde{V}) plays the role of a space of 1-jets over X . First assume that V is *non-singular*. We let

$$(2.4) \quad \tilde{X} = P(V), \quad \tilde{V} \subset T_{\tilde{X}}$$

be the projectivized bundle of lines of V , together with a subbundle \tilde{V} of $T_{\tilde{X}}$ defined as follows: for every point $(x, [v]) \in \tilde{X}$ associated with a vector $v \in V_x \setminus \{0\}$,

$$(2.4') \quad \tilde{V}_{(x,[v])} = \{\xi \in T_{\tilde{X},(x,[v])}; \pi_*\xi \in \mathbb{C}v\}, \quad \mathbb{C}v \subset V_x \subset T_{X,x},$$

where $\pi : \tilde{X} = P(V) \rightarrow X$ is the natural projection and $\pi_* : T_{\tilde{X}} \rightarrow \pi^*T_X$ is its differential. On $\tilde{X} = P(V)$ we have the tautological line bundle $\mathcal{O}_{\tilde{X}}(-1) \subset \pi^*V$ such that $\mathcal{O}_{\tilde{X}}(-1)_{(x,[v])} = \mathbb{C}v$. The bundle \tilde{V} is characterized by the two exact sequences

$$(2.5) \quad 0 \rightarrow T_{\tilde{X}/X} \rightarrow \tilde{V} \xrightarrow{\pi_*} \mathcal{O}_{\tilde{X}}(-1) \rightarrow 0,$$

$$(2.5') \quad 0 \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow \pi^*V \otimes \mathcal{O}_{\tilde{X}}(1) \rightarrow T_{\tilde{X}/X} \rightarrow 0,$$

where $T_{\tilde{X}/X}$ denotes the relative tangent bundle of the fibration $\pi : \tilde{X} \rightarrow X$. The first sequence is a direct consequence of the definition of \tilde{V} , whereas the second is a relative version of the Euler exact sequence describing the tangent bundle of the fibers $P(V_x)$. From these exact sequences we infer

$$(2.6) \quad \dim \tilde{X} = n + r - 1, \quad \text{rank } \tilde{V} = \text{rank } V = r,$$

and by taking determinants we find $\det(T_{\tilde{X}/X}) = \pi^* \det V \otimes \mathcal{O}_{\tilde{X}}(r)$, thus

$$(2.7) \quad \det \tilde{V} = \pi^* \det V \otimes \mathcal{O}_{\tilde{X}}(r - 1).$$

By definition, $\pi : (\tilde{X}, \tilde{V}) \rightarrow (X, V)$ is a morphism of complex directed manifolds. Clearly, our construction is functorial, i.e., for every morphism of directed manifolds $\Phi : (X, V) \rightarrow (Y, W)$, there is a commutative diagram

$$\begin{array}{ccc} (\tilde{X}, \tilde{V}) & \xrightarrow{\pi} & (X, V) \\ \tilde{\Phi} \downarrow & & \downarrow \Phi \\ (\tilde{Y}, \tilde{W}) & \xrightarrow{\pi} & (Y, W), \end{array}$$

where the left vertical arrow is the meromorphic map $P(V) \dashrightarrow P(W)$ induced by the differential $\Phi_* : V \rightarrow \Phi^*W$ ($\tilde{\Phi}$ is actually holomorphic if $\Phi_* : V \rightarrow \Phi^*W$ is injective).

2.C. LIFTING OF CURVES TO THE 1-JET BUNDLE

Suppose that we are given a holomorphic curve $f : D_R \rightarrow X$ parametrized by the disk D_R of centre 0 and radius R in the complex plane, and that f is a tangent curve of the directed manifold, i.e., $f'(t) \in V_{f(t)}$ for every $t \in D_R$. If f is nonconstant, there is a well defined and unique tangent line $[f'(t)]$ for every t , even at stationary points, and the map

$$(2.8) \quad \tilde{f} : D_R \rightarrow \tilde{X}, \quad t \mapsto \tilde{f}(t) := (f(t), [f'(t)])$$

is holomorphic (at a stationary point t_0 , we just write $f'(t) = (t - t_0)^s u(t)$ with $s \in \mathbb{N}^*$ and $u(t_0) \neq 0$, and we define the tangent line at t_0 to be $[u(t_0)]$, hence $\tilde{f}(t) = (f(t), [u(t)])$ near t_0 ; even for $t = t_0$, we still denote $[f'(t_0)] = [u(t_0)]$ for simplicity of notation). By definition $f'(t) \in \mathcal{O}_{\tilde{X}}(-1)_{\tilde{f}(t)} = \mathbb{C}u(t)$, hence the derivative f' defines a section

$$(2.9) \quad f' : T_{D_R} \rightarrow \tilde{f}^* \mathcal{O}_{\tilde{X}}(-1).$$

Moreover $\pi \circ \tilde{f} = f$, therefore

$$\pi_* \tilde{f}'(t) = f'(t) \in \mathbb{C}u(t) \implies \tilde{f}'(t) \in \tilde{V}_{(f(t), u(t))} = \tilde{V}_{\tilde{f}(t)}$$

and we see that \tilde{f} is a tangent trajectory of (\tilde{X}, \tilde{V}) . We say that \tilde{f} is the *canonical lifting* of f to \tilde{X} . Conversely, if $g : D_R \rightarrow \tilde{X}$ is a tangent trajectory of (\tilde{X}, \tilde{V}) , then by definition of \tilde{V} we see that $f = \pi \circ g$ is a tangent trajectory of (X, V) and that $g = \tilde{f}$ (unless g is contained in a vertical fiber $P(V_x)$, in which case f is constant).

For any point $x_0 \in X$, there are local coordinates (z_1, \dots, z_n) on a neighborhood Ω of x_0 such that the fibers $(V_z)_{z \in \Omega}$ can be defined by linear equations

$$(2.10) \quad V_z = \left\{ \xi = \sum_{1 \leq j \leq n} \xi_j \frac{\partial}{\partial z_j}; \xi_j = \sum_{1 \leq k \leq r} a_{jk}(z) \xi_k \text{ for } j = r+1, \dots, n \right\},$$

where (a_{jk}) is a holomorphic $(n-r) \times r$ matrix. It follows that a vector $\xi \in V_z$ is completely determined by its first r components (ξ_1, \dots, ξ_r) , and the affine chart $\xi_j \neq 0$ of $P(V)|_\Omega$ can be described by the coordinate system

$$(2.11) \quad \left(z_1, \dots, z_n; \frac{\xi_1}{\xi_j}, \dots, \frac{\xi_{j-1}}{\xi_j}, \frac{\xi_{j+1}}{\xi_j}, \dots, \frac{\xi_r}{\xi_j} \right).$$

Let $f \simeq (f_1, \dots, f_n)$ be the components of f in the coordinates (z_1, \dots, z_n) (we suppose here R so small that $f(D_R) \subset \Omega$). It should be observed that f is uniquely determined by its initial value x and by the first r components (f_1, \dots, f_r) . Indeed, as $f'(t) \in V_{f(t)}$, we can recover the other components by integrating the system of ordinary differential equations

$$(2.12) \quad f'_j(t) = \sum_{1 \leq k \leq r} a_{jk}(f(t)) f'_k(t), \quad j > r,$$

on a neighborhood of 0, with initial data $f(0) = x$. We denote by $m = m(f, t_0)$ the *multiplicity* of f at any point $t_0 \in D_R$, that is, $m(f, t_0)$ is the smallest integer $m \in \mathbb{N}^*$ such that $f_j^{(m)}(t_0) \neq 0$ for some j . By (2.12), we can always suppose $j \in \{1, \dots, r\}$, for example $f_r^{(m)}(t_0) \neq 0$. Then $f'(t) = (t - t_0)^{m-1} u(t)$ with $u_r(t_0) \neq 0$, and the lifting f is described in the coordinates of the affine chart $\xi_r \neq 0$ of $P(V)|_\Omega$ by

$$(2.13) \quad \tilde{f} \simeq \left(f_1, \dots, f_n; \frac{f'_1}{f'_r}, \dots, \frac{f'_{r-1}}{f'_r} \right).$$

2.D. THE SEMPLE TOWER

Let X be a complex n -dimensional manifold. Following ideas of Green-Griffiths [GrGr80], we let $J_k X \rightarrow X$ be the bundle of k -jets of germs of parametrized curves in X , that is, the set of equivalence classes of holomorphic maps $f : (\mathbb{C}, 0) \rightarrow (X, x)$, with the equivalence relation $f \sim g$ if and only if all derivatives $f^{(j)}(0) = g^{(j)}(0)$ coincide for $0 \leq j \leq k$, when computed in some local coordinate system of X near x . The projection map $J_k X \rightarrow X$ is simply $f \mapsto f(0)$. If (z_1, \dots, z_n) are local holomorphic coordinates on an open set $\Omega \subset X$, the elements f of any fiber $J_k X_x$, $x \in \Omega$, can be seen as \mathbb{C}^n -valued maps

$$f = (f_1, \dots, f_n) : (\mathbb{C}, 0) \rightarrow \Omega \subset \mathbb{C}^n,$$

and they are completely determined by their Taylor expansion of order k at $t = 0$

$$f(t) = x + t f'(0) + \frac{t^2}{2!} f''(0) + \dots + \frac{t^k}{k!} f^{(k)}(0) + O(t^{k+1}).$$

In these coordinates, the fiber $J_k X_x$ can thus be identified with the set of k -tuples of vectors $(\xi_1, \dots, \xi_k) = (f'(0), \dots, f^{(k)}(0)) \in (\mathbb{C}^n)^k$. It follows that $J_k X$ is a holomorphic fiber bundle with typical fiber $(\mathbb{C}^n)^k$ over X (however, $J_k X$ is not a vector bundle for $k \geq 2$, because of the nonlinearity of coordinate changes. According to the philosophy of directed structures, one can also introduce the concept of jet bundle in the general situation of complex directed manifolds. If X is equipped with a holomorphic subbundle $V \subset T_X$, one associates to V a k -jet bundle $J_k V$ as follows.

2.14. Definition. *Let (X, V) be a complex directed manifold. We define $J_k V \rightarrow X$ to be the bundle of k -jets of curves $f : (\mathbb{C}, 0) \rightarrow X$ which are tangent to V , i.e., such that $f'(t) \in V_{f(t)}$ for all t in a neighborhood of 0, together with the projection map $f \mapsto f(0)$ onto X .*

It is easy to check that $J_k V$ is actually a subbundle of $J_k X$. In fact, by using (2.10) and (2.12), we see that the fibers $J_k V_x$ are parametrized by

$$((f'_1(0), \dots, f'_r(0)); (f''_1(0), \dots, f''_r(0)); \dots; (f_1^{(k)}(0), \dots, f_r^{(k)}(0))) \in (\mathbb{C}^r)^k$$

for all $x \in \Omega$, hence $J_k V$ is a locally trivial $(\mathbb{C}^r)^k$ -subbundle of $J_k X$. Alternatively, we can pick a local holomorphic connection ∇ on V such that for any germs $w = \sum_{1 \leq j \leq n} w_j \frac{\partial}{\partial z_j} \in \mathcal{O}(T_{X,x})$ and $v = \sum_{1 \leq \lambda \leq r} v_\lambda e_\lambda \in \mathcal{O}(V)_x$ in a local trivializing frame (e_1, \dots, e_r) of $V|_\Omega$ we have

$$(2.15) \quad \nabla_w v(x) = \sum_{1 \leq j \leq n, 1 \leq \lambda \leq r} w_j \frac{\partial v_\lambda}{\partial z_j} e_\lambda(x) + \sum_{1 \leq j \leq n, 1 \leq \lambda, \mu \leq r} \Gamma_{j\lambda}^\mu(x) w_j v_\lambda e_\mu(x).$$

We can of course take the frame obtained from (2.10) by lifting the vector fields $\partial/\partial z_1, \dots, \partial/\partial z_r$, and the “trivial connection” given by the zero Christoffel symbols $\Gamma = 0$. One then obtains a trivialization $J^k V|_\Omega \simeq V|_\Omega^{\oplus k}$ by considering

$$J_k V_x \ni f \mapsto (\xi_1, \xi_2, \dots, \xi_k) = (\nabla f(0), \nabla^2 f(0), \dots, \nabla^k f(0)) \in V_x^{\oplus k}$$

and computing inductively the successive derivatives $\nabla f(t) = f'(t)$ and $\nabla^s f(t)$ via

$$\nabla^s f = (f^* \nabla)_{d/dt} (\nabla^{s-1} f) = \sum_{1 \leq \lambda \leq r} \frac{d}{dt} \left(\nabla^{s-1} f \right)_\lambda e_\lambda(f) + \sum_{1 \leq j \leq n, 1 \leq \lambda, \mu \leq r} \Gamma_{j\lambda}^\mu(f) f'_j \left(\nabla^{s-1} f \right)_\lambda e_\mu(f).$$

This identification depends of course on the choice of ∇ and cannot be defined globally in general (unless we are in the rare situation where V has a global holomorphic connection. \square)

We now describe a convenient process for constructing “projectivized jet bundles”, which will later appear as natural quotients of our jet bundles $J_k V$ (or rather, as suitable desingularized compactifications of the quotients). Such spaces have already been considered since a long time, at least in the special case $X = \mathbb{P}^2$, $V = T_{\mathbb{P}^2}$ (see Gherardelli [Ghe41], Semple [Sem54]), and they have been mostly used as a tool for establishing enumerative formulas dealing with the order of contact of plane curves (see [Coll88], [CoKe94]); the article [ASS97] is also concerned with such generalizations of jet bundles, as well as [LaTh96] by Laksov and Thorup. One defines inductively the *projectivized k -jet bundle* X_k (or *Semple k -jet bundle*) and the associated subbundle $V_k \subset T_{X_k}$ by

$$(2.16) \quad (X_0, V_0) = (X, V), \quad (X_k, V_k) = (\tilde{X}_{k-1}, \tilde{V}_{k-1}).$$

In other words, (X_k, V_k) is obtained from (X, V) by iterating k -times the lifting construction $(X, V) \mapsto (\tilde{X}, \tilde{V})$ described in § 2.B. By (2.4–2.9), we find

$$(2.17) \quad \dim X_k = n + k(r - 1), \quad \text{rank } V_k = r,$$

together with exact sequences

$$(2.18) \quad 0 \longrightarrow T_{X_k/X_{k-1}} \longrightarrow V_k \xrightarrow{(\pi_k)_*} \mathcal{O}_{X_k}(-1) \longrightarrow 0,$$

$$(2.18') \quad 0 \longrightarrow \mathcal{O}_{X_k} \longrightarrow \pi_k^* V_{k-1} \otimes \mathcal{O}_{X_k}(1) \longrightarrow T_{X_k/X_{k-1}} \longrightarrow 0,$$

where π_k is the natural projection $\pi_k : X_k \rightarrow X_{k-1}$ and $(\pi_k)_*$ its differential. Formula (5.4) yields

$$(2.19) \quad \det V_k = \pi_k^* \det V_{k-1} \otimes \mathcal{O}_{X_k}(r - 1).$$

Every nonconstant tangent trajectory $f : D_R \rightarrow X$ of (X, V) lifts to a well defined and unique tangent trajectory $f_{[k]} : D_R \rightarrow X_k$ of (X_k, V_k) . Moreover, the derivative $f'_{[k-1]}$ gives rise to a section

$$(2.20) \quad f'_{[k-1]} : T_{D_R} \rightarrow f_{[k]}^* \mathcal{O}_{X_k}(-1).$$

In coordinates, one can compute $f_{[k]}$ in terms of its components in the various affine charts (5.9) occurring at each step: we get inductively

$$(2.21) \quad f_{[k]} = (F_1, \dots, F_N), \quad f_{[k+1]} = \left(F_1, \dots, F_N, \frac{F'_{s_1}}{F'_{s_r}}, \dots, \frac{F'_{s_{r-1}}}{F'_{s_r}} \right),$$

where $N = n + k(r - 1)$ and $\{s_1, \dots, s_r\} \subset \{1, \dots, N\}$. If $k \geq 1$, $\{s_1, \dots, s_r\}$ contains the last $r - 1$ indices of $\{1, \dots, N\}$ corresponding to the ‘‘vertical’’ components of the projection $X_k \rightarrow X_{k-1}$, and in general, s_r is an index such that $m(F_{s_r}, 0) = m(f_{[k]}, 0)$, that is, F_{s_r} has the smallest vanishing order among all components F_s (s_r may be vertical or not, and the choice of $\{s_1, \dots, s_r\}$ need not be unique).

By definition, there is a canonical injection $\mathcal{O}_{X_k}(-1) \hookrightarrow \pi_k^* V_{k-1}$, and a composition with the projection $(\pi_{k-1})_*$ (analogue for order $k - 1$ of the arrow $(\pi_k)_*$ in the sequence (2.18)) yields for all $k \geq 2$ a canonical line bundle morphism

$$(2.22) \quad \mathcal{O}_{X_k}(-1) \hookrightarrow \pi_k^* V_{k-1} \xrightarrow{(\pi_k)^*(\pi_{k-1})_*} \pi_k^* \mathcal{O}_{X_{k-1}}(-1),$$

which admits precisely $D_k = P(T_{X_{k-1}/X_{k-2}}) \subset P(V_{k-1}) = X_k$ as its zero divisor (clearly, D_k is a hyperplane subbundle of X_k). Hence we find

$$(2.23) \quad \mathcal{O}_{X_k}(1) = \pi_k^* \mathcal{O}_{X_{k-1}}(1) \otimes \mathcal{O}(D_k).$$

Now, we consider the composition of projections

$$(2.24) \quad \pi_{j,k} = \pi_{j+1} \circ \dots \circ \pi_{k-1} \circ \pi_k : X_k \longrightarrow X_j.$$

Then $\pi_{0,k} : X_k \rightarrow X_0 = X$ is a locally trivial holomorphic fiber bundle over X , and the fibers $X_{k,x} = \pi_{0,k}^{-1}(x)$ are k -stage towers of \mathbb{P}^{r-1} -bundles. Since we have (in both directions) morphisms $(\mathbb{C}^r, T_{\mathbb{C}^r}) \leftrightarrow (X, V)$ of directed manifolds which are bijective on the level of bundle morphisms, the fibers are all isomorphic to a ‘‘universal’’ non-singular projective algebraic variety of dimension $k(r - 1)$ which we will denote by $\mathcal{R}_{r,k}$; it is not hard to see that $\mathcal{R}_{r,k}$ is rational (as will indeed follow from the proof of Theorem 3.11 below).

2.25. Remark. When (X, V) is singular, one can easily extend the construction of the Semple tower by functoriality. In fact, assume that X is a closed analytic subset of some open set $Z \subset \mathbb{C}^N$, and that $X' \subset X$ is a Zariski open subset on which $V|_{X'}$ is a subbundle of $T_{X'}$. Then we consider the injection of the nonsingular directed manifold (X', V') into the absolute structure (Z, W) , $W = T_Z$. This yields an injection $(X'_k, V'_k) \hookrightarrow (Z_k, W_k)$, and we simply define (X_k, V_k) to be the closure of (X'_k, V'_k) into (Z_k, W_k) . It is not hard to see that this is indeed a closed analytic subset of the same dimension $n + k(r - 1)$, where $r = \text{rank } V'$.

3. JET DIFFERENTIALS AND GREEN-GRIFFITHS BUNDLES

3.A. GREEN-GRIFFITHS JET DIFFERENTIALS

We first introduce the concept of jet differentials in the sense of Green-Griffiths [GrGr80]. The goal is to provide an intrinsic geometric description of holomorphic differential equations that a germ of curve $f : (\mathbb{C}, 0) \rightarrow X$ may satisfy. In the sequel, we fix a directed manifold (X, V) and suppose implicitly that all germs of curves f are tangent to V .

Let \mathbb{G}_k be the group of germs of k -jets of biholomorphisms of $(\mathbb{C}, 0)$, that is, the group of germs of biholomorphic maps

$$t \mapsto \varphi(t) = a_1 t + a_2 t^2 + \dots + a_k t^k, \quad a_1 \in \mathbb{C}^*, \quad a_j \in \mathbb{C}, \quad j \geq 2,$$

in which the composition law is taken modulo terms t^j of degree $j > k$. Then \mathbb{G}_k is a k -dimensional nilpotent complex Lie group, which admits a natural fiberwise right action on $J_k V$. The action

consists of reparametrizing k -jets of maps $f : (\mathbb{C}, 0) \rightarrow X$ by a biholomorphic change of parameter $\varphi : (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$, that is, $(f, \varphi) \mapsto f \circ \varphi$. There is an exact sequence of groups

$$1 \rightarrow \mathbb{G}'_k \rightarrow \mathbb{G}_k \rightarrow \mathbb{C}^* \rightarrow 1,$$

where $\mathbb{G}_k \rightarrow \mathbb{C}^*$ is the obvious morphism $\varphi \mapsto \varphi'(0)$, and $\mathbb{G}'_k = [\mathbb{G}_k, \mathbb{G}_k]$ is the group of k -jets of biholomorphisms tangent to the identity. Moreover, the subgroup $\mathbb{H} \simeq \mathbb{C}^*$ of homotheties $\varphi(t) = \lambda t$ is a (non-normal) subgroup of \mathbb{G}_k , and we have a semidirect decomposition $\mathbb{G}_k = \mathbb{G}'_k \rtimes \mathbb{H}$. The corresponding action on k -jets is described in coordinates by

$$\lambda \cdot (f', f'', \dots, f^{(k)}) = (\lambda f', \lambda^2 f'', \dots, \lambda^k f^{(k)}).$$

Following [GrGr80], we introduce the vector bundle $E_{k,m}^{\text{GG}}V^* \rightarrow X$ whose fibers are complex valued polynomials $Q(f', f'', \dots, f^{(k)})$ on the fibers of J_kV , of weighted degree m with respect to the \mathbb{C}^* action defined by H , that is, such that

$$(3.1) \quad Q(\lambda f', \lambda^2 f'', \dots, \lambda^k f^{(k)}) = \lambda^m Q(f', f'', \dots, f^{(k)})$$

for all $\lambda \in \mathbb{C}^*$ and $(f', f'', \dots, f^{(k)}) \in J_kV$. Here we view $(f', f'', \dots, f^{(k)})$ as indeterminates with components

$$((f'_1, \dots, f'_r); (f''_1, \dots, f''_r); \dots; (f^{(k)}_1, \dots, f^{(k)}_r)) \in (\mathbb{C}^r)^k.$$

Notice that the concept of polynomial on the fibers of J_kV makes sense, for all coordinate changes $z \mapsto w = \Psi(z)$ on X induce polynomial transition automorphisms on the fibers of J_kV , given by a formula

$$(3.2) \quad (\Psi \circ f)^{(j)} = \Psi'(f) \cdot f^{(j)} + \sum_{s=2}^{s=j} \sum_{j_1+j_2+\dots+j_s=j} c_{j_1 \dots j_s} \Psi^{(s)}(f) \cdot (f^{(j_1)}, \dots, f^{(j_s)})$$

with suitable integer constants $c_{j_1 \dots j_s}$ (this is easily checked by induction on s). In the case $V = T_X$, we get the bundle of ‘‘absolute’’ jet differentials $E_{k,m}^{\text{GG}}T_X^*$. If $Q \in E_{k,m}^{\text{GG}}V^*$ is decomposed into multihomogeneous components of multidegree $(\ell_1, \ell_2, \dots, \ell_k)$ in $f', f'', \dots, f^{(k)}$ (the decomposition is of course coordinate dependent), these multidegrees must satisfy the relation

$$\ell_1 + 2\ell_2 + \dots + k\ell_k = m.$$

The bundle $E_{k,m}^{\text{GG}}V^*$ will be called the *bundle of jet differentials of order k and weighted degree m* . It is clear from (3.2) that a coordinate change $f \mapsto \Psi \circ f$ transforms every monomial $(f^{(\bullet)})^\ell = (f')^{\ell_1} (f'')^{\ell_2} \dots (f^{(k)})^{\ell_k}$ of partial weighted degree $|\ell|_s := \ell_1 + 2\ell_2 + \dots + s\ell_s$, $1 \leq s \leq k$, into a polynomial $((\Psi \circ f)^{(\bullet)})^\ell$ in $(f', f'', \dots, f^{(k)})$ which has the same partial weighted degree of order s if $\ell_{s+1} = \dots = \ell_k = 0$, and a larger or equal partial degree of order s otherwise. Hence, for each $s = 1, \dots, k$, we get a well defined (i.e., coordinate invariant) decreasing filtration F_s^\bullet on $E_{k,m}^{\text{GG}}V^*$ as follows:

$$(3.3) \quad F_s^p(E_{k,m}^{\text{GG}}V^*) = \left\{ \begin{array}{l} Q(f', f'', \dots, f^{(k)}) \in E_{k,m}^{\text{GG}}V^* \text{ involving} \\ \text{only monomials } (f^{(\bullet)})^\ell \text{ with } |\ell|_s \geq p \end{array} \right\}, \quad \forall p \in \mathbb{N}.$$

The graded terms $\text{Gr}_{k-1}^p(E_{k,m}^{\text{GG}}V^*)$ associated with the filtration $F_{k-1}^p(E_{k,m}^{\text{GG}}V^*)$ are precisely the homogeneous polynomials $Q(f', \dots, f^{(k)})$ whose monomials $(f^{(\bullet)})^\ell$ all have partial weighted degree $|\ell|_{k-1} = p$ (hence their degree ℓ_k in $f^{(k)}$ is such that $m - p = k\ell_k$, and $\text{Gr}_{k-1}^p(E_{k,m}^{\text{GG}}V^*) = 0$ unless k divides $m - p$). The transition automorphisms of the graded bundle are induced by coordinate changes $f \mapsto \Psi \circ f$, and they are described by substituting the arguments of $Q(f', \dots, f^{(k)})$ according to formula (3.2), namely $f^{(j)} \mapsto (\Psi \circ f)^{(j)}$ for $j < k$, and $f^{(k)} \mapsto \Psi'(f) \circ f^{(k)}$ for $j = k$ (when $j = k$, the other terms fall in the next stage F_{k-1}^{p+1} of the filtration). Therefore $f^{(k)}$ behaves as an element of $V \subset T_X$ under coordinate changes. We thus find

$$(3.4) \quad \text{Gr}_{k-1}^{m-k\ell_k}(E_{k,m}^{\text{GG}}V^*) = E_{k-1, m-k\ell_k}^{\text{GG}}V^* \otimes S^{\ell_k}V^*.$$

Combining all filtrations F_s^\bullet together, we find inductively a filtration F^\bullet on $E_{k,m}^{\text{GG}}V^*$ such that the graded terms are

$$(3.5) \quad \text{Gr}^\ell(E_{k,m}^{\text{GG}}V^*) = S^{\ell_1}V^* \otimes S^{\ell_2}V^* \otimes \cdots \otimes S^{\ell_k}V^*, \quad \ell \in \mathbb{N}^k, \quad |\ell|_k = m.$$

The bundles $E_{k,m}^{\text{GG}}V^*$ have other interesting properties. In fact,

$$E_{k,\bullet}^{\text{GG}}V^* := \bigoplus_{m \geq 0} E_{k,m}^{\text{GG}}V^*$$

is in a natural way a bundle of graded algebras (the product is obtained simply by taking the product of polynomials). There are natural inclusions $E_{k,\bullet}^{\text{GG}}V^* \subset E_{k+1,\bullet}^{\text{GG}}V^*$ of algebras, hence $E_{\infty,\bullet}^{\text{GG}}V^* = \bigcup_{k \geq 0} E_{k,\bullet}^{\text{GG}}V^*$ is also an algebra. Moreover, the sheaf of holomorphic sections $\mathcal{O}(E_{\infty,\bullet}^{\text{GG}}V^*)$ admits a canonical derivation D^{GG} given by a collection of \mathbb{C} -linear maps

$$D^{\text{GG}} : \mathcal{O}(E_{k,m}^{\text{GG}}V^*) \rightarrow \mathcal{O}(E_{k+1,m+1}^{\text{GG}}V^*),$$

constructed in the following way. A holomorphic section of $E_{k,m}^{\text{GG}}V^*$ on a coordinate open set $\Omega \subset X$ can be seen as a differential operator on the space of germs $f : (\mathbb{C}, 0) \rightarrow \Omega$ of the form

$$(3.6) \quad Q(f) = \sum_{|\alpha_1|+2|\alpha_2|+\cdots+k|\alpha_k|=m} a_{\alpha_1 \dots \alpha_k}(f) (f')^{\alpha_1} (f'')^{\alpha_2} \cdots (f^{(k)})^{\alpha_k}$$

in which the coefficients $a_{\alpha_1 \dots \alpha_k}$ are holomorphic functions on Ω . Then $D^{\text{GG}}Q$ is given by the formal derivative $(D^{\text{GG}}Q)(f)(t) = d(Q(f))/dt$ with respect to the 1-dimensional parameter t in $f(t)$. For example, in dimension 2, if $Q \in H^0(\Omega, \mathcal{O}(E_{2,4}^{\text{GG}}))$ is the section of weighted degree 4

$$Q(f) = a(f_1, f_2) f_1^3 f_2' + b(f_1, f_2) f_1''^2,$$

we find that $D^{\text{GG}}Q \in H^0(\Omega, \mathcal{O}(E_{3,5}^{\text{GG}}))$ is given by

$$\begin{aligned} (D^{\text{GG}}Q)(f) &= \frac{\partial a}{\partial z_1}(f_1, f_2) f_1^4 f_2' + \frac{\partial a}{\partial z_2}(f_1, f_2) f_1^3 f_2'' + \frac{\partial b}{\partial z_1}(f_1, f_2) f_1' f_1''^2 \\ &+ \frac{\partial b}{\partial z_2}(f_1, f_2) f_2' f_1''^2 + a(f_1, f_2) (3f_1^2 f_1'' f_2' + f_1^3 f_2'') + b(f_1, f_2) 2f_1'' f_1'''. \end{aligned}$$

Associated with the graded algebra bundle $E_{k,\bullet}^{\text{GG}}V^*$, we define an analytic fiber bundle

$$(3.7) \quad X_k^{\text{GG}} := \text{Proj}(E_{k,\bullet}^{\text{GG}}V^*) = (J_k V \setminus \{0\})/\mathbb{C}^*$$

over X , which has weighted projective spaces $\mathbb{P}(1^{[r]}, 2^{[r]}, \dots, k^{[r]})$ as fibers (these weighted projective spaces are singular for $k > 1$, but they only have quotient singularities, see [Dol81]; here $J_k V \setminus \{0\}$ is the set of nonconstant jets of order k ; we refer e.g. to Hartshorne's book [Har77] for a definition of the Proj functor). As such, it possesses a canonical sheaf $\mathcal{O}_{X_k^{\text{GG}}}(1)$ such that $\mathcal{O}_{X_k^{\text{GG}}}(m)$ is invertible when m is a multiple of $\text{lcm}(1, 2, \dots, k)$. Under the natural projection $\pi_k : X_k^{\text{GG}} \rightarrow X$, the direct image $(\pi_k)_* \mathcal{O}_{X_k^{\text{GG}}}(m)$ coincides with polynomials

$$(3.8) \quad P(z; \xi_1, \dots, \xi_k) = \sum_{\alpha_\ell \in \mathbb{N}^r, 1 \leq \ell \leq k} a_{\alpha_1 \dots \alpha_k}(z) \xi_1^{\alpha_1} \cdots \xi_k^{\alpha_k}$$

of weighted degree $|\alpha_1| + 2|\alpha_2| + \cdots + k|\alpha_k| = m$ on $J^k V$ with holomorphic coefficients; in other words, we obtain precisely the sheaf of sections of the bundle $E_{k,m}^{\text{GG}}V^*$ of jet differentials of order k and degree m .

3.9. Proposition. *By construction, if $\pi_k : X_k^{\text{GG}} \rightarrow X$ is the natural projection, we have the direct image formula*

$$(\pi_k)_* \mathcal{O}_{X_k^{\text{GG}}}(m) = \mathcal{O}(E_{k,m}^{\text{GG}}V^*)$$

for all k and m .

3.B. INVARIANT JET DIFFERENTIALS

In the geometric context, we are not really interested in the bundles $(J_k V \setminus \{0\})/\mathbb{C}^*$ themselves, but rather on their quotients $(J_k V \setminus \{0\})/\mathbb{G}_k$ (would such nice complex space quotients exist!). We will see that the Semple bundle X_k constructed in §2.D plays the role of such a quotient. First we introduce a canonical bundle subalgebra of $E_{k,\bullet}^{\text{GG}}V^*$.

3.10. Definition. *We introduce a subbundle $E_{k,m}V^* \subset E_{k,m}^{\text{GG}}V^*$, called the bundle of invariant jet differentials of order k and degree m , defined as follows: $E_{k,m}V^*$ is the set of polynomial differential operators $Q(f', f'', \dots, f^{(k)})$ which are invariant under arbitrary changes of parametrization, i.e., for every $\varphi \in \mathbb{G}_k$*

$$Q((f \circ \varphi)', (f \circ \varphi)'', \dots, (f \circ \varphi)^{(k)}) = \varphi'(0)^m Q(f', f'', \dots, f^{(k)}).$$

Alternatively, $E_{k,m}V^* = (E_{k,m}^{\text{GG}}V^*)^{\mathbb{G}'_k}$ is the set of invariants of $E_{k,m}^{\text{GG}}V^*$ under the action of \mathbb{G}'_k . Clearly, $E_{\infty,\bullet}V^* = \bigcup_{k \geq 0} \bigoplus_{m \geq 0} E_{k,m}V^*$ is a subalgebra of $E_{k,m}^{\text{GG}}V^*$ (observe however that this algebra is not invariant under the derivation D^{GG} , since e.g. $f_j'' = D^{\text{GG}}f_j$ is not an invariant polynomial).

3.11. Theorem. *Suppose that V has rank $r \geq 2$. Let $\pi_{0,k} : X_k \rightarrow X$ be the Semple jet bundles constructed in section 2.B, and let $J_k V^{\text{reg}}$ be the bundle of regular k -jets of maps $f : (\mathbb{C}, 0) \rightarrow X$, that is, jets f such that $f'(0) \neq 0$.*

- (i) *The quotient $J_k V^{\text{reg}}/\mathbb{G}_k$ has the structure of a locally trivial bundle over X , and there is a holomorphic embedding $J_k V^{\text{reg}}/\mathbb{G}_k \hookrightarrow X_k$ over X , which identifies $J_k V^{\text{reg}}/\mathbb{G}_k$ with X_k^{reg} (thus X_k is a relative compactification of $J_k V^{\text{reg}}/\mathbb{G}_k$ over X).*
- (ii) *The direct image sheaf*

$$(\pi_{0,k})_* \mathcal{O}_{X_k}(m) \simeq \mathcal{O}(E_{k,m}V^*)$$

can be identified with the sheaf of holomorphic sections of $E_{k,m}V^$.*

- (iii) *For every $m > 0$, the relative base locus of the linear system $|\mathcal{O}_{X_k}(m)|$ is equal to the set X_k^{sing} of singular k -jets. Moreover, $\mathcal{O}_{X_k}(1)$ is relatively big over X .*

Proof. (i) For $f \in J_k V^{\text{reg}}$, the lifting \tilde{f} is obtained by taking the derivative $(f, [f'])$ without any cancellation of zeroes in f' , hence we get a uniquely defined $(k-1)$ -jet $\tilde{f} : (\mathbb{C}, 0) \rightarrow X$. Inductively, we get a well defined $(k-j)$ -jet $f_{[j]}$ in X_j , and the value $f_{[k]}(0)$ is independent of the choice of the representative f for the k -jet. As the lifting process commutes with reparametrization, i.e., $(f \circ \varphi)^\sim = \tilde{f} \circ \varphi$ and more generally $(f \circ \varphi)_{[k]} = f_{[k]} \circ \varphi$, we conclude that there is a well defined set-theoretic map

$$J_k V^{\text{reg}}/\mathbb{G}_k \rightarrow X_k^{\text{reg}}, \quad f \bmod \mathbb{G}_k \mapsto f_{[k]}(0).$$

This map is better understood in coordinates as follows. Fix coordinates (z_1, \dots, z_n) near a point $x_0 \in X$, such that $V_{x_0} = \text{Vect}(\partial/\partial z_1, \dots, \partial/\partial z_r)$. Let $f = (f_1, \dots, f_n)$ be a regular k -jet tangent to V . Then there exists $i \in \{1, 2, \dots, r\}$ such that $f'_i(0) \neq 0$, and there is a unique reparametrization $t = \varphi(\tau)$ such that $f \circ \varphi = g = (g_1, g_2, \dots, g_n)$ with $g_i(\tau) = \tau$ (we just express the curve as a graph over the z_i -axis, by means of a change of parameter $\tau = f_i(t)$, i.e. $t = \varphi(\tau) = f_i^{-1}(\tau)$). Suppose $i = r$ for the simplicity of notation. The space X_k is a k -stage tower of \mathbb{P}^{r-1} -bundles. In the corresponding inhomogeneous coordinates on these \mathbb{P}^{r-1} 's, the point $f_{[k]}(0)$ is given by the collection of derivatives

$$((g'_1(0), \dots, g'_{r-1}(0)); (g''_1(0), \dots, g''_{r-1}(0)); \dots; (g_1^{(k)}(0), \dots, g_{r-1}^{(k)}(0))).$$

[Recall that the other components (g_{r+1}, \dots, g_n) can be recovered from (g_1, \dots, g_r) by integrating the differential system (5.10)]. Thus the map $J_k V^{\text{reg}}/\mathbb{G}_k \rightarrow X_k$ is a bijection onto X_k^{reg} , and the fibers of these isomorphic bundles can be seen as unions of r affine charts $\simeq (\mathbb{C}^{r-1})^k$, associated with

each choice of the axis z_i used to describe the curve as a graph. The change of parameter formula $\frac{d}{d\tau} = \frac{1}{f'_r(t)} \frac{d}{dt}$ expresses all derivatives $g_i^{(j)}(\tau) = d^j g_i/d\tau^j$ in terms of the derivatives $f_i^{(j)}(t) = d^j f_i/dt^j$

$$(3.12) \quad \begin{aligned} (g'_1, \dots, g'_{r-1}) &= \left(\frac{f'_1}{f'_r}, \dots, \frac{f'_{r-1}}{f'_r} \right); \\ (g''_1, \dots, g''_{r-1}) &= \left(\frac{f''_1 f'_r - f'_r f''_1}{f_r'^3}, \dots, \frac{f''_{r-1} f'_r - f'_r f''_{r-1}}{f_r'^3} \right); \dots; \\ (g_1^{(k)}, \dots, g_{r-1}^{(k)}) &= \left(\frac{f_1^{(k)} f'_r - f'_r f_1^{(k)}}{f_r'^{k+1}}, \dots, \frac{f_{r-1}^{(k)} f'_r - f'_r f_{r-1}^{(k)}}{f_r'^{k+1}} \right) + (\text{order} < k). \end{aligned}$$

Also, it is easy to check that $f_r'^{2k-1} g_i^{(k)}$ is an invariant polynomial in $f', f'', \dots, f^{(k)}$ of total degree $2k-1$, i.e., a section of $E_{k,2k-1}$.

(ii) Since the bundles X_k and $E_{k,m}V^*$ are both locally trivial over X , it is sufficient to identify sections σ of $\mathcal{O}_{X_k}(m)$ over a fiber $X_{k,x} = \pi_{0,k}^{-1}(x)$ with the fiber $E_{k,m}V_x^*$, at any point $x \in X$. Let $f \in J_k V_x^{\text{reg}}$ be a regular k -jet at x . By (6.6), the derivative $f'_{[k-1]}(0)$ defines an element of the fiber of $\mathcal{O}_{X_k}(-1)$ at $f_{[k]}(0) \in X_k$. Hence we get a well defined complex valued operator

$$(3.13) \quad Q(f', f'', \dots, f^{(k)}) = \sigma(f_{[k]}(0)) \cdot (f'_{[k-1]}(0))^m.$$

Clearly, Q is holomorphic on $J_k V_x^{\text{reg}}$ (by the holomorphicity of σ), and the \mathbb{G}_k -invariance condition of Definition 3.10 is satisfied since $f_{[k]}(0)$ does not depend on reparametrization and

$$(f \circ \varphi)'_{[k-1]}(0) = f'_{[k-1]}(0) \varphi'(0).$$

Now, $J_k V_x^{\text{reg}}$ is the complement of a linear subspace of codimension n in $J_k V_x$, hence Q extends holomorphically to all of $J_k V_x \simeq (\mathbb{C}^r)^k$ by Riemann's extension theorem (here we use the hypothesis $r \geq 2$; if $r = 1$, the situation is anyway not interesting since $X_k = X$ for all k). Thus Q admits an everywhere convergent power series

$$Q(f', f'', \dots, f^{(k)}) = \sum_{\alpha_1, \alpha_2, \dots, \alpha_k \in \mathbb{N}^r} a_{\alpha_1 \dots \alpha_k} (f')^{\alpha_1} (f'')^{\alpha_2} \dots (f^{(k)})^{\alpha_k}.$$

The \mathbb{G}_k -invariance (3.10) implies in particular that Q must be multihomogeneous in the sense of (3.1), and thus Q must be a polynomial. We conclude that $Q \in E_{k,m}V_x^*$, as desired.

Conversely, for all w in a neighborhood of any given point $w_0 \in X_{k,x}$, we can find a holomorphic family of germs $f_w : (\mathbb{C}, 0) \rightarrow X$ such that $(f_w)_{[k]}(0) = w$ and $(f_w)'_{[k-1]}(0) \neq 0$ (just take the projections to X of integral curves of (X_k, V_k) integrating a nonvanishing local holomorphic section of V_k near w_0). Then every $Q \in E_{k,m}V_x^*$ yields a holomorphic section σ of $\mathcal{O}_{X_k}(m)$ over the fiber $X_{k,x}$ by putting

$$(3.14) \quad \sigma(w) = Q(f'_w, f''_w, \dots, f_w^{(k)})(0) \cdot ((f_w)'_{[k-1]}(0))^{-m}.$$

(iii) By what we saw in (i)–(ii), every section σ of $\mathcal{O}_{X_k}(m)$ over the fiber $X_{k,x}$ is given by a polynomial $Q \in E_{k,m}V_x^*$, and this polynomial can be expressed on the Zariski open chart $f'_r \neq 0$ of $X_{k,x}^{\text{reg}}$ as

$$(3.15) \quad Q(f', f'', \dots, f^{(k)}) = f_r'^m \widehat{Q}(g', g'', \dots, g^{(k)}),$$

where \widehat{Q} is a polynomial and g is the reparametrization of f such that $g_r(\tau) = \tau$. In fact \widehat{Q} is obtained from Q by substituting $f'_r = 1$ and $f_r^{(j)} = 0$ for $j \geq 2$, and conversely Q can be recovered easily from \widehat{Q} by using the substitutions (3.12).

In this context, the jet differentials $f \mapsto f'_1, \dots, f \mapsto f'_r$ can be viewed as sections of $\mathcal{O}_{X_k}(1)$ on a neighborhood of the fiber $X_{k,x}$. Since these sections vanish exactly on X_k^{sing} , the relative base locus of $\mathcal{O}_{X_k}(m)$ is contained in X_k^{sing} for every $m > 0$. We see that $\mathcal{O}_{X_k}(1)$ is big by considering the

sections of $\mathcal{O}_{X_k}(2k-1)$ associated with the polynomials $Q(f', \dots, f^{(k)}) = f_r^{2k-1} g_i^{(j)}$, $1 \leq i \leq r-1$, $1 \leq j \leq k$; indeed, these sections separate all points in the open chart $f'_r \neq 0$ of $X_{k,x}^{\text{reg}}$.

Now, we check that every section σ of $\mathcal{O}_{X_k}(m)$ over $X_{k,x}$ must vanish on $X_{k,x}^{\text{sing}}$. Pick an arbitrary element $w \in X_k^{\text{sing}}$ and a germ of curve $f : (\mathbb{C}, 0) \rightarrow X$ such that $f_{[k]}(0) = w$, $f'_{[k-1]}(0) \neq 0$ and $s = m(f, 0) \gg 0$ (such an f exists by Corollary 6.14). There are local coordinates (z_1, \dots, z_n) on X such that $f(t) = (f_1(t), \dots, f_n(t))$ where $f_r(t) = t^s$. Let Q, \widehat{Q} be the polynomials associated with σ in these coordinates and let $(f')^{\alpha_1} (f'')^{\alpha_2} \dots (f^{(k)})^{\alpha_k}$ be a monomial occurring in Q , with $\alpha_j \in \mathbb{N}^r$, $|\alpha_j| = \ell_j$, $\ell_1 + 2\ell_2 + \dots + k\ell_k = m$. Putting $\tau = t^s$, the curve $t \mapsto f(t)$ becomes a Puiseux expansion $\tau \mapsto g(\tau) = (g_1(\tau), \dots, g_{r-1}(\tau), \tau)$ in which g_i is a power series in $\tau^{1/s}$, starting with exponents of τ at least equal to 1. The derivative $g^{(j)}(\tau)$ may involve negative powers of τ , but the exponent is always $\geq 1 + \frac{1}{s} - j$. Hence the Puiseux expansion of $\widehat{Q}(g', g'', \dots, g^{(k)})$ can only involve powers of τ of exponent $\geq -\max_{\ell}((1 - \frac{1}{s})\ell_2 + \dots + (k-1 - \frac{1}{s})\ell_k)$. Finally $f'_r(t) = st^{s-1} = s\tau^{1-1/s}$, thus the lowest exponent of τ in $Q(f', \dots, f^{(k)})$ is at least equal to

$$\begin{aligned} & \left(1 - \frac{1}{s}\right)m - \max_{\ell} \left(\left(1 - \frac{1}{s}\right)\ell_2 + \dots + \left(k-1 - \frac{1}{s}\right)\ell_k \right) \\ & \geq \min_{\ell} \left(1 - \frac{1}{s}\right)\ell_1 + \left(1 - \frac{1}{s}\right)\ell_2 + \dots + \left(1 - \frac{k-1}{s}\right)\ell_k, \end{aligned}$$

where the minimum is taken over all monomials $(f')^{\alpha_1} (f'')^{\alpha_2} \dots (f^{(k)})^{\alpha_k}$, $|\alpha_j| = \ell_j$, occurring in Q . Choosing $s \geq k$, we already find that the minimal exponent is positive, hence $Q(f', \dots, f^{(k)})(0) = 0$ and $\sigma(w) = 0$ by (3.14). \square

Theorem 3.11 (iii) shows that $\mathcal{O}_{X_k}(1)$ is never relatively ample over X for $k \geq 2$. In order to overcome this difficulty, we define for every $a_{\bullet} = (a_1, \dots, a_k) \in \mathbb{Z}^k$ a line bundle $\mathcal{O}_{X_k}(a_{\bullet})$ on X_k such that

$$(3.16) \quad \mathcal{O}_{X_k}(a_{\bullet}) = \pi_{1,k}^* \mathcal{O}_{X_1}(a_1) \otimes \pi_{2,k}^* \mathcal{O}_{X_2}(a_2) \otimes \dots \otimes \mathcal{O}_{X_k}(a_k).$$

By (6.9), we have $\pi_{j,k}^* \mathcal{O}_{X_j}(1) = \mathcal{O}_{X_k}(1) \otimes \mathcal{O}_{X_k}(-\pi_{j+1,k}^* D_{j+1} - \dots - D_k)$, thus by putting $D_j^* = \pi_{j+1,k}^* D_{j+1}$ for $1 \leq j \leq k-1$ and $D_k^* = 0$, we find an identity

$$(3.17) \quad \begin{aligned} \mathcal{O}_{X_k}(a_{\bullet}) &= \mathcal{O}_{X_k}(b_k) \otimes \mathcal{O}_{X_k}(-b_{\bullet} \cdot D^*), \quad \text{where} \\ b_{\bullet} &= (b_1, \dots, b_k) \in \mathbb{Z}^k, \quad b_j = a_1 + \dots + a_j, \\ b_{\bullet} \cdot D^* &= \sum_{1 \leq j \leq k-1} b_j \pi_{j+1,k}^* D_{j+1}. \end{aligned}$$

In particular, if $b_{\bullet} \in \mathbb{N}^k$, i.e., $a_1 + \dots + a_j \geq 0$, we get a morphism

$$(3.18) \quad \mathcal{O}_{X_k}(a_{\bullet}) = \mathcal{O}_{X_k}(b_k) \otimes \mathcal{O}_{X_k}(-b_{\bullet} \cdot D^*) \rightarrow \mathcal{O}_{X_k}(b_k).$$

The following result gives a sufficient condition for the relative nefness or ampleness of weighted jet bundles.

3.19. Proposition. *Take a very ample line bundle A on X , and consider on X_k the line bundle*

$$L_k = \mathcal{O}_{X_k}(3^{k-1}, 3^{k-2}, \dots, 3, 1) \otimes \pi_{k,0}^* A^{\otimes 3^k}$$

defined inductively by $L_0 = A$ and $L_k = \mathcal{O}_{X_k}(1) \otimes \pi_{k,k-1}^ L_{k-1}^{\otimes 3}$. Then $V_k^* \otimes L_k^{\otimes 2}$ is a nef vector bundle on X_k , which is in fact generated by its global sections, for all $k \geq 0$. Equivalently*

$$L'_k = \mathcal{O}_{X_k}(1) \otimes \pi_{k,k-1}^* L_{k-1}^{\otimes 2} = \mathcal{O}_{X_k}(2 \cdot 3^{k-2}, 2 \cdot 3^{k-3}, \dots, 6, 2, 1) \otimes \pi_{k,0}^* A^{\otimes 2 \cdot 3^{k-1}}$$

is nef over X_k (and generated by sections) for all $k \geq 1$.

Let us recall that a line bundle $L \rightarrow X$ on a projective variety X is said to nef if $L \cdot C \geq 0$ for all irreducible algebraic curves $C \subset X$, and that a vector bundle $E \rightarrow X$ is said to be nef if $\mathcal{O}_{\mathbb{P}(E)}(1)$ is nef on $\mathbb{P}(E) := P(E^*)$; any vector bundle generated by global sections is nef

(cf. [DePS94] for more details). The statement concerning L'_k is obtained by projectivizing the vector bundle $E = V_{k-1}^* \otimes L_{k-1}^{\otimes 2}$ on X_{k-1} , whose associated tautological line bundle is $\mathcal{O}_{\mathbb{P}(E)}(1) = L'_k$ on $\mathbb{P}(E) = P(V_{k-1}) = X_k$. Also one gets inductively that

$$(3.20) \quad L_k = \mathcal{O}_{\mathbb{P}(V_{k-1} \otimes L_{k-1}^{\otimes 2})}(1) \otimes \pi_{k,k-1}^* L_{k-1} \quad \text{is very ample on } X_k.$$

Proof. Let $X \subset \mathbb{P}^N$ be the embedding provided by A , so that $A = \mathcal{O}_{\mathbb{P}^N}(1)|_X$. As is well known, if Q is the tautological quotient vector bundle on \mathbb{P}^N , the twisted cotangent bundle

$$T_{\mathbb{P}^N}^* \otimes \mathcal{O}_{\mathbb{P}^N}(2) = \Lambda^{N-1} Q$$

is nef; hence its quotients $T_X^* \otimes A^{\otimes 2}$ and $V_0^* \otimes L_0^{\otimes 2} = V^* \otimes A^{\otimes 2}$ are nef (any tensor power of nef vector bundles is nef, and so is any quotient). We now proceed by induction, assuming $V_{k-1}^* \otimes L_{k-1}^{\otimes 2}$ to be nef, $k \geq 1$. By taking the second wedge power of the central term in (6.4'), we get an injection

$$0 \longrightarrow T_{X_k/X_{k-1}} \longrightarrow \Lambda^2(\pi_k^* V_{k-1} \otimes \mathcal{O}_{X_k}(1)).$$

By dualizing and twisting with $\mathcal{O}_{X_{k-1}}(2) \otimes \pi_k^* L_{k-1}^{\otimes 2}$, we find a surjection

$$\pi_k^* \Lambda^2(V_{k-1}^* \otimes L_{k-1}) \longrightarrow T_{X_k/X_{k-1}}^* \otimes \mathcal{O}_{X_k}(2) \otimes \pi_k^* L_{k-1}^{\otimes 2} \longrightarrow 0.$$

By the induction hypothesis, we see that $T_{X_k/X_{k-1}}^* \otimes \mathcal{O}_{X_k}(2) \otimes \pi_k^* L_{k-1}^{\otimes 2}$ is nef. Next, the dual of (6.4) yields an exact sequence

$$0 \longrightarrow \mathcal{O}_{X_k}(1) \longrightarrow V_k^* \longrightarrow T_{X_k/X_{k-1}}^* \longrightarrow 0.$$

As an extension of nef vector bundles is nef, the nefness of $V_k^* \otimes L_k^{\otimes 2}$ will follow if we check that $\mathcal{O}_{X_k}(1) \otimes L_k^{\otimes 2}$ and $T_{X_k/X_{k-1}}^* \otimes L_k^{\otimes 2}$ are both nef. However, this follows again from the induction hypothesis if we observe that the latter implies

$$L_k \geq \pi_{k,k-1}^* L_{k-1} \quad \text{and} \quad L_k \geq \mathcal{O}_{X_k}(1) \otimes \pi_{k,k-1}^* L_{k-1}$$

in the sense that $L'' \geq L'$ if the ‘‘difference’’ $L'' \otimes (L')^{-1}$ is nef. All statements remain valid if we replace ‘‘nef’’ with ‘‘generated by sections’’ in the above arguments. \square

3.21. Corollary. *A \mathbb{Q} -line bundle $\mathcal{O}_{X_k}(a_\bullet) \otimes \pi_{k,0}^* A^{\otimes p}$, $a_\bullet \in \mathbb{Q}^k$, $p \in \mathbb{Q}$, is nef (resp. ample) on X_k as soon as*

$$a_j \geq 3a_{j+1} \text{ for } j = 1, 2, \dots, k-2 \text{ and } a_{k-1} \geq 2a_k \geq 0, p \geq 2 \sum a_j,$$

resp.

$$a_j \geq 3a_{j+1} \text{ for } j = 1, 2, \dots, k-2 \text{ and } a_{k-1} > 2a_k > 0, p > 2 \sum a_j.$$

Proof. This follows easily by taking convex combinations of the L_j and L'_j and applying Proposition 3.19 and our observation (3.20). \square

3.22. Remark. As \mathbb{G}_k is a non-reductive group, it is a priori unclear whether the graded ring $\mathcal{A}_{n,k,r} = \bigoplus_{m \in \mathbb{Z}} E_{k,m} V^*$ (taken pointwise over X) is finitely generated. This can be checked manually ([Dem07a], [Dem07b]) for $n = 2$ and $k \leq 4$. Rousseau [Rou06] also checked the case $n = 3$, $k = 3$, and then Merker [Mer08, Mer10] proved the finiteness for $n = 2, 3, 4$, $k \leq 4$ and $n = 2$, $k = 5$. Recently, Bérczi and Kirwan [BeKi12] made an attempt to prove the finiteness in full generality, but it appears that the general case is still unsettled.

3.C. FUNDAMENTAL VANISHING THEOREM

We prove here a fundamental vanishing theorem due to Siu and Yeung ([SiYe96, SiYe97], [Siu97]). Their original proof makes use of Nevanlinna theory, especially of the logarithmic derivative lemma, see also [Dem97] for a more detailed account (in French). An alternative simpler proof based on the Ahlfors lemma and on algebraic properties of jet differentials can be found in [Dem18] (cf. also [Dem95]).

3.23. Fundamental vanishing theorem. *Let (X, V) be a projective directed manifold and A an ample divisor on X . Then $P(f; f', f'', \dots, f^{(k)}) = 0$ for every entire curve $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ and every global section $P \in H^0(X, E_{k,m}^{\text{GG}} V^* \otimes \mathcal{O}(-A))$.*

Proof. We first give a proof of 3.23 in the special case where f is a Brody curve, i.e. $\sup_{t \in \mathbb{C}} \|f'(t)\|_{\omega} < +\infty$ with respect to a given Hermitian metric ω on X . In fact, the proof is much simpler in that case, and thanks to the Brody criterion 1.8, this is sufficient to establish the hyperbolicity of (X, V) . After raising P to a power P^s and replacing $\mathcal{O}(-A)$ with $\mathcal{O}(-sA)$, one can always assume that A is a very ample divisor. We interpret $E_{k,m}^{\text{GG}} V^* \otimes \mathcal{O}(-A)$ as the bundle of complex valued differential operators whose coefficients $a_{\alpha}(z)$ vanish along A .

Fix a finite open covering of X by coordinate balls $B(p_j, R_j)$ such that the balls $B_j(p_j, R_j/4)$ still cover X . As f' is bounded, there exists $\delta > 0$ such that for $f(t_0) \in B(p_j, R_j/4)$ we have $f(t) \in B(p_j, R_j/2)$ whenever $|t - t_0| < \delta$, uniformly for every $t_0 \in \mathbb{C}$. The Cauchy inequalities applied to the components of f in each of the balls imply that the derivatives $f^{(j)}(t)$ are bounded on \mathbb{C} , and therefore, since the coefficients $a_{\alpha}(z)$ of P are also uniformly bounded on each of the balls $B(p_j, R_j/2)$ we conclude that $g := P(f; f', f'', \dots, f^{(k)})$ is a bounded holomorphic function on \mathbb{C} . After moving A in the linear system $|A|$, we may further assume that $\text{Supp } A$ intersects $f(\mathbb{C})$. Then g vanishes somewhere, hence $g \equiv 0$ by Liouville's theorem, as expected.

Next we consider the case where $P \in H^0(X, E_{k,m} V^* \otimes \mathcal{O}(-A))$ is an invariant differential operator. We may of course assume $P \neq 0$. Then we get an associated non-zero section $\sigma \in H^0(X_k, \mathcal{O}_{X_k}(m) \otimes \pi_{k,0}^* \mathcal{O}(-A))$. Thanks to Corollary 3.21, the line bundle

$$L = \mathcal{O}_{X_k}(a_{\bullet}) \otimes \pi_{k,0}^* \mathcal{O}(pA) = \mathcal{O}_{X_k}(m') \otimes \mathcal{O}_{X_k}(-b_{\bullet} \cdot D^*) \otimes \pi_{k,0}^* \mathcal{O}(pA)$$

is ample on X_k for suitable $b_{\bullet} \geq 0$ and $m', p > 0$. Let h_L be a smooth metric on L such that $\omega_k = \Theta_{L, h_L}$ is a Kähler metric on X_k . Then we can produce a singular hermitian metric h on $\mathcal{O}_{X_k}(-1)$ by putting

$$\|\xi\|_h = (\|\sigma^p \cdot \xi^{pm+m'}\|_{h_L^{-1}})^{1/(pm+m')}, \xi \in \mathcal{O}_{X_k}(-1),$$

and viewing $\sigma^p \cdot \xi^{pm+m'}$ as an element in $\mathcal{O}_{X_k}(-m') \otimes \pi_{k,0}^* \mathcal{O}(-pA) \subset \mathcal{O}(L^{-1})$. The metric h has a weight e^{φ} that is continuous, with zeroes contained in the union of $\{\sigma = 0\}$ and of the vertical divisor D^* . Moreover the curvature tensor $\Theta_{\mathcal{O}_{X_k}(1), h^{-1}} = \frac{i}{2\pi} \partial \bar{\partial} \log h$ satisfies by construction $\Theta_{\mathcal{O}_{X_k}(1), h^{-1}} \geq (pm + m')^{-1} \omega_k$. On the other hand, the continuity of the weight of h and the compactness of X_k imply that there exists a constant $C > 0$ such that $\|d\pi_{k,k-1}(\eta)\|_h \leq C \|\eta\|_{\omega_k}$ for all vectors $\eta \in V_k$ (notice that $\xi = d\pi_{k,k-1}(\eta) \in \mathcal{O}_{X_k}(-1)$). Now, the derivative $f'_{[k-1]}$ can be seen as a section of $f_{[k]}^* \mathcal{O}_{X_k}(-1)$, and we use this to define a singular hermitian metric $\gamma(t) i dt \wedge \bar{d}\bar{t}$ on \mathbb{C} by taking

$$\gamma(t) = \|f'_{[k-1]}(t)\|_{h(f_{[k]}(t))}^2.$$

If $f_{[k]}(\mathbb{C})$ is not contained in the divisor $\{\sigma = 0\}$, then γ is not identically zero and, in the sense of distributions, we find

$$\frac{i}{2\pi} \partial \bar{\partial} \log \gamma \geq f_{[k]}^* \Theta_{\mathcal{O}_{X_k}(1), h^{-1}} \geq (pm + m')^{-1} f_{[k]}^* \omega_k \geq C^{-1} (pm + m')^{-1} \gamma.$$

The final inequality comes from the inequality relating h and ω_k when we take $\eta = f'_{[k]}(t)$ and $\xi = f'_{[k-1]}(t)$. However, the Ahlfors lemma shows that a hermitian metric on \mathbb{C} with negative curvature bounded away from 0 cannot exist, thus we must have $f_{[k]}(\mathbb{C}) \subset \{\sigma = 0\}$. This proves our vanishing theorem in the case where P is invariant. The general case of a nonnecessarily invariant operator P will not be used here; a proof can be obtained by decomposing P into invariant parts and using an induction on m (cf. [Dem18] for details), or alternatively by means of Nevanlinna theory arguments ([SiYe97], [Siu97], see also [Dem97]). \square

Especially, we can apply the above vanishing theorem for any global invariant jet differential $P \in H^0(X, E_{k,m} V^* \otimes \mathcal{O}(-A))$. In that case, P corresponds bijectively to a section

$$(3.24) \quad \sigma \in H^0(X_k, \mathcal{O}_{X_k}(m) \otimes \pi_{k,0}^* \mathcal{O}(-A)),$$

and assuming $P \neq 0$, the vanishing theorem can be reinterpreted by stating that $f_{[k]}(\mathbb{C})$ is contained in the zero divisor $Z_\sigma \subset X_k$. Let $\Delta_k = \bigcup_{2 \leq \ell \leq k} \pi_{k,\ell}^{-1}(D_\ell)$ be the union of the vertical divisors (see (2.22) and (2.23)). Then $f_{[k]}(\mathbb{C})$ cannot be contained in Δ_k (as otherwise we would have $f'(t) = 0$ identically). We define the k -stage Green-Griffiths locus of (X, V) to be the Zariski closure

$$(3.25) \quad \text{GG}_k(X, V) = \overline{(X_k \setminus \Delta_k) \cap \bigcap_{m \in \mathbb{N}} \left(\text{base locus of } \mathcal{O}_{X_k}(m) \otimes \pi_{k,0}^* \mathcal{O}(-A) \right)}$$

(trivially independent of the choice of A), and

$$(3.26) \quad \text{GG}(X, V) = \bigcap_{k \in \mathbb{N}^*} \pi_{k,0}(\text{GG}_k(X, V)).$$

Then Theorem 3.23 implies that $f_{[k]}(\mathbb{C})$ must be contained in $\text{GG}_k(X, V)$ for every entire curve $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$, and also that $f(\mathbb{C}) \subset \text{GG}(X, V)$.

3.27. Corollary. *If $\text{GG}(X, V) = \emptyset$, then (X, V) is hyperbolic. In particular, if there exists $k \geq 1$ and a weight $a_\bullet \in \mathbb{N}^k$ such that $\mathcal{O}_{X_k}(a_\bullet)$ is ample on X_k , then (X, V) is hyperbolic.*

It should be observed that Corollary 3.27 yields a sufficient condition for hyperbolicity, but this is not a necessary condition. In fact, if we take $X = C_1 \times C_2$ to be a product of curves of genus ≥ 2 and $V = T_X$, it is easily checked that $\text{GG}(X) = \text{GG}(X, T_X) = X$. More general examples have been found by Diverio and Rousseau [DR15]. In a similar way, the Green-Griffiths-Lang conjecture holds for (X, V) if $Y := \text{GG}(X, V) \subsetneq X$, but this is only a sufficient condition. The following fundamental existence theorem, however, has been proved in [Dem11], using holomorphic Morse inequalities of [Dem85] as an essential tool. We only state the main result, as it will not be used here.

3.28. Theorem. *Let (X, V) be a projective directed manifold of general type, in the sense that the sheaf K_V of locally bounded sections of $\mathcal{O}(\det V^*)$ is big. Let A be an ample \mathbb{Q} -divisor on X such that $\mathcal{O}(\det V^*) \otimes \mathcal{O}(-A)$ is still ample. Then*

$$H^0 \left(X_k, \mathcal{O}_{X_k}(m) \otimes \pi_{k,0}^* \mathcal{O} \left(-\frac{m}{kr} \left(1 + \frac{1}{2} + \dots + \frac{1}{k} \right) A \right) \right) \neq 0$$

for $m \gg k \gg 1$ and m sufficiently divisible (so that the multiple of A is an integral divisor). In particular $\text{GG}_k(X, V) \subsetneq X_k$ for $k \gg 1$.

4. EXISTENCE OF HYPERBOLIC HYPERSURFACES OF LOW DEGREE

We give here a self-contained proof of the existence of hyperbolic surfaces of low degree in \mathbb{P}^{n+1} , using various techniques borrowed from the work of Toda [Toda71], Fujimoto [Fuj74], Green [Gre75], Nadel [Nad89], Siu-Yeung [SiYe96], Masuda-Noguchi [MaNo96] and Shiffman-Zaidenberg [ShZa02]. The main idea is to produce ad hoc differential equations for entire curves by means of Wronskian operators. This can be seen as a variation of Nadel's approach, that was actually based on Wronskians associated with meromorphic connections – Wronskian operators have the advantage of being much easier to handle than general jet differentials, thanks to their straightforward relationship with linear degeneracy.

4.A. GENERAL WRONSKIAN OPERATORS

This section follows closely the work of D. Brotbek [Brot17]. Let U be an open set of a complex manifold X , $\dim X = n$, and $s_0, \dots, s_k \in \mathcal{O}_X(U)$ be holomorphic functions. To these functions, we

can associate a Wronskian operator of order k defined by

$$(4.1) \quad W_k(s_0, \dots, s_k)(f) = \begin{vmatrix} s_0(f) & s_1(f) & \dots & s_k(f) \\ D(s_0(f)) & D(s_1(f)) & \dots & D(s_k(f)) \\ \vdots & \vdots & & \vdots \\ D^k(s_0(f)) & D^k(s_1(f)) & \dots & D^k(s_k(f)) \end{vmatrix},$$

where $f : t \mapsto f(t) \in U \subset X$ is a germ of holomorphic curve (or a k -jet of curve), and $D = \frac{d}{dt}$. For a biholomorphic change of variable φ of $(\mathbb{C}, 0)$, we find by induction on ℓ polynomial differential operators $Q_{\ell, i}$ of order $\leq \ell$ acting on φ satisfying

$$D^\ell(s_j(f \circ \varphi)) = \varphi'^\ell D^\ell(s_j(f)) \circ \varphi + \sum_{i < \ell} Q_{\ell, i}(\varphi', \dots, \varphi^{(\ell)}) D^i(s_j(f)) \circ \varphi.$$

It follows easily from this that

$$W_k(s_0, \dots, s_k)(f \circ \varphi) = (\varphi')^{1+2+\dots+k} W_k(s_0, \dots, s_k)(f) \circ \varphi,$$

hence $W_k(s_0, \dots, s_k)(f)$ is an invariant differential operator of degree $k' = \frac{1}{2}k(k+1)$. Especially, we get in this way a section that we denote

$$(4.2) \quad W_k(s_0, \dots, s_k) = \begin{vmatrix} s_0 & s_1 & \dots & s_k \\ D(s_0) & D(s_1) & \dots & D(s_k) \\ \vdots & \vdots & & \vdots \\ D^k(s_0) & D^k(s_1) & \dots & D^k(s_k) \end{vmatrix} \in H^0(U, E_{k, k'} T_X^*).$$

4.3. Proposition. *These Wronskian operators satisfy the following properties.*

(a) $W_k(s_0, \dots, s_k)$ is \mathbb{C} -multilinear and alternate in (s_0, \dots, s_k) .

(b) For any $g \in \mathcal{O}_X(U)$, we have

$$W_k(g s_0, \dots, g s_k) = g^{k+1} W_k(s_0, \dots, s_k).$$

Property 4.3 (b) is an easy consequence of the Leibniz formula

$$D^\ell(g(f) s_j(f)) = \sum_{k=0}^{\ell} \binom{\ell}{k} D^k(g(f)) D^{\ell-k}(s_j(f)),$$

by performing linear combinations of rows in the determinants. This property implies in its turn that one can define more generally an operator

$$(4.4) \quad W_k(s_0, \dots, s_k) \in H^0(U, E_{k, k'} T_X^* \otimes L^{k+1})$$

for any $(k+1)$ -tuple of sections $s_0, \dots, s_k \in H^0(U, L)$ of a holomorphic line bundle $L \rightarrow X$. In fact, when we compute the Wronskian in a local trivialization of $L|_U$, Property 4.3 (b) shows that the determinant is independent of the trivialization. Moreover, if $g \in H^0(U, G)$ for some line bundle $G \rightarrow X$, we have

$$(4.5) \quad W_k(g s_0, \dots, g s_k) = g^{k+1} W_k(s_0, \dots, s_k) \in H^0(U, E_{k, k'} T_X^* \otimes L^{k+1} \otimes G^{k+1}).$$

For global sections $\sigma_0, \dots, \sigma_k \in H^0(X, L)$, we thus get a Wronskian operator

$$(4.6) \quad W_k(s_0, \dots, s_k) \in H^0(X_k, \mathcal{O}_{X_k}(k') \otimes \pi_{k,0}^* L^{k+1})$$

on the k -stage X_k of the Semple tower. Very roughly, the idea for the construction of hyperbolic hypersurfaces is apply the fundamental vanishing theorem 3.23 to show that all entire curves have to satisfy certain Wronskian equations, leading in fine to exclude their existence. However, the vanishing theorem only holds for jet differentials in $H^0(X_k, \mathcal{O}_{X_k}(k') \otimes \pi_{k,0}^* A^{-1})$ with $A > 0$, while the existence of suitable sections $s_j \in H^0(X, L)$ can be achieved only when L is ample, so the

strategy seems a priori unapplicable. It turns out that one can sometimes arrange the Wronkian operator coefficients to be divisible by a section $\sigma_\Delta \in H^0(X, \mathcal{O}_X(\Delta))$ possessing a large zero divisor Δ , so that

$$(4.7) \quad \sigma_\Delta^{-1} W_k(s_0, \dots, s_k) \in H^0(X_k, \mathcal{O}_{X_k}(k') \otimes \pi_{k,0}^*(L^{k+1} \otimes \mathcal{O}_X(-\Delta))),$$

and we can then hope that $L^{k+1} \otimes \mathcal{O}_X(-\Delta) < 0$. The strategy is to find a variety X and sections $\sigma_0, \dots, \sigma_k \in H^0(X, L)$ for which the associated Wronskian $W_k(s_0, \dots, s_k)$ is highly divisible.

4.B. HYPERBOLICITY OF CERTAIN FERMAT-WARING HYPERSURFACES

Let Z be a non-singular $(n+1)$ -dimensional projective variety, and let A be a very ample divisor on Z ; the fundamental example is of course $Z = \mathbb{P}^{n+1}$ and $A = \mathcal{O}_{\mathbb{P}^{n+1}}(1)$. Our goal is to show that a well chosen $(n$ -dimensional) hypersurface $X = \{x \in Z; \sigma(x) = 0\}$ defined by a section $\sigma \in H^0(Z, A^d)$, $d \gg 1$, is Kobayashi hyperbolic. The construction explained below follows closely the ideas of Shiffman-Zaidenberg [ShZa02] and is based similarly on a use of Fermat-Waring type hypersurfaces. Our proof is however completely self-contained. The reader can consult Brody-Green [BrGr77], Nadel [Nad89] and Masuda-Noguchi [MaNo96] for constructions based on other techniques.

4.8. Theorem. *Let Z be a non-singular $(n+1)$ -dimensional projective variety, A a very ample divisor on Z , and $\tau_j \in H^0(Z, A)$, $0 \leq j \leq N$, sufficiently general sections. Then for $N \geq 2n$ and $d \geq N^2$, the hypersurface $X = \sigma^{-1}(0)$ associated with $\sigma = \sum_{0 \leq j \leq N} \tau_j^d \in H^0(Z, A^d)$ is Kobayashi hyperbolic.*

In particular, Theorem 4.8 provides examples of hyperbolic hypersurfaces of \mathbb{P}^{n+1} for all $n \geq 1$ and all degrees $d \geq 4n^2$. A substantially improved bound $d \geq [(n+3)^2/4]$ has been obtained recently by [DTH16] via a deformation argument for certain unions of hyperplanes, but the methods are quite different from the techniques used here. As in [ShZa02], the main step of our proof is the following proposition due to Toda [Toda71], Fujimoto [Fuj74] and Green [Gre75].

4.9. Proposition. *Let $g_j : \mathbb{C} \rightarrow \mathbb{C}$, $0 \leq j \leq N$, be non-zero entire functions such that the curve $g = [g_0 : \dots : g_N] : \mathbb{C} \rightarrow \mathbb{P}^N$ satisfies $\sum_{0 \leq j \leq N} g_j^d = 0$. If $d \geq N^2$, there exists a partition J_1, \dots, J_q of $\{0, 1, \dots, N\}$ such that $|J_s| \geq 2$, g_j/g_i is constant for all $i, j \in J_s$, and $\sum_{j \in J_s} g_j^d = 0$ for all $s = 1, 2, \dots, q$. If g is nonconstant, we must have $q \geq 2$.*

Proof. The result is true for $N = 1$ (with a single $J_1 = \{0, 1\}$), and for higher values $N \geq 2$ we apply induction and use vanishing arguments for Wronskians. The map $g = [g_0 : \dots : g_N] : \mathbb{C} \rightarrow \mathbb{P}^N$ can be seen as an entire curve drawn in the (smooth, irreducible) Fermat hypersurface $Y = \sum_{0 \leq j \leq N} z_j^d$ of \mathbb{P}^N . We set $k = N - 1$ and consider on Y the Wronskian operator

$$W_k(s_0, \dots, s_k) \quad \text{where } s_j(z) = z_j^d, \quad s_j \in H^0(Y, \mathcal{O}(d)).$$

Then

$$W_{N-1}(s_0, \dots, s_{N-1}) \in H^0(Y, E_{k,k'} T_Y^* \otimes \mathcal{O}(Nd)).$$

Since $D^\ell(s_j)$ is divisible by z_j^{d-k} for $\ell \leq k$, we conclude that $W_{N-1}(s_0, \dots, s_{N-1})$ is divisible by $\prod_{j < N} z_j^{d-k}$. However, as $s_0 = -(s_1 + \dots + s_N)$ on Y , we get

$$W_{N-1}(s_0, \dots, s_{N-1}) = (-1)^N W_{N-1}(s_1, \dots, s_N)$$

and conclude that $W_{N-1}(s_0, \dots, s_{N-1})$ must be also divisible by z_N^{d-k} . Since the $\{z_j = 0\}$, $0 \leq j \leq N$, form a normal crossing divisor on Y , we infer that

$$\widetilde{W} := \prod_{0 \leq j \leq N} z_j^{-(d-k)} W_{N-1}(s_0, \dots, s_{N-1}) \in H^0(Y, E_{k,k'} T_Y^* \otimes \mathcal{O}(Nd - (N+1)(d-k)))$$

i.e. $\widetilde{W} \in H^0(Y, E_{k,k'} T_Y^* \otimes \mathcal{O}(N^2 - 1 - d))$. By the fundamental vanishing theorem, we must have $\widetilde{W}(g) = 0$. Since this is equivalent to the vanishing of the determinant $\det(D^\ell(g_j^d))$, we conclude

that the functions g_0^d, \dots, g_{N-1}^d must be linearly dependent. After eliminating zero coefficients, we find a linear relation $\sum_{0 \leq k \leq p} c_k g_{j(k)}^d = 0$ with $c_k \in \mathbb{C}^*$, $j(k) \leq N-1$ and $1 \leq p \leq N-1$. The induction hypothesis applied to the functions $c_k^{1/d} g_{j(k)}$, implies that at least two of them are proportional. By grouping together the g_j 's that are proportional in the identity $\sum_{0 \leq j \leq N} g_j^d = 0$, we find a partition $(J_s)_{1 \leq s \leq q}$ of $\{0, 1, \dots, N\}$ and relations of the form $\sum_{j \in J_s} g_j^d = \lambda_s g_{j_s}^d$, $j_s \in J_s$. Moreover we get $\sum_{1 \leq s \leq q} \lambda_s g_{j_s}^d = 0$ with strictly less than $N+1$ functions g_{j_s} involved, all of them being pairwise nonproportional. This contradicts the induction hypothesis unless all coefficients λ_s are zero, and we must then have $|J_s| \geq 2$. The case $q = 1$ corresponds to g being constant. Proposition 4.9 follows. \square

Proof of Theorem 4.8. We argue by induction on $n \geq 1$. For $n = 1$, an easy adjunction argument shows that it is enough to take $d \geq 4$: sections of A can be used to embed the polarized surface (Z, A) in \mathbb{P}^N (e.g. with $N = 5$), and whenever $X = \sigma^{-1}(0)$ is a smooth curve, we have $K_X = K_{Z|X} \otimes A^d$ and a surjective restriction morphism $\Omega_{\mathbb{P}^N}^2 \rightarrow K_Z = \Lambda^2 T_Z^*$. As $\Omega_{\mathbb{P}^N}^2 \otimes \mathcal{O}(3) = \Lambda^{N-2}(T_{\mathbb{P}^N} \otimes \mathcal{O}(-1))$ is generated by sections, one sees that $K_Z \otimes A^3$ is also generated by sections, hence K_X is ample for $d \geq 4$.

Now, assume that the result is already proved for $n-1$ and consider a (non-constant) entire curve $f : \mathbb{C} \rightarrow X$ where $X = \{\sum_{0 \leq j \leq N} \tau_j^d = 0\} \subset Z$. For suitably chosen sections $\tau_j \in H^0(Z, A)$, $0 \leq j \leq N$ and $N \geq \dim Z = n+1$, the map $\tau := [\tau_0 : \dots : \tau_N] : Z \rightarrow \mathbb{P}^N$ can be taken to be a generically finite morphism. If $\tau_j \circ f$ vanishes for some j , say $j = N$, then f is drawn in the hypersurface X' of $Z' = \tau_N^{-1}(0)$ associated with $\sigma' = \sum_{0 \leq j \leq N-1} \tau_j^d$. We can suppose that Z' is smooth and, by the induction hypothesis for $(n-1, N-1)$, that X' is hyperbolic (notice that $N-1 \geq 2(n-1)$ and $d \geq (N-1)^2$); this is a contradiction.

Without loss of generality, we can thus assume that all sections $g_j := \tau_j \circ f$ are non-zero. Also suppose that $g = \tau \circ f$ is nonconstant. By definition of X , we have $\sum_{0 \leq j \leq N} g_j^d = 0$, and Proposition 4.9 shows that there exists a partition $J = \{J_1, \dots, J_q\}$ of $\{0, 1, \dots, N\}$ such that $q \geq 2$, $|J_s| \geq 2$, and the ratios $g_{j'}/g_j$ are constant for $j, j' \in J_s$, and $\sum_{j \in J_s} g_j^d = 0$ for all $s = 1, 2, \dots, q$. Set $j_s = \min J_s$ and $w_j = g_j/g_{j_s} \in \mathbb{C}^*$ for $j \in J_s \setminus \{j_s\}$. Then $g = [g_0 : \dots : g_N] = \tau \circ f$ is drawn in a projective linear subspace $Y_{J,w} \subset \mathbb{P}^{N-1}$ of dimension $q-1$ defined by the equations

$$(4.10) \quad Y_{J,w} : z_j = w_j z_{j_s} \quad \text{for } j \in J_s \setminus \{j_s\}, \quad 1 + \sum_{j \in J_s \setminus \{j_s\}} w_j^d = 0, \quad 1 \leq s \leq q.$$

Theorem 4.8 is now a consequence of the following lemma, which forces $g = \tau \circ f$, and hence f , to be constant. \square

4.11. Lemma. *For $N \geq 2n$ and $\tau_j \in H^0(Z, A)$ sufficiently general, $0 \leq j \leq N$, the hypersurface $X = \{\sum_{0 \leq j \leq N} \tau_j^d = 0\}$ is smooth and the map $\tau = [\tau_0 : \dots : \tau_N] : Z \rightarrow \mathbb{P}^N$ has a restriction $\tau : X \rightarrow \mathbb{P}^N$ that is a finite morphism. Moreover, for all partitions $J = \{J_s\}$ and all choices of $w = (w_j) \in (\mathbb{C}^*)^{N+1-q}$ as in (4.10), the preimage $\tau^{-1}(Y_{J,w})$ in Z is finite.*

Proof. Let $(\sigma_1, \dots, \sigma_m)$ be a basis of $H^0(Z, A)$. We write $\tau_j = \sum_{1 \leq \ell \leq m} a_{j\ell} \sigma_\ell$ and consider the matrix $a = (a_{j\ell}) \in \mathbb{C}^{m(N+1)}$. The singular locus of $X = \{\sum_{0 \leq j \leq N} \tau_j^d = 0\}$ is described by the equations

$$\sum_{0 \leq j \leq N} \left(\sum_{1 \leq \ell \leq m} a_{j\ell} \sigma_\ell(x) \right)^d = 0, \quad \frac{\partial}{\partial x_s} \left(\sum_{0 \leq j \leq N} \left(\sum_{1 \leq \ell \leq m} a_{j\ell} \sigma_\ell(x) \right)^d \right) = 0, \quad 1 \leq s \leq n+1$$

in coordinates. As the σ_ℓ 's generate all 1-jets at every point $x \in X$, we have $(n+2)$ independent equations in terms of a , hence the bad locus L of points $(x, a) \in Z \times \mathbb{C}^{m(N+1)}$ admits a fibration $\text{pr}_1 : L \rightarrow Z$ whose fibers are of dimension $m(N+1) - (n+2)$ in $\mathbb{C}^{m(N+1)}$. Therefore we get

$\dim L \leq m(N+1) - 1$ and $\text{pr}_2(L)$ does not cover $\mathbb{C}^{m(N+1)}$. Any matrix a taken in the complement $\mathbb{C}^{m(N+1)} \setminus \text{pr}_2(L)$ will produce a smooth hypersurface X .

Similarly, as the σ_ℓ 's separate points of Z , the set S of triples $(x_1, x_2, a) \in Z \times Z \times \mathbb{C}^{m(N+1)}$ with $x_1, x_2 \in X$, $x_1 \neq x_2$ and $\tau(x_1) = \tau(x_2)$ is such that the fibers of $S \rightarrow Z \times Z$ in $\mathbb{C}^{m(N+1)}$ are described by $N+1$ independent equations

$$\sum_{0 \leq j \leq N} \left(\sum_{1 \leq \ell \leq m} a_{j\ell} \sigma_\ell(x_1) \right)^d = 0, \quad \left[\sum_{1 \leq \ell \leq m} a_{j\ell} \sigma_\ell(x_1) \right]_{0 \leq j \leq N} = \left[\sum_{1 \leq \ell \leq m} a_{j\ell} \sigma_\ell(x_2) \right]_{0 \leq j \leq N} \in \mathbb{P}^N.$$

Therefore $\dim S = \dim(Z \times Z) + m(N+1) - (N+1) \leq m(N+1) + 1$ and the projection $S \rightarrow \mathbb{C}^{m(N+1)}$ has a fiber of dimension at most 1 over a generic point $a \in \mathbb{C}^{m(N+1)}$. For such a choice of a , if $F = \tau^{-1}(y)$ is a fiber of $\tau : X \rightarrow \mathbb{P}^N$, then S contains $F \times F \setminus \Delta_F$, hence we must have $\dim F = 0$, and all fibers F are finite.

In order to study the finiteness of $\tau^{-1}(Y_{J,w})$, we look at the incidence variety V_J of 4-tuples $(x_1, x_2, a, w) \in Z^2 \times \mathbb{C}^{m(N+1)} \times W_J$ such that $x_1 \neq x_2$ and $\tau(x_1) = \tau(x_2) \in Y_{J,w}$, where W_J is the set of points $w = (w_j)$ such that $1 + \sum_{j \in J_s \setminus \{j_s\}} w_j^d = 0$, $1 \leq s \leq q$. This variety will detect the fibers $\tau^{-1}(Y_{J,w})$ that contain at least two distinct points. Notice also that we have only finitely many subvarieties W_J involved, and that $\dim W_J = \sum (|J_s| - 2) = N + 1 - 2q$. The variety V_J is defined by $2(N+1-q) + q - 1$ linear equations in the $a_{j\ell}$:

$$\sum_{1 \leq \ell \leq m} (a_{j\ell} - w_j a_{j_s \ell}) \sigma_\ell(x_i) = 0, \quad j \in J_s \setminus \{j_s\}, \quad 1 \leq s \leq q, \quad i = 1, 2,$$

$$\left[\sum_{1 \leq \ell \leq m} a_{j_s \ell} \sigma_\ell(x_1) \right]_{1 \leq s \leq q} = \left[\sum_{1 \leq \ell \leq m} a_{j_s \ell} \sigma_\ell(x_2) \right]_{1 \leq s \leq q} \in \mathbb{P}^{q-1}.$$

These equations are independent: this is again a consequence of the fact that the σ_ℓ 's separate points of Z . The dimension of V_J is thus

$$\begin{aligned} \dim V_J &= m(N+1) + 2(n+1) + (N+1-2q) - (2(N+1-q) + (q-1)) \\ &= m(N+1) + 2n + 2 - N - q. \end{aligned}$$

For $q \geq 2$ and $N \geq 2n$, we have $\dim V_J \leq n(N+1)$, therefore the projection $V_J \rightarrow \mathbb{C}^{m(N+1)}$ has finite fibers over a Zariski open set $\mathbb{C}^{m(N+1)} \setminus S_J$. Hence, for $a \in \mathbb{C}^{m(N+1)} \setminus \bigcup S_J$, we infer that all sets $\tau^{-1}(Y_{J,w})$ are finite. (For $N \geq 2n+1$, we could even take a outside of the projections of the incidence varieties V_J , and in that case, for a generic, the sets $\tau^{-1}(Y_{J,w})$ have at most one point). \square

5. PROOF OF THE KOBAYASHI CONJECTURE ON THE HYPERBOLICITY OF GENERAL HYPERSURFACES

In this section, our more ambitious goal is to give a simple proof of the Kobayashi conjecture, combining ideas of Green-Griffiths [GrGr80], Demailly [Dem95], Brotbek [Brot17] and Ya Deng [Deng16], in chronological order. Related ideas had been used earlier in [Xie18] and then in [BrDa18], to establish Debarre's conjecture on the ampleness of the cotangent bundle of generic complete intersections, when their codimension is at least equal to the dimension.

5.A. USING BLOW-UPS OF WRONSKIAN IDEAL SHEAVES

Let X be a projective non-singular algebraic variety and $L \rightarrow X$ a line bundle over X . We consider a linear system $\Sigma \subset H^0(X, L)$ producing some non-zero Wronskian sections $W_k(s_0, \dots, s_k)$, so that $\dim \Sigma \geq k+1$. As the Wronskian is alternate and multilinear in the arguments s_j , we get a meromorphic map $X_k \dashrightarrow P(\Lambda^{k+1} \Sigma^*)$ by sending a k -jet $\gamma = f_{[k]}(0) \in X_k$ to the point of projective coordinates $[W_k(u_{i_0}, \dots, u_{i_k})(f)(0)]_{i_0, \dots, i_k}$, where $(u_j)_{j \in J}$ is a basis of Σ and $i_0, \dots, i_k \in J$ are in

increasing order. This assignment factorizes through the Plücker embedding into a meromorphic map

$$\Phi : X_k \dashrightarrow \text{Gr}_{k+1}(\Sigma)$$

into the Grassmannian of dimension $k + 1$ subspaces of Σ^* (or codimension $k + 1$ subspaces of Σ , alternatively). In fact, if $L|_U \simeq U \times \mathbb{C}$ is a trivialization of L in a neighborhood of a point $x_0 = f(0) \in X$, we can consider the map $\Psi_U : X_k \rightarrow \text{Hom}(\Sigma, \mathbb{C}^{k+1})$ given by

$$\pi_{k,0}^{-1}(U) \ni f_{[k]} \mapsto (s \mapsto (D^\ell(s(f)))_{0 \leq \ell \leq k}),$$

and associate either the kernel $\Xi \subset \Sigma$ of $\Psi_U(f_{[k]})$, seen as a point $\Xi \in \text{Gr}_{k+1}(\Sigma)$, or $\Lambda^{k+1}\Xi^\perp \subset \Lambda^{k+1}\Sigma^*$, seen as a point of $P(\Lambda^{k+1}\Sigma^*)$ (assuming that we are at a point where the rank is equal to $k + 1$). Let $\mathcal{O}_{\text{Gr}}(1)$ be the tautological very ample line bundle on $\text{Gr}_{k+1}(\Sigma)$ (equal to the restriction of $\mathcal{O}_{P(\Lambda^{k+1}\Sigma^*)}(1)$). By construction, Φ is induced by the linear system of sections

$$W_k(u_{i_0}, \dots, u_{i_k}) \in H^0(X_k, \mathcal{O}_{X_k}(k') \otimes \pi_{k,0}^* L^{k+1}),$$

and we thus get a natural isomorphism

$$(5.1) \quad \mathcal{O}_{X_k}(k') \otimes \pi_{k,0}^* L^{k+1} \simeq \Phi^* \mathcal{O}_{\text{Gr}}(1) \quad \text{on } X_k \setminus B_k,$$

where $B_k \subset X_k$ is the base locus of our linear system of Wronskians. The presence of the indeterminacy set B_k may create trouble in analyzing the positivity of our line bundles, so we are going to use an appropriate blow-up to resolve the indeterminacies. For this purpose, we introduce the ideal sheaf $\mathcal{J}_{k,\Sigma} \subset \mathcal{O}_{X_k}$ generated by the linear system Σ , and take a modification $\mu_{k,\Sigma} : \widehat{X}_{k,\Sigma} \rightarrow X_k$ in such a way that $\mu_{k,\Sigma}^* \mathcal{J}_{k,\Sigma} = \mathcal{O}_{\widehat{X}_{k,\Sigma}}(-F_{k,\Sigma})$ for some divisor $F_{k,\Sigma}$ in $\widehat{X}_{k,\Sigma}$. Then Φ is resolved into a morphism $\Phi \circ \mu_{k,\Sigma} : \widehat{X}_{k,\Sigma} \rightarrow \text{Gr}_{k+1}(\Sigma)$, and on $\widehat{X}_{k,\Sigma}$, (5.1) becomes an everywhere defined isomorphism

$$(5.2) \quad \mu_{k,\Sigma}^* (\mathcal{O}_{X_k}(k') \otimes \pi_{k,0}^* L^{k+1}) \otimes \mathcal{O}_{\widehat{X}_{k,\Sigma}}(-F_{k,\Sigma}) \simeq (\Phi \circ \mu_{k,\Sigma})^* \mathcal{O}_{\text{Gr}}(1).$$

In fact, we can simply take \widehat{X}_k to be the normalized blow-up of $\mathcal{J}_{k,\Sigma}$, i.e. the normalization of the closure $\Gamma \subset X_k \times \text{Gr}_{k+1}(\Sigma)$ of the graph of Φ and $\mu_{k,\Sigma} : \widehat{X}_k \rightarrow X_k$ to be the composition of the normalization map $\widehat{X}_k \rightarrow \Gamma$ with the first projection $\Gamma \rightarrow X_k$. [The Hironaka desingularization theorem would possibly allow us to replace \widehat{X}_k by a nonsingular modification, and $F_{k,\Sigma}$ by a simple normal crossing divisor on the desingularization; we will avoid doing so here, as we would otherwise need to show the existence of universal desingularizations when (X_t, Σ_t) is a family of linear systems of k -jets of sections associated with a family of algebraic varieties]. The following basic lemma was observed by Ya Deng [Deng16].

5.3. Lemma. *Locally over coordinate open sets $U \subset X$ on which $L|_U$ is trivial, there is a maximal “Wronskian ideal sheaf” $\mathcal{J}_k^X \supset \mathcal{J}_{k,\Sigma}$ in \mathcal{O}_{X_k} achieved by linear systems $\Sigma \subset H^0(U, L)$. It is attained globally on X whenever the linear system $\Sigma \subset H^0(X, L)$ generates k -jets of sections of L at every point. Finally, it is “universal” in the sense that it does not depend on L and behaves functorially under immersions: if $\psi : X \rightarrow Y$ is an immersion and $\mathcal{J}_k^X, \mathcal{J}_k^Y$ are the corresponding Wronskian ideal sheaves in $\mathcal{O}_{X_k}, \mathcal{O}_{Y_k}$, then $\psi_k^* \mathcal{J}_k^Y = \mathcal{J}_k^X$ with respect to the induced immersion $\psi_k : X_k \rightarrow Y_k$.*

Proof. The (local) existence of such a maximal ideal sheaf is merely a consequence of the strong Noetherian property of coherent ideals. As observed at the end of section 2.D, the bundle $X_k \rightarrow X$ is a locally trivial tower of \mathbb{P}^{n-1} -bundles, with a fiber $\mathcal{R}_{n,k}$ that is a rational $k(n-1)$ -dimensional variety; over any coordinate open set $U \subset X$ equipped with local coordinates $(z_1, \dots, z_n) \in B(0, r) \subset \mathbb{C}^n$, it is isomorphic to the product $U \times \mathcal{R}_{n,k}$, the fiber over a point $x_0 \in U$ being identified with the central fiber through a translation $(t \mapsto f(t)) \mapsto (t \mapsto x_0 + f(t))$ of germs of curves. In this setting, \mathcal{J}_k^X is generated by the functions in \mathcal{O}_{X_k} associated with Wronskians

$$X_k|_U \ni \xi = f_{[k]} \mapsto W_k(s_0, \dots, s_k)(f) \in \mathcal{O}_{X_k}(k')|_{\mathcal{R}_{n,k}}, \quad s_j \in H^0(U, \mathcal{O}_X),$$

by taking local trivializations $\mathcal{O}_{X_k}(k')_{\xi_0} \simeq \mathcal{O}_{X_k, \xi_0}$ at points $\xi_0 \in X_k$. In fact, it is enough to take Wronskians associated with *polynomials* $s_j \in \mathbb{C}[z_1, \dots, z_n]$. To see this, one can e.g. invoke Krull's lemma for local rings, which implies $\mathcal{J}_{k, \xi_0}^X = \bigcap_{\ell \geq 0} (\mathcal{J}_{k, \xi_0}^X + \mathfrak{m}_{\xi_0}^{\ell+1})$, and to observe that ℓ -jets of Wronskians $W_k(s_0, \dots, s_k) \pmod{\mathfrak{m}_{\xi_0}^{\ell+1}}$ depend only on the $(k + \ell)$ -jets of the sections s_j in $\mathcal{O}_{X, x_0}/\mathfrak{m}_{x_0}^{k+\ell+1}$, where $x_0 = \pi_{k,0}(\xi_0)$. Therefore, polynomial sections s_j or arbitrary holomorphic functions s_j define the same ℓ -jets of Wronskians for any ℓ . Now, in the case of polynomials, it is clear that translations $(t \mapsto f(t)) \mapsto (t \mapsto x_0 + f(t))$ leave \mathcal{J}_k^X invariant, hence \mathcal{J}_k^X is the pull-back by the second projection $X_k|_U \simeq U \times \mathcal{R}_{n,k} \rightarrow \mathcal{R}_{n,k}$ of its restriction to any of the fibers $\pi_{k,0}^{-1}(x_0) \simeq \mathcal{R}_{n,k}$. As the k -jets of the s_j 's at x_0 are sufficient to determine the restriction of our Wronskians to $\pi_{k,0}^{-1}(x_0)$, the first two claims of Lemma 5.3 follow. The universality property comes from the fact that $L|_U$ is trivial (cf. Property 4.3 b) and that germs of sections of \mathcal{O}_X extend to germs of sections of \mathcal{O}_Y via the immersion ψ . (Notice that in this discussion, one may have to pick Taylor expansions of order $> k$ for f to reach all points of the fiber $\pi_{k,0}^{-1}(x_0)$, the order $2k - 1$ being sufficient by [Dem95, Proposition 5.11], but this fact does not play any role here). A consequence of universality is that \mathcal{J}_k^X does not depend on coordinates nor on the geometry of X . \square

The above discussion combined with Lemma 5.3 leads to the following statement.

5.4. Proposition. *Assume that L generates all k -jets of sections (e.g. take $L = A^p$ with A very ample and $p \geq k$), and let $\Sigma \subset H^0(X, L)$ be a linear system that also generates k -jets of sections at any point of X . Then we have a universal isomorphism*

$$\mu_k^*(\mathcal{O}_{X_k}(k') \otimes \pi_{k,0}^* L^{k+1}) \otimes \mathcal{O}_{\widehat{X}_k, \Sigma}(-F_k) \simeq (\Phi \circ \mu_k)^* \mathcal{O}_{\text{Gr}_{k+1}(\Sigma)}(1),$$

where $\mu_k : \widehat{X}_k \rightarrow X_k$ is the normalized blow-up of the (maximal) ideal sheaf $\mathcal{J}_k^X \subset \mathcal{O}_{X_k}$ associated with order k Wronskians, and F_k the universal divisor of \widehat{X}_k resolving \mathcal{J}_k^X .

5.B. SPECIALIZATION TO SUITABLE HYPERSURFACES

As in §4.B, let Z be a non-singular $(n + 1)$ -dimensional projective variety polarized with a very ample divisor A . We are going to show that a sufficiently general algebraic hypersurface $X = \{x \in Z; \sigma(x) = 0\}$ defined by $\sigma \in H^0(Z, A^d)$ is Kobayashi hyperbolic when d is large. Brotbek's main idea developed in [Brot17] is that a carefully selected hypersurface (of a more complicated type than the Fermat-Waring hypersurfaces considered in §4) may have enough Wronskian sections to directly imply the ampleness of some tautological jet line bundle – a Zariski open property. Here, we take σ be a sum of terms

$$(5.5) \quad \sigma = \sum_{0 \leq j \leq N} a_j m_j^\delta, \quad a_j \in H^0(Z, A^p), \quad m_j \in H^0(Z, A^b), \quad n < N \leq k, \quad d = \delta b + \rho,$$

where $\delta \gg 1$ and the m_j are “monomials” of the same degree b , i.e. product of b “linear” sections $\tau_I \in H^0(Z, A)$, and the factors a_j are general enough. The integer ρ is taken in the range $[k, k + b - 1]$, first to ensure that $H^0(Z, A^p)$ generates k -jets of sections, and second, to allow d to be an arbitrary large integer (once $\delta \geq \delta_0$ has been chosen large enough).

The monomials m_j will be chosen in such a way that for suitable $c \in \mathbb{N}$, $1 \leq c \leq N$, any subfamily of c terms m_j shares a common factor $\tau_I \in H^0(X, A)$. To this end, we consider all subsets $I \subset \{0, 1, \dots, N\}$ with $\text{card } I = c$; there are $B = \binom{N+1}{c}$ subsets of this type. For all such I , we select sections $\tau_I \in H^0(Z, A)$ such that $\prod_I \tau_I = 0$ is a simple normal crossing divisor in Z (with all of its components of multiplicity 1). For $j = 0, 1, \dots, N$ given, the number of subsets I containing j is $b = \binom{N}{c-1}$. We put

$$(5.6) \quad m_j = \prod_{I \ni j} \tau_I \in H^0(Z, A^b).$$

The first step consists in checking that we can achieve X to be smooth with these constraints.

5.7. Lemma. *Assume $N \geq c(n+1)$. Then, for a generic choice of the sections $a_j \in H^0(Z, A^\rho)$ and $\tau_I \in H^0(Z, A)$, the hypersurface $X = \sigma^{-1}(0) \subset Z$ defined by (5.5), (5.6) is non-singular. Moreover, under the same condition for N , the intersection of $\prod \tau_I = 0$ with X can be taken to be a simple normal crossing divisor in X .*

Proof. As the properties considered in the Lemma are Zariski open properties in terms of the $(N + B + 1)$ -tuple (a_j, τ_I) , it is sufficient to prove the result for a specific choice of the a_j 's: we fix here $a_j = \tilde{\tau}_j \tau_{I(j)}^{\rho-1}$ where $\tilde{\tau}_j \in H^0(X, A)$, $0 \leq j \leq N$ are new sections such that $\prod \tilde{\tau}_j \prod \tau_I = 0$ is a simple normal crossing divisor, and $I(j)$ is any subset of cardinal c containing j . Let H be the hypersurface of degree d of \mathbb{P}^{N+B} defined in homogeneous coordinates $(z_j, z_I) \in \mathbb{C}^{N+B+1}$ by $h(z) = 0$ where

$$h(z) = \sum_{0 \leq j \leq N} z_j z_{I(j)}^{\rho-1} \prod_{I \ni j} z_I^\delta,$$

and consider the morphism $\Phi : Z \rightarrow \mathbb{P}^{N+B}$ such that $\Phi(x) = (\tilde{\tau}_j(x), \tau_I(x))$. With our choice of the a_j 's, we have $\sigma = h \circ \Phi$. Now, when the $\tilde{\tau}_j$ and τ_I are general enough, the map Φ defines an embedding of Z into \mathbb{P}^{N+B} (for this, one needs $N + B \geq 2 \dim Z + 1 = 2n + 3$, which is the case by our assumptions). Then, by definition, X is isomorphic to the intersection of H with $\Phi(Z)$. Changing generically the $\tilde{\tau}_j$ and τ_I 's can be achieved by composing Φ with a generic automorphism $g \in \text{Aut}(\mathbb{P}^{N+B}) = \text{PGL}_{N+B+1}(\mathbb{C})$ (as $\text{GL}_{N+B+1}(\mathbb{C})$ acts transitively on $(N + B + 1)$ -tuples of linearly independent linear forms). As $\dim g \circ \Phi(Z) = \dim Z = n + 1$, Lemma 5.7 will follow from a standard Bertini argument if we can check that $\text{Sing}(H)$ has codimension at least $n + 2$ in \mathbb{P}^{N+B} . In fact, this condition implies $\text{Sing}(H) \cap (g \circ \Phi(Z)) = \emptyset$ for g generic, while $g \circ \Phi(Z)$ can be chosen transverse to $\text{Reg}(H)$. Now, a sufficient condition for smoothness is that one of the differentials dz_j , $0 \leq j \leq N$, appears with a non-zero factor in $dh(z)$ (just neglect the other differentials $*dz_I$ in this argument). We infer from this and the fact that $\delta \geq 2$ that $\text{Sing}(H)$ consists of the locus defined by $\prod_{I \ni j} z_I = 0$ for all $j = 0, 1, \dots, N$. It is the union of the linear subspaces $z_{I_0} = \dots = z_{I_N} = 0$ for all possible choices of subsets I_j such that $I_j \ni j$. Since $\text{card } I_j = c$, the equality $\bigcup I_j = \{0, 1, \dots, N\}$ implies that there are at least $\lceil (N + 1)/c \rceil$ distinct subsets I_j involved in each of these linear subspaces, and the equality can be reached. Therefore $\text{codim } \text{Sing}(H) = \lceil (N + 1)/c \rceil \geq n + 2$ as soon as $N \geq c(n + 1)$. By the same argument, we can assume that the intersection of $g \circ \Phi(Z)$ with at least $(n + 2)$ distinct hyperplanes $z_I = 0$ is empty. In order that $\prod \tau_I = 0$ defines a normal crossing divisor at a point $x \in X$, it is sufficient to ensure that for any family \mathcal{G} of coordinate hyperplanes $z_I = 0$, $I \in \mathcal{G}$, with $\text{card } \mathcal{G} \leq n + 1$, we have a ‘‘free’’ index $j \notin \bigcup_{I \in \mathcal{G}} I$ such that $x_I \neq 0$ for all $I \ni j$, so that dh involves a non-zero term $*dz_j$ independent of the dz_I , $I \in \mathcal{G}$. If this fails, there must be at least $(n + 2)$ hyperplanes $z_I = 0$ containing x , associated either with $I \in \mathcal{G}$, or with other I 's covering $\mathbb{C}(\bigcup_{I \in \mathcal{G}} I)$. The corresponding bad locus is of codimension at least $(n + 2)$ in \mathbb{P}^{N+B} and can be avoided by $g \circ \Phi(Z)$ for a generic choice of $g \in \text{Aut}(\mathbb{P}^{N+B})$. Then $X \cap \bigcap_{I \in \mathcal{G}} \tau_I^{-1}(0)$ is smooth of codimension equal to $\text{card } \mathcal{G}$. \square

5.C. CONSTRUCTION OF HIGHLY DIVISIBLE WRONSKIANs

To any families $s, \hat{\tau}$ of sections $s_1, \dots, s_r \in H^0(Z, A^k)$, $\hat{\tau}_1, \dots, \hat{\tau}_r \in H^0(Z, A)$, and any subset $J \subset \{0, 1, \dots, N\}$ with $\text{card } J = c$, we associate a Wronskian operator of order k (i.e. a $(k + 1) \times (k + 1)$ -determinant)

$$(5.8) \quad W_{k,s,\hat{\tau},a,J} = W_k(s_1 \hat{\tau}_1^{d-k}, \dots, s_r \hat{\tau}_r^{d-k}, (a_j m_j^\delta)_{j \in \mathbb{C}J}), \quad r = k + c - N, \quad |\mathbb{C}J| = N - c + 1.$$

We assume here again that the $\hat{\tau}_j$ are chosen so that $\prod \hat{\tau}_j \prod \tau_I = 0$ defines a simple normal crossing divisor in Z and X . Since $s_j \hat{\tau}_j^{d-k}, a_j m_j^\delta \in H^0(Z, A^d)$, formula (4.6) applied with $L = A^d$ implies that

$$(5.9) \quad W_{k,s,\hat{\tau},a,J} \in H^0(Z, E_{k,k'} T_Z^* \otimes A^{(k+1)d}).$$

However, we are going to see that $W_{k,s,\hat{\tau},a,J}$ and its restriction $W_{k,s,\hat{\tau},a,J|X}$ are divisible by monomials $\hat{\tau}^\alpha \tau^\beta$ of very large degree, where $\hat{\tau}$, resp. τ , denotes the collection of sections $\hat{\tau}_j$, resp. τ_I in $H^0(Z, A)$. In this way, we will see that we can even obtain a negative exponent of A after simplifying $\hat{\tau}^\alpha \tau^\beta$ in $W_{k,s,\hat{\tau},a,J|X}$. This simplification process is a generalization of techniques already considered by [Siu87] and [Nad89] (and later [DeEG97]), in relation with the use of meromorphic connections of low pole order.

5.10. Lemma. *Assume that $\delta \geq k$. Then the Wronskian operator $W_{k,s,\hat{\tau},a,J}$, resp. $W_{k,s,\hat{\tau},a,J|X}$, is divisible by a monomial $\hat{\tau}^\alpha \tau^\beta$, resp. $\hat{\tau}^\alpha \tau^\beta \tau_J^{\delta-k}$ (with a multi-index notation $\hat{\tau}^\alpha \tau^\beta = \prod \hat{\tau}_j^{\alpha_j} \prod \tau_I^{\beta_I}$), and*

$$\alpha, \beta \geq 0, \quad |\alpha| = r(d - 2k), \quad |\beta| = (N + 1 - c)(\delta - k)b.$$

Proof. $W_{k,s,\hat{\tau},a,J}$ is obtained as a determinant whose r first columns are the derivatives $D^\ell(s_j \hat{\tau}_j^{d-k})$ and the last $N + 1 - c$ columns are the $D^\ell(a_j m_j^\delta)$, divisible respectively by $\hat{\tau}_j^{d-2k}$ and $m_j^{\delta-k}$. As m_j is of the form τ^γ , $|\gamma| = b$, this implies the divisibility of $W_{k,s,\hat{\tau},a,J}$ by a monomial of the form $\hat{\tau}^\alpha \tau^\beta$, as asserted. Now, we explain why one can gain the additional factor $\tau_J^{\delta-k}$ dividing the restriction $W_{k,s,\hat{\tau},a,J|X}$. First notice that τ_J does not appear as a factor in $\hat{\tau}^\alpha \tau^\beta$, precisely because the Wronskian involves only terms $a_j m_j^\delta$ with $j \notin J$, hence these m_j 's do not contain τ_J . Let us pick $j_0 = \min(\mathbb{C}J) \in \{0, 1, \dots, N\}$. Since X is defined by $\sum_{0 \leq j \leq N} a_j m_j^\delta = 0$, we have identically

$$a_{j_0} m_{j_0}^\delta = - \sum_{i \in J} a_i m_i^\delta - \sum_{i \in \mathbb{C}J \setminus \{j_0\}} a_i m_i^\delta$$

in restriction to X , whence (by the alternate property of $W_k(\bullet)$)

$$W_{k,s,\hat{\tau},a,J|X} = - \sum_{i \in J} W_k(s_1 \hat{\tau}_1^{d-k}, \dots, s_r \hat{\tau}_r^{d-k}, a_i m_i^\delta, (a_j m_j^\delta)_{j \in \mathbb{C}J \setminus \{j_0\}})_{|X}.$$

However, all terms m_i , $i \in J$, contain by definition the factor τ_J , and the derivatives $D^\ell(\bullet)$ leave us a factor $m_i^{\delta-k}$ at least. Therefore, the above restricted Wronskian is also divisible by $\tau_J^{\delta-k}$, thanks to the fact that $\prod \hat{\tau}_j \prod \tau_I = 0$ forms a simple normal crossing divisor in X . \square

5.11. Corollary. *For $\delta \geq k$, there exists a monomial $\hat{\tau}^{\alpha_J} \tau^{\beta_J}$ dividing $W_{k,s,\hat{\tau},a,J|X}$ such that*

$$|\alpha_J| + |\beta_J| = (k + c - N)(d - 2k) + (N + 1 - c)(\delta - k)b + (\delta - k)$$

and we have

$$\widetilde{W}_{k,s,\hat{\tau},a,J|X} := (\hat{\tau}^{\alpha_J} \tau^{\beta_J})^{-1} W_{k,s,\hat{\tau},a,J|X} \in H^0(X, E_{k,k'} T_X^* \otimes A^{-p}),$$

where

$$(5.12) \quad p = |\alpha_J| + |\beta_J| - (k + 1)d = (\delta - k) - (k + c - N)2k - (N + 1 - c)(kb + \rho).$$

In particular, we have $p > 0$ for δ large enough (all other parameters being fixed or bounded), and under this assumption, the fundamental vanishing theorem implies that all entire curves $f : \mathbb{C} \rightarrow X$ are annihilated by these Wronskian operators.

Proof. In fact,

$$(k + 1)d = (k + c - N)d + (N + 1 - c)d = (k + c - N)d + (N + 1 - c)(\delta b + \rho)$$

and we get (5.12) by subtraction. \square

5.D. CONTROL OF THE BASE LOCUS FOR SUFFICIENTLY GENERAL COEFFICIENTS a_j IN σ

The next step is to control more precisely the base locus of these Wronskians and to find conditions on $N, k, c, d = b\delta + \rho$ ensuring that the base locus is empty for a generic choice of the sections a_j in $\sigma = \sum a_j m_j$. Although we will not formally use it, the next lemma is useful to realize that the base locus is related to a natural rank condition.

5.13. Lemma. *Set $u_j := a_j m_j^\delta$. The base locus in X_k^{reg} of the above Wronskians $W_{k,s,\hat{\tau},a,J|X}$, when $s, \hat{\tau}$ vary, consists of jets $f_{[k]}(0) \in X_k^{\text{reg}}$ such that the matrix $(D^\ell(u_j \circ f)(0))_{0 \leq \ell \leq k, j \in \mathbb{C}J}$ is not of maximal rank (i.e., of rank $< \text{card } \mathbb{C}J = N + 1 - c$); if $\delta > k$, this includes all jets $f_{[k]}(0)$ such that $f(0) \in \bigcup_{I \neq J} \tau_I^{-1}(0)$. When J also varies, the base locus of all $W_{k,s,\hat{\tau},a,J|X}$ in the Zariski open set $X'_k := X_k^{\text{reg}} \setminus \bigcup_{|I|=c} \tau_I^{-1}(0)$ consists of all k -jets such that $\text{rank}(D^\ell(u_j \circ f)(0))_{0 \leq \ell \leq k, 0 \leq j \leq N} \leq N - c$.*

Proof. If $\delta > k$ and $m_j \circ f(0) = 0$ for some $j \in J$, we have in fact $D^\ell(u_j \circ f)(0) = 0$ for all derivatives $\ell \leq k$, because the exponents involved in all factors of the differentiated monomial $a_j m_j^\delta$ are at least equal to $\delta - k > 0$. Hence the rank of the matrix cannot be maximal. Now, assume that $m_j \circ f(0) \neq 0$ for all $j \in \mathbb{C}J$, i.e.

$$(5.14) \quad x_0 := f(0) \in X \setminus \bigcup_{j \in \mathbb{C}J} m_j^{-1}(0) = X \setminus \bigcup_{I \neq J} \tau_I^{-1}(0).$$

We take sections $\hat{\tau}_j$ so that $\hat{\tau}_j(x_0) \neq 0$, and then adjust the k -jet of the sections s_1, \dots, s_r in order to generate any matrix of derivatives $(D^\ell(s_j(f)\hat{\tau}_j(f)^{d-k})(0))_{0 \leq \ell \leq k, j \in \mathbb{C}J}$ (the fact that $f'(0) \neq 0$ is used for this!). Therefore, by expanding the determinant according to the last $N + 1 - c$ columns, we see that the base locus is defined by the equations

$$(5.15) \quad \det(D^\ell(u_j(f))(0))_{\ell \in L, j \in \mathbb{C}J} = 0, \quad \forall L \subset \{0, 1, \dots, k\}, \quad |L| = N + 1 - c,$$

equivalent to the non-maximality of the rank. The last assertion follows by a simple linear algebra argument. \square

For a finer control of the base locus, we adjust the family of coefficients

$$(5.16) \quad a = (a_j)_{0 \leq j \leq N} \in S := H^0(Z, A^\rho)^{\oplus(N+1)}$$

in our section $\sigma = \sum a_j m_j^\delta \in H^0(Z, A^d)$, and denote by $X_a = \sigma^{-1}(0) \subset Z$ the corresponding hypersurface. By Lemma 5.7, we know that there is a Zariski open set $U \subset S$ such that X_a is smooth and $\prod \tau_I = 0$ is a simple normal crossing divisor in X_a for all $a \in U$. We consider the Semple tower $X_{a,k} := (X_a)_k$ of X_a , the ‘‘universal blow-up’’ $\mu_{a,k} : \widehat{X}_{a,k} \rightarrow X_{a,k}$ of the Wronskian ideal sheaf $\mathcal{J}_{a,k}$ such that $\mu_{a,k}^* \mathcal{J}_{a,k} = \mathcal{O}_{\widehat{X}_{a,k}}(-F_{a,k})$ for some ‘‘Wronskian divisor’’ $F_{a,k}$ in $\widehat{X}_{a,k}$. By the universality of this construction, we can also embed $X_{a,k}$ in the Semple tower Z_k of Z , blow up the Wronskian ideal sheaf \mathcal{J}_k^Z of Z_k to get a Wronskian divisor F_k in \widehat{Z}_k where $\mu_k : \widehat{Z}_k \rightarrow Z_k$ is the blow-up map. Then $F_{a,k}$ is the restriction of F_k to $\widehat{X}_{a,k} \subset \widehat{Z}_k$. Our section $\widetilde{W}_{k,a,\hat{\tau},s,J|X_a}$ is the restriction of a *meromorphic* section defined on Z , namely

$$(5.17) \quad (\hat{\tau}^{\alpha_J} \tau^{\beta_J})^{-1} W_{k,s,\hat{\tau},a,J} = (\hat{\tau}^{\alpha_J} \tau^{\beta_J})^{-1} W_k(s_1 \hat{\tau}_1^{d-k}, \dots, s_r \hat{\tau}_r^{d-k}, (a_j m_j^\delta)_{j \in \mathbb{C}J}).$$

It induces over the Zariski open set $Z' = Z \setminus \bigcup_I \tau_I^{-1}(0)$ a holomorphic section

$$(5.18) \quad \sigma_{k,s,\hat{\tau},a,J} \in H^0(\widehat{Z}'_k, \mu_k^*(\mathcal{O}_{Z_k}(k') \otimes \pi_{k,0}^* A^{-p}) \otimes \mathcal{O}_{\widehat{Z}'_k}(-F_k))$$

(notice that the relevant factors $\hat{\tau}_j$ remain divisible on the whole variety Z). By construction, thanks to the divisibility property explained in Lemma 5.10, the restriction of this section to $\widehat{X}'_{a,k} = \widehat{X}_{a,k} \cap \widehat{Z}'_k$ extends holomorphically to $\widehat{X}_{a,k}$, i.e.

$$(5.19) \quad \sigma_{k,s,\hat{\tau},a,J|_{\widehat{X}_{a,k}}} \in H^0(\widehat{X}_{a,k}, \mu_{a,k}^*(\mathcal{O}_{X_{a,k}}(k') \otimes \pi_{k,0}^* A^{-p}) \otimes \mathcal{O}_{\widehat{X}_{a,k}}(-F_{a,k})).$$

(Here the fact that we took $\widehat{X}_{k,a}$ to be normal avoids any potential issue in the division process, as $\widehat{X}_{k,a} \cap \mu_k^{-1}(\pi_{k,0}^{-1} \cap_{I \in \mathcal{G}} \tau_I^{-1}(0))$ has the expected codimension = card \mathcal{G} for any family \mathcal{G}).

5.20. Lemma. *Let V be a finite dimensional vector space over \mathbb{C} , $\Psi : V^p \rightarrow \mathbb{C}$ a non-zero alternating multilinear form, and let $m, c \in \mathbb{N}$, $c < m \leq p$, $r = p + c - m \geq 0$. Then the subset $T \subset V^m$ of vectors $(v_1, \dots, v_m) \in V^m$ such that*

$$(*) \quad \Psi(h_1, \dots, h_r, (v_j)_{j \in \mathcal{C}J}) = 0 \quad \text{for all } J \subset \{1, \dots, m\}, |J| = c, \text{ and all } h_1, \dots, h_r \in V,$$

is a closed algebraic subset of codimension $\geq (c+1)(r+1)$.

Proof. A typical example is $\Psi = \det$ on a p -dimensional vector space V , then T consists of m -tuples of vectors of rank $< p - r$, and the assertion concerning the codimension is well known (we will reprove it anyway). In general, the algebraicity of T is obvious. We argue by induction on p , the result being trivial for $p = 1$ (the kernel of a non-zero linear form is indeed of codimension ≥ 1). If K is the kernel of Ψ , i.e. the subspace of vectors $v \in V$ such that $\Psi(h_1, \dots, h_{p-1}, v) = 0$ for all $h_j \in V$, then Ψ induces an alternating multilinear form $\overline{\Psi}$ on V/K , whose kernel is equal to $\{0\}$. The proof is thus reduced to the case when $\text{Ker } \Psi = \{0\}$. Notice that we must have $\dim V \geq p$, otherwise Ψ would vanish. If $\text{card } \mathcal{C}J = m - c = 1$, condition $(*)$ implies that $v_j \in \text{Ker } \Psi = \{0\}$ for all j , hence $\text{codim } T = \dim V^m \geq mp = (c+1)(r+1)$, as desired. Now, assume $m - c \geq 2$, fix $v_m \in V \setminus \{0\}$ and consider the non-zero alternating multilinear form on V^{p-1} such that

$$\Psi'_{v_m}(w_1, \dots, w_{p-1}) := \Psi(w_1, \dots, w_{p-1}, v_m).$$

If $(v_1, \dots, v_m) \in T$, then (v_1, \dots, v_{m-1}) belongs to the set T'_{v_m} associated with the new data $(\Psi'_{v_m}, p-1, m-1, c, r)$. The induction hypothesis implies that $\text{codim } T'_{v_m} \geq (c+1)(r+1)$, and since the projection $T \rightarrow V$ to the first factor admits the T'_{v_m} as its fibers, we conclude that

$$\text{codim } T \cap ((V \setminus \{0\}) \times V^{m-1}) \geq (c+1)(r+1).$$

By permuting the arguments v_j , we also conclude that

$$\text{codim } T \cap (V^{k-1} \times (V \setminus \{0\}) \times V^{m-k}) \geq (c+1)(r+1)$$

for all $k = 1, \dots, m$. The union $\bigcup_k (V^{k-1} \times (V \setminus \{0\}) \times V^{m-k}) \subset V^m$ leaves out only $\{0\} \subset V^m$ whose codimension is at least $mp \geq (c+1)(r+1)$, so Lemma 5.20 follows. \square

5.21. Proposition. *Consider in $U \times \widehat{Z}'_k$ the set Γ of pairs (a, ξ) such that $\sigma_{k,s,\hat{\tau},a,J}(\xi) = 0$ for all choices of $s, \hat{\tau}$ and $J \subset \{0, 1, \dots, N\}$ with $\text{card } J = c$. Then Γ is an algebraic set of dimension*

$$\dim \Gamma \leq \dim S - (c+1)(k+c-N+1) + n+1 + kn.$$

As a consequence, if $(c+1)(k+c-N+1) > n+1+kn$, there exists $a \in U \subset S$ such that the base locus of the family of sections $\sigma_{k,s,\hat{\tau},a,J}$ in $\widehat{X}_{a,k}$ lies over $\bigcup_I X_a \cap \tau_I^{-1}(0)$.

Proof. The idea is similar to [Brot17, Lemma 3.8], but somewhat simpler in the present context. Let us consider a point $\xi \in \widehat{Z}'_k$ and the k -jet $f_{[k]} = \mu_k(\xi) \in Z'_k$, so that $x = f(0) \in Z' = Z \setminus \bigcup_I \tau_I^{-1}(0)$. Let us take the $\hat{\tau}_j$ such that $\hat{\tau}_j(x) \neq 0$. Then, we do not have to pay attention to the non-vanishing factors $\hat{\tau}^{\alpha_J} \tau^{\beta_J}$, and the k -jets of sections m_j and $\hat{\tau}_j^{d-k}$ are invertible near x . Let e_A be a local generator of A near x and $e_{\mathcal{L}}$ a local generator of the invertible sheaf

$$\mathcal{L} = \mu_k^* \mathcal{O}_{Z_k}(k') \otimes \mathcal{O}_{\widehat{Z}'_k}(-F_k)$$

near $\xi \in \widehat{Z}'_k$. Let $J^k \mathcal{O}_{Z,x} = \mathcal{O}_{Z,x} / \mathfrak{m}_{Z,x}^{k+1}$ be the vector space of k -jets of functions on Z at x . By definition of the Wronskian ideal and of the associated divisor F_k , we have a *non-zero* alternating multilinear form

$$\Psi : (J^k \mathcal{O}_{Z,x})^{k+1} \rightarrow \mathbb{C}, \quad (g_0, \dots, g_k) \mapsto \mu_k^* W_k(g_0, \dots, g_k)(\xi) / e_{\mathcal{L}}(\xi).$$

The simultaneous vanishing of our sections at ξ is equivalent to the vanishing of

$$(5.22) \quad \Psi(s_1 \hat{\tau}_1^{d-k} e_A^{-d}, \dots, s_r \hat{\tau}_r^{d-k} e_A^{-d}, (a_j m_j^\delta e_A^{-d})_{j \in \mathbb{C}J})$$

for all (s_1, \dots, s_r) . Since A is very ample and $\rho \geq k$, the power A^ρ generates k -jets at every point $x \in Z$, hence the morphisms

$$H^0(Z, A^\rho) \rightarrow J^k \mathcal{O}_{Z,x}, \quad a \mapsto a m_j^\delta e_A^{-d} \quad \text{and} \quad H^0(Z, A^k) \rightarrow J^k \mathcal{O}_{Z,x}, \quad s \mapsto s \hat{\tau}_j^{d-k} e_A^{-d}$$

are surjective. Lemma 5.20 applied with $r = k + c - N$ and (p, m) replaced by $(k + 1, N + 1)$ implies that the codimension of families $a = (a_0, \dots, a_N) \in S = H^0(Z, A^\rho)^{\oplus(N+1)}$ for which $\sigma_{k,s,\hat{\tau},a,J}(\xi) = 0$ for all choices of $s, \hat{\tau}$ and J is at least $(c+1)(k+c-N+1)$, i.e. the dimension is at most $\dim S - (c+1)(k+c-N+1)$. When we let ξ vary over \hat{Z}'_k which has dimension $(n+1) + kn$ and take into account the fibration $(a, \xi) \mapsto \xi$, the dimension estimate of Proposition 5.21 follows. Under the assumption

$$(5.23) \quad (c+1)(k+c-N+1) > n+1+kn$$

we have $\dim \Gamma < \dim S$, hence the image of the projection $\Gamma \rightarrow S, (a, \xi) \mapsto a$ is a constructible algebraic subset distinct from S . This concludes the proof. \square

Our final goal is to completely eliminate the base locus. Proposition 5.21 indicates that we have to pay attention to the intersections $X_a \cap \tau_I^{-1}(0)$. For $x \in Z$, we let \mathcal{G} be the family of hyperplane sections $\tau_I = 0$ that contain x . We introduce the set $P = \{0, 1, \dots, N\} \setminus \bigcup_{I \in \mathcal{G}} I$ and the smooth intersection

$$Z_{\mathcal{G}} = Z \cap \bigcap_{I \in \mathcal{G}} \tau_I^{-1}(0),$$

so that $N' + 1 := \text{card } P \geq N + 1 - c \text{ card } \mathcal{G}$ and $\dim Z_{\mathcal{G}} = n + 1 - \text{card } \mathcal{G}$. If $a \in U$ is such that $x \in X_a$, we also look at the intersection

$$X_{\mathcal{G},a} = X_a \cap \bigcap_{I \in \mathcal{G}} \tau_I^{-1}(0),$$

which is a smooth hypersurface of $Z_{\mathcal{G}}$. In that situation, we consider Wronskians $W_{k,s,\hat{\tau},a,J}$ as defined above, but we now take $J \subset P$, $\text{card } J = c$, $\mathbb{C}J = P \setminus J$, $r' = k + c - N'$.

5.24. Lemma. *In the above setting, if we assume $\delta > k$, the restriction $W_{k,s,\hat{\tau},a,J}|_{X_{\mathcal{G},a}}$ is still divisible by a monomial $\hat{\tau}^{\alpha_J} \tau^{\beta_J}$ such that*

$$|\alpha_J| + |\beta_J| = (k + c - N')(d - 2k) + (N' + 1 - c)(\delta - k)b + (\delta - k).$$

Therefore, if

$$p' = |\alpha_J| + |\beta_J| - (k + 1)d = (\delta - k) - (k + c - N')2k - (N' + 1 - c)(kb + \rho)$$

as in (5.12), we obtain again holomorphic sections

$$\begin{aligned} \widetilde{W}_{k,s,\hat{\tau},a,J}|_{X_{\mathcal{G},a}} &:= (\hat{\tau}^{\alpha_J} \tau^{\beta_J})^{-1} W_{k,s,\hat{\tau},a,J}|_{X_{\mathcal{G},a}} \in H^0(X_{\mathcal{G},a}, E_{k,k'} T_X^* \otimes A^{-p'}), \\ \sigma_{k,s,\hat{\tau},a,J}|_{\pi_{k,0}^{-1}(X_{\mathcal{G},a})} &\in H^0(\pi_{k,0}^{-1}(X_{\mathcal{G},a}), \mu_{a,k}^*(\mathcal{O}_{X_{a,k}}(k')) \otimes \pi_{k,0}^* A^{-p'}) \otimes \mathcal{O}_{\hat{X}_{a,k}}(-F_{a,k}). \end{aligned}$$

Proof. The arguments are similar to those employed in the proof of Lemma 5.10. Let $f_{[k]} \in X_{a,k}$ be a k -jet such that $f(0) \in X_{\mathcal{G},a}$ (the k -jet need not be entirely contained in $X_{\mathcal{G},a}$). Putting $j_0 = \min(\mathbb{C}J)$, we observe that we have on $X_{\mathcal{G},a}$ an identity

$$a_{j_0} m_{j_0}^\delta = - \sum_{i \in P \setminus \{j_0\}} a_i m_i^\delta = - \sum_{i \in J} a_i m_i^\delta - \sum_{P \setminus (J \cup \{j_0\})} a_i m_i^\delta$$

because $m_i = \prod_{I \ni i} \tau_I = 0$ on $X_{\mathcal{G},a}$ when $i \in \mathbb{C}P = \bigcup_{I \in \mathcal{G}} I$ (one of the factors τ_I is such that $I \in \mathcal{G}$, hence $\tau_I = 0$). If we compose with a germ $t \mapsto f(t)$ such that $f(0) \in X_{\mathcal{G},a}$ (even though f does not necessarily lie entirely in $X_{\mathcal{G},a}$), we get

$$a_{j_0} m_{j_0}^\delta(f(t)) = - \sum_{i \in J} a_i m_i^\delta(f(t)) - \sum_{P \setminus (J \cup \{j_0\})} a_i m_i^\delta(f(t)) + O(t^{k+1})$$

as soon as $\delta > k$. Hence we have an equality for all derivatives $D^\ell(\bullet)$, $\ell \leq k$ at $t = 0$, and

$$W_{k,s,\hat{\tau},a,J|X_{\mathcal{G},a}}(f|_k) = - \sum_{i \in J} W_k(s_1 \hat{\tau}_1^{d-k}, \dots, s_{r'} \hat{\tau}_{r'}^{d-k}, a_i m_i^\delta, (a_j m_j^\delta)_{j \in P \setminus (J \cup \{j_0\})})|_{X_{\mathcal{G},a}}(f|_k).$$

Then, again, $\tau_J^{\delta-k}$ is a new additional common factor of all terms in the sum, and we conclude as in Lemma 5.10 and Corollary 5.11. \square

Now, we analyze the base locus of these new sections on

$$\bigcup_{a \in U} \mu_{a,k}^{-1} \pi_{k,0}^{-1}(X_{\mathcal{G},a}) \subset \mu_k^{-1} \pi_{k,0}^{-1}(Z_{\mathcal{G}}) \subset \widehat{Z}_k.$$

As x runs in $Z_{\mathcal{G}}$ and $N' < N$, Lemma 5.20 shows that (5.23) can be replaced by the less demanding condition

$$(5.23') \quad (c+1)(k+c-N'+1) > n+1 - \text{card } \mathcal{G} + kn = \dim \mu_k^{-1} \pi_{k,0}^{-1}(Z_{\mathcal{G}}).$$

A proof entirely similar to that of Proposition 5.21 shows that for a generic choice of $a \in U$, the base locus of these sections on $\widehat{X}_{\mathcal{G},a,k}$ projects onto $\bigcup_{I \in \mathbb{C}\mathcal{G}} X_{\mathcal{G},a} \cap \tau_I^{-1}(0)$. Arguing inductively on $\text{card } \mathcal{G}$, the base locus can be shrunk step by step down to empty set (but it is in fact sufficient to stop when $X_{\mathcal{G},a} \cap \tau_I^{-1}(0)$ reaches dimension 0).

5.E. NEFNESS AND AMPLENESS OF APPROPRIATE TAUTOLOGICAL LINE BUNDLES

At this point, we have produced a smooth family $\mathcal{X}_S \rightarrow U \subset S$ of particular hypersurfaces in Z , namely $X_a = \{\sigma_a(z) = 0\}$, $a \in U$, for which a certain ‘‘tautological’’ line bundle has an empty base locus for sufficiently general coefficients:

5.25. Corollary. *Under condition (5.23) and the hypothesis $p > 0$ in (5.12), the following properties hold.*

(a) *The line bundle*

$$\mathcal{L}_a := \mu_{a,k}^*(\mathcal{O}_{X_{a,k}}(k') \otimes \pi_{k,0}^* A^{-1}) \otimes \mathcal{O}_{\widehat{X}_{a,k}}(-F_{a,k})$$

is nef on $\widehat{X}_{a,k}$ for general $a \in U'$, where $U' \subset U$ is a dense Zariski open set.

(b) *Let $\Delta_a = \sum_{2 \leq \ell \leq k} \lambda_\ell D_{a,\ell}$ be a positive rational combination of vertical divisors of the Semple tower and $q \in \mathbb{N}$, $q \gg 1$, an integer such that*

$$\mathcal{L}'_a := \mathcal{O}_{X_{a,k}}(1) \otimes \mathcal{O}_{a,k}(-\Delta_a) \otimes \pi_{k,0}^* A^q$$

is ample on $X_{a,k}$. Then the \mathbb{Q} -line bundle

$$\mathcal{L}_{a,\varepsilon,\eta} := \mu_{a,k}^*(\mathcal{O}_{X_{a,k}}(k') \otimes \mathcal{O}_{X_{a,k}}(-\varepsilon \Delta_a) \otimes \pi_{k,0}^* A^{-1+q\varepsilon}) \otimes \mathcal{O}_{\widehat{X}_{a,k}}(-(1+\varepsilon\eta)F_{a,k})$$

is ample on $\widehat{X}_{a,k}$ for $a \in U'$, for some $q \in \mathbb{N}$ and $\varepsilon, \eta \in \mathbb{Q}_{>0}$ arbitrarily small.

Proof. (a) This would be obvious if we had global sections generating \mathcal{L}_a on the whole of $\widehat{X}_{a,k}$, but our sections are only defined on a stratification of $\widehat{X}_{a,k}$. In any case, if $C \subset \widehat{X}_{a,k}$ is an irreducible curve, we take a maximal family \mathcal{G} such that $C \subset X_{\mathcal{G},a,k}$. Then, by what we have seen, for $a \in U$ general enough, we can find global sections of \mathcal{L}_a on $\widehat{X}_{\mathcal{G},a,k}$ such that C is not contained in their base locus. Hence $\mathcal{L}_a \cdot C \geq 0$ and \mathcal{L}_a is nef for a in a dense Zariski open set $U' \subset U$.

(b) The existence of Δ_a and q follows from Proposition 3.19 and Corollary 3.21, which even provide universal values for λ_ℓ and q . After taking the blow up $\mu_{a,k} : \widehat{X}_{a,k} \rightarrow X_{a,k}$ (cf. (4.8)), we infer that

$$\mathcal{L}'_{a,\eta} := \mu_{a,k}^* \mathcal{L}'_a \otimes \mathcal{O}_{\widehat{X}_{a,k}}(-\eta F_{a,k}) = \mu_{a,k}^* (\mathcal{O}_{X_{a,k}}(1) \otimes \mathcal{O}_{X_{a,k}}(-\Delta_a) \otimes \pi_{k,0}^* A^q) \otimes \mathcal{O}_{\widehat{X}_{a,k}}(-\eta F_{a,k})$$

is ample for $\eta > 0$ small. The result now follows by taking a combination

$$\mathcal{L}_{a,\varepsilon,\eta} = \mathcal{L}'_a^{1-\varepsilon/k'} \otimes (\mathcal{L}'_{a,\eta})^\varepsilon. \quad \square$$

5.26. Corollary. *Let $\mathcal{X} \rightarrow \Omega$ be the universal family of hypersurfaces $X_\sigma = \{\sigma(z) = 0\}$, $\sigma \in \Omega$, where $\Omega \subset P(H^0(Z, A^d))$ is the dense Zariski open set over which the family is smooth. On the “Wronskian blow-up” $\widehat{X}_{\sigma,k}$ of $X_{\sigma,k}$, let us consider the line bundle*

$$\mathcal{L}_{\sigma,\varepsilon,\eta} := \mu_{\sigma,k}^* (\mathcal{O}_{X_{\sigma,k}}(k') \otimes \mathcal{O}_{X_{\sigma,k}}(-\varepsilon \Delta_\sigma) \otimes \pi_{k,0}^* A^{-1+q\varepsilon}) \otimes \mathcal{O}_{\widehat{X}_{\sigma,k}}(-(1+\varepsilon\eta)F_{\sigma,k})$$

associated with the same choice of constants as in Cor. 5.25. Then $\mathcal{L}'_{\sigma,\varepsilon,\eta}$ is ample on $\widehat{X}_{\sigma,k}$ for σ in a dense Zariski open set $\Omega' \subset \Omega$.

Proof. By 5.25 (b), we can find $\sigma_0 \in H^0(Z, A^d)$ such that $X_{\sigma_0} = \sigma_0^{-1}(0)$ is smooth and $\mathcal{L}_{\sigma_0,\varepsilon,\eta}^m$ is an ample line bundle on $\widehat{X}_{\sigma_0,k}$ ($m \in \mathbb{N}^*$). As ampleness is a Zariski open condition, we infer that $\mathcal{L}_{\sigma,\varepsilon,\eta}^m$ remains ample for a general section $\sigma \in H^0(Z, A^d)$, i.e. for $[\sigma]$ in some Zariski open set $\Omega' \subset \Omega$. Since $\mu_{\sigma,k}(F_{\sigma,k})$ is contained in the vertical divisor of $X_{\sigma,k}$, we conclude by Corollary 3.27 that X_σ is Kobayashi hyperbolic for $[\sigma] \in \Omega$. \square

5.F. FINAL CONCLUSION AND COMPUTATION OF DEGREE BOUNDS

At this point, we fix our integer parameters to meet all conditions that have been found. We must have $N \geq c(n+1)$ by Lemma 5.7, and for such a large value of N , condition (5.23) can hold only when $c \geq n$, so we take $c = n$ and $N = n(n+1)$. Inequality (5.23) then requires k large enough, $k = n^3 + n^2 + 1$ being the smallest possible value. We find

$$b = \binom{N}{c-1} = \binom{n^2+n}{n-1} = n \frac{(n^2+n) \dots (n^2+2)}{n!}.$$

We have $n^2 + k = n^2(1 + k/n^2) < n^2 \exp(k/n^2)$ and by Stirling's formula, $n! > \sqrt{2\pi n} (n/e)^n$, hence

$$b < \frac{n^{2n-1} \exp((2+\dots+n)/n^2)}{\sqrt{2\pi n} (n/e)^n} < \frac{e^{n+\frac{1}{2}+\frac{1}{2n}}}{\sqrt{2\pi}} n^{n-\frac{3}{2}}.$$

Finally, we divide $d-k$ by b , get in this way $d-k = b\delta + \lambda$, $0 \leq \lambda < b$, and put $\rho = \lambda + k \geq k$. Then $\delta + 1 \geq (d-k+1)/b$ and formula (5.12) yields

$$\begin{aligned} p &= (\delta - k) - (n^3 + 1)2k - (n^2 + 2n + 1)(kb + \rho) \\ &\geq (d - k + 1)/b - 1 - (2n^3 + 3)k - (n^2 + 2n + 1)(kb + k + b - 1), \end{aligned}$$

therefore $p > 0$ is achieved as soon as

$$d \geq d_n = k + b(1 + (2n^3 + 3)k + (n^2 + 2n + 1)(kb + k + b - 1)),$$

where

$$k = n^3 + n^2 + 1, \quad b = \binom{n^2 + n}{n - 1}.$$

The dominant term in d_n is $k(n^2 + 2n + 1)b^2 \sim e^{2n+1}n^{2n+2}/2\pi$. By means of more precise numerical calculations and of Stirling's asymptotic expansion for $n!$, one can check in fact that $d_n \leq \lfloor (n+4)(en)^{2n+1}/2\pi \rfloor$ for $n \geq 4$ (which is also an equivalent and a close approximation as $n \rightarrow +\infty$), while $d_1 = 61$, $d_2 = 6685$, $d_3 = 2825761$. We can now state the main result of this section.

5.27. Theorem. *Let Z be a projective $(n + 1)$ -dimensional manifold and A a very ample line bundle on Z . Then, for a general section $\sigma \in H^0(Z, A^d)$ and $d \geq d_n$, the hypersurface $X_\sigma = \sigma^{-1}(0)$ is Kobayashi hyperbolic. The bound d_n for the degree can be taken to be*

$$d_n = \lfloor (n + 4)(en)^{2n+1}/2\pi \rfloor \quad \text{for } n \geq 4,$$

and for $n \leq 3$, one can take $d_1 = 4$, $d_2 = 6685$, $d_3 = 2825761$.

For $n = 1$, we have already seen in § 4.B that $d_1 = 4$ works (rather than the insane value $d_1 = 61$). A simpler (and less refined) choice is $\tilde{d}_n = \lfloor \frac{1}{3}(en)^{2n+2} \rfloor$, which is valid for all n . These bounds are only slightly weaker than the ones found by Ya Deng in his PhD thesis [Deng16, Deng17], namely $\tilde{d}_n = O(n^{2n+6})$.

5.G. FURTHER COMMENTS

5.28. Our bound d_n is rather large, but just as in Ya Deng's effective approach of Brotbek's theorem [Deng17], the bound holds for a property that looks substantially stronger than hyperbolicity, namely the ampleness of the pull-back of some (twisted) jet bundle $\mu_k^* \mathcal{O}_{\widehat{X}_k}(a_\bullet) \otimes \mathcal{O}_{\widehat{X}_k}(-F'_k)$. It is certainly desirable to look for more general jet differentials than Wronskians, and to relax the positivity demands on tautological line bundles to ensure hyperbolicity (see e.g. [Dem14]). However, the required calculations appear to be much more involved.

5.29. After this chapter was written, Riedl and Yang [RiYa18] proved the important and somewhat surprising result that the lower bound estimates $d_{\text{GG}}(n)$ and $d_{\text{Kob}}(n)$, respectively for the Green-Griffiths-Lang and Kobayashi conjectures for general hypersurfaces in \mathbb{P}^{n+1} , can be related by $d_{\text{Kob}}(n) := d_{\text{GG}}(2n - 2)$. This should be understood in the sense that a solution of the generic $(2n - 2)$ -dimensional Green-Griffiths conjecture for $d \geq d_{\text{GG}}(2n - 2)$ implies a solution of the n -dimensional Kobayashi conjecture for the same lower bound. We refer to [RiYa18] for the precise statement, which requires an extra assumption on the algebraic dependence of the Green-Griffiths locus with respect to a variation of coefficients in the defining polynomials. In combination with [DMR10], this gives a completely new proof of the Kobayashi conjecture, and the order 1 bound $d_{\text{GG}}(n) = O(\exp(n^{1+\varepsilon}))$ of [Dem12] implies a similar bound $d_{\text{Kob}}(n) = O(\exp(n^{1+\varepsilon}))$ for the Kobayashi conjecture – just a little bit weaker than what our direct proof gave (Theorem 5.26). In [MeTa19], Merker and Ta were able to improve the Green-Griffiths bound to $d_{\text{GG}}(n) = o(\sqrt{n} \log n)^n$, using a strengthening of Darondeau's estimates [Dar16a, Dar16b], along with very delicate calculations. The Riedl-Yang result then implies $d_{\text{Kob}}(n) = O((n \log n)^{n+1})$, which is the best bound known at this time.

5.30. In the unpublished preprint [Dem15], we introduced an alternative strategy for the proof of the Kobayashi conjecture which appears to be still incomplete at this point. We nevertheless hope that a refined version could one day lead to linear bounds such as $d_{\text{Kob}}(n) = 2n + 1$. The rough idea was to establish a k -jet analogue of Claire Voisin's proof [Voi96] of the Clemens conjecture. Unfortunately, Lemma 5.1.18 as stated in [Dem15] is incorrect – the assertion concerning the Δ divisor introduced there simply does not hold. It is however conceivable that a weaker statement holds, in the form of a control of the degree of the divisor Δ , and in a way that would still be sufficient to imply similar consequences for the generic positivity of tautological jet bundles.

5.31. In [Ber19], G. Bérczi stated a positivity conjecture for Thom polynomials of Morin singularities (see also [BeSz12]), and showed that it would imply a polynomial bound $d_n = 2n^9 + 1$ for the generic hyperbolicity of hypersurfaces.

5.32. In September 2019, Bérczi and Kirwan [BeKi19] introduced new deep ideas in non-reductive geometric invariant theory that actually lead to polynomial bounds for the Kobayashi conjecture. Their technique is based on the use of alternative compactifications for the jet spaces.

REFERENCES

- [ASS97] Arrondo, E., Sols, I., Speiser, R.: *Global moduli for contacts*. Ark. Mat. **35** (1997), 1–57.
- [Ber19] Bérczi, G.: *Thom polynomials and the Green-Griffiths conjecture for hypersurfaces with polynomial degree*. Intern. Math. Res. Not. **2019/22** (2019), 7037–7092.
- [BeKi12] Bérczi, G., Kirwan, F.: *A geometric construction for invariant jet differentials*. Surveys in Diff. Geom., Vol XVII (2012), 79–126.
- [BeKi19] Bérczi, G., Kirwan, F.: *Non-reductive geometric invariant theory and hyperbolicity*. Preprint in arXiv, Math.AG:1909.11417.
- [BeSz12] Bérczi, G., Szenes, A.: *Thom polynomials of Morin singularities*. Annals of Math. **175** (2012), 567–629.
- [Bro78] Brody, R.: *Compact manifolds and hyperbolicity*. Trans. Amer. Math. Soc. **235** (1978), 213–219.
- [Brot17] Brotbek, D.: *On the hyperbolicity of general hypersurfaces*. Publications mathématiques de l’IHÉS, **126** (2017), 1–34.
- [BrDa18] Brotbek, D., Darondeau, L.: *Complete intersection varieties with ample cotangent bundles*. Inventiones Math. **212** (2018), 913–940.
- [Cle86] Clemens, H.: *Curves on generic hypersurfaces*. Ann. Sci. Ec. Norm. Sup. **19** (1986), 629–636.
- [CoKe94] Colley, S.J., Kennedy, G.: *The enumeration of simultaneous higher order contacts between plane curves*. Compositio Math. **93** (1994), 171–209.
- [Coll88] Collino, A.: *Evidence for a conjecture of Ellingsrud and Strømme on the Chow ring of $\mathbf{Hilb}_d(\mathbb{P}^2)$* . Illinois J. Math. **32** (1988), 171–210.
- [Dar16a] Darondeau, L.: *Fiber integration on the Demailly tower*. Ann. Inst. Fourier **66** (2016), 29–54.
- [Dar16b] Darondeau, L.: *On the logarithmic Green-Griffiths conjecture*. Int. Math. Res. Not. (2016), no. 6, 1871–1923.
- [Dem85] Demailly, J.-P.: *Champs magnétiques et inégalités de Morse pour la d'' -cohomologie*. Ann. Inst. Fourier (Grenoble) **35** (1985) 189–229.
- [Dem95] Demailly, J.-P.: *Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials*. AMS Summer School on Algebraic Geometry, Santa Cruz 1995, Proc. Symposia in Pure Math., ed. by J. Kollár and R. Lazarsfeld, Amer. Math. Soc., Providence, RI (1997), 285–360.
- [Dem97] Demailly, J.-P.: *Variétés hyperboliques et équations différentielles algébriques*. Gaz. Math. **73** (juillet 1997) 3–23.
- [Dem07a] Demailly, J.-P.: *Structure of jet differential rings and holomorphic Morse inequalities*. Talk at the CRM Workshop “The geometry of holomorphic and algebraic curves in complex algebraic varieties”, Montréal, May 2007.
- [Dem07b] Demailly, J.-P.: *On the algebraic structure of the ring of jet differential operators*. Talk at the conference “Effective aspects of complex hyperbolic varieties”, Aber Wrac’h, France, September 10–14, 2007.
- [Dem11] Demailly, J.-P.: *Holomorphic Morse Inequalities and the Green-Griffiths-Lang Conjecture*. Pure and Applied Math. Quarterly **7** (2011), 1165–1208.
- [Dem14] Demailly, J.-P.: *Towards the Green-Griffiths-Lang conjecture*. Conference “Analysis and Geometry”, Tunis, March 2014, in honor of Mohammed Salah Baouendi, ed. by A. Baklouti, A. El Kacimi, S. Kallel, N. Mir, Springer (2015), 141–159.
- [Dem15] Demailly, J.-P.: *Proof of the Kobayashi conjecture on the hyperbolicity of very general hypersurfaces*. arXiv:1501.07625, math.CV, unpublished.
- [Dem18] Demailly, J.-P.: *Recent results on the Kobayashi and Green-Griffiths-Lang conjectures*, Contribution to the 16th Takagi lectures in celebration of the 100th anniversary of K.Kodaira’s birth, 2015, Japanese Journal of Mathematics **15** (2020), 1–120.
- [DeEG97] Demailly, J.-P., El Goul, J.: *Connexions méromorphes projectives et variétés algébriques hyperboliques*. C. R. Acad. Sci. Paris, Série I, Math. (janvier 1997).
- [DeEG00] Demailly, J.-P., El Goul, J.: *Hyperbolicity of generic surfaces of high degree in projective 3-space*. Amer. J. Math. **122** (2000), 515–546.
- [DeLS94] Demailly, J.-P., Lempert L., Shiffman, B.: *Algebraic approximation of holomorphic maps from Stein domains to projective manifolds*. Duke Math. J. **76** (1994) 333–363.

- [Deng16] Deng, Ya.: *Effectivity in the hyperbolicity related problems*. Chap. 4 of the PhD memoir “Generalized Okounkov Bodies, Hyperbolicity-Related and Direct Image Problems” defended on June 26, 2017 at Université Grenoble Alpes, Institut Fourier, arXiv:1606.03831, math.CV.
- [Deng17] Deng, Ya.: *On the Diverio-Trapani Conjecture* arXiv:1703.07560, math.CV.
- [DTH16] Dinh, Tuan Huynh: *Construction of hyperbolic hypersurfaces of low degree in $\mathbb{P}^n(\mathbb{C})$* . Int. J. Math. **27** (2016) 1650059 (9 pages).
- [DMR10] Diverio, S., Merker, J., Rousseau, E.: *Effective algebraic degeneracy*. Invent. Math. **180** (2010) 161–223.
- [DR15] Diverio, S., Rousseau, E.: *The exceptional set and the Green–Griffiths locus do not always coincide*. Enseign. Math. **61** (2015) 417–452.
- [DT10] Diverio, S., Trapani, S.: *A remark on the codimension of the Green–Griffiths locus of generic projective hypersurfaces of high degree*. J. Reine Angew. Math. **649** (2010) 55–61.
- [Dol81] Dolgachev, I.: *Weighted projective varieties*. Proceedings Polish–North Amer. Sem. on Group Actions and Vector Fields, Vancouver, 1981, J.B. Carrels editor, Lecture Notes in Math. **956**, Springer-Verlag (1982), 34–71.
- [Duv04] Duval J.: *Une sextique hyperbolique dans $\mathbb{P}^3(\mathbb{C})$* . Math. Ann. **330** (2004) 473–476.
- [Ein88] Ein L.: *Subvarieties of generic complete intersections*. Invent. Math. **94** (1988), 163–169.
- [Ein91] Ein L.: *Subvarieties of generic complete intersections, II*. Math. Ann. **289** (1991), 465–471.
- [Fuj74] Fujimoto H.: *On meromorphic maps into the complex projective space*. J. Math. Soc. Japan **26** (1974), 272–288.
- [Ghe41] Gherardelli, G.: *Sul modello minimo della varieta degli elementi differenziali del 2° ordine del piano proiettivo*. Atti Accad. Italia. Rend., Cl. Sci. Fis. Mat. Nat. (7) **2** (1941), 821–828.
- [Gr75] Green M.: *Some Picard theorems for holomorphic maps to algebraic varieties*. Amer. J. Math. **97** (1975), 43–75.
- [GrGr80] Green, M., Griffiths, P.: *Two applications of algebraic geometry to entire holomorphic mappings*. The Chern Symposium 1979, Proc. Internal. Sympos. Berkeley, CA, 1979, Springer-Verlag, New York (1980), 41–74.
- [Har77] Hartshorne, R.: *Algebraic geometry*. Springer-Verlag, Berlin (1977).
- [Ko67a] Kobayashi, S.: *Intrinsic metrics on complex manifolds*. Bull. Amer. Math. Soc. **73** (3) (1967), 347–349.
- [Ko67b] Kobayashi, S.: *Invariant distances on complex manifolds and holomorphic mappings*. J. Math. Soc. Japan **19** (1967), 460–480.
- [Kob70] Kobayashi, S.: *Hyperbolic manifolds and holomorphic mappings*. Marcel Dekker, New York 1970, 2nd Edition, World Sci. 2005.
- [Kob76] Kobayashi, S.: *Intrinsic distances, measures and geometric function theory*. Bull. Amer. Math. Soc. **82** (1976), 357–416.
- [Kob98] Kobayashi, S.: *Hyperbolic Complex Spaces*. Grundle. der Math. Wissen. vol. 318, Springer-Verlag, Berlin-Heidelberg, 1998.
- [LaTh96] Laksov, D., Thorup, A.: *These are the differentials of order n* . Trans. Amer. Math. Soc. **351** (1999), 1293–1353.
- [Lang86] Lang, S.: *Hyperbolic and Diophantine analysis*. Bull. Amer. Math. Soc. **14** (1986), 159–205.
- [MaNo96] Masuda, K., Noguchi, J.: *A construction of hyperbolic hypersurface of $\mathbb{P}^n(\mathbb{C})$* . Math. Ann. **304** (1996), 339–362.
- [McQ99] McQuillan, M.: *Holomorphic curves on hyperplane sections of 3-folds*. Geom. Funct. Anal. **9** (1999), 370–392.
- [Mer08] Merker, J.: *Jets de Demailly–Semple d’ordres 4 et 5 en dimension 2*. Int. J. Contemp. Math. Sci. **3-18** (2008), 861–933.
- [Mer10] Merker, J.: *Application of computational invariant theory to Kobayashi hyperbolicity and to Green–Griffiths algebraic degeneracy* J. of Symbolic Computation, **45** (2010), 986–1074.
- [MeTa19] Merker, J., Ta, The-Anh: *Degrees $d \geq \sqrt{n} \log n^n$ and $d \geq (n \log n)^n$ in the Conjectures of Green–Griffiths and of Kobayashi*. arXiv:1901.04042, math.AG.
- [Nad89] Nadel, A.: *Hyperbolic surfaces in \mathbb{P}^3* . Duke Math. J. **58** (1989), 749–771.
- [Pac04] Pacienza, G.: *Subvarieties of general type on a general projective hypersurface*. Trans. Amer. Math. Soc. **356** (2004), 2649–2661.

- [Pau08] Păun, M.: *Vector fields on the total space of hypersurfaces in the projective space and hyperbolicity*. Math. Ann. **340** (2008) 875–892.
- [RiYa18] Riedl, E., Yang, D.: *Applications of a grassmannian technique in hypersurfaces*. June 2018, arXiv:1806.02364, math.AG.
- [Rou06] Rousseau, E.: *Étude des jets de Demailly-Semple en dimension 3*. Ann. Inst. Fourier (Grenoble) **56** (2006), 397–421.
- [Roy71] Royden, H.: *Remarks on the Kobayashi metric*. Proc. Maryland Conference on Several Complex Variables, Lecture Notes, Vol. 185, Springer-Verlag, Berlin (1971).
- [Roy74] Royden, H.: *The extension of regular holomorphic maps*. Proc. Amer. Math. Soc. **43** (1974), 306–310.
- [Sem54] Semple, J.G.: *Some investigations in the geometry of curves and surface elements*. Proc. London Math. Soc. (3) **4** (1954), 24–49.
- [ShZa02] Shiffman, B., Zaidenberg, M.: *Hyperbolic hypersurfaces in \mathbb{P}^n of Fermat-Waring type*. Proc. Amer. Math. Soc. **130** (2002) 2031–2035.
- [Siu87] Siu, Y.T.: *Defect relations for holomorphic maps between spaces of different dimensions*. Duke Math. J. **55** (1987), 213–251.
- [Siu97] Siu, Y.T.: *A proof of the general Schwarz lemma using the logarithmic derivative lemma*. Personal communication, April 1997.
- [Siu15] Siu, Y.T.: *Hyperbolicity of generic high-degree hypersurfaces in complex projective spaces*. Inventiones Math. **202** (2015) 1069–1166.
- [SiYe96] Siu, Y.T., Yeung, S.K.: *Hyperbolicity of the complement of a generic smooth curve of high degree in the complex projective plane*. Invent. Math. **124** (1996), 573–618.
- [SiYe97] Siu, Y.T., Yeung, S.K.: *Defects for ample divisors of Abelian varieties, Schwarz lemma and hyperbolic surfaces of low degree*. Amer. J. Math. **119** (1997), 1139–1172.
- [Toda71] Toda N.: *On the functional equation $\sum_{i=0}^p a_i f_i^{n_i} = 1$* . Tôhoku Math. J. (2) **23** (1971), 289–299.
- [Ven96] Venturini S.: *The Kobayashi metric on complex spaces*. Math. Ann. **305** (1996), 25–44.
- [Voi96] Voisin, C.: *On a conjecture of Clemens on rational curves on hypersurfaces*. J. Diff. Geom. **44** (1996) 200–213, Correction: J. Diff. Geom. **49** (1998), 601–611.
- [Xie18] Xie, S.-Y.: *On the ampleness of the cotangent bundles of complete intersections*. Inventiones Math. **212** (2018) 941–996.
- [Zai87] Zaidenberg, M.: *The complement of a generic hypersurface of degree $2n$ in $\mathbb{C}P^n$ is not hyperbolic*. Siberian Math. J. **28** (1987), 425–432.
- [Zai93] Zaidenberg, M.: *Hyperbolicity in projective spaces*. International Symposium on Holomorphic mappings, Diophantine Geometry and Related topics, R.I.M.S. Lecture Notes ser. **819**, R.I.M.S. Kyoto University (1993), 136–156.

(version of January 21, 2021, printed on January 21, 2021, 16:33)

Jean-Pierre Demailly

Université Grenoble Alpes, Institut Fourier (Mathématiques)

UMR 5582 du C.N.R.S., 100 rue des Maths, 38610 Gières, France

e-mail: jean-pierre.demailly@univ-grenoble-alpes.fr