



INSTITUT DE FRANCE
Académie des sciences



A sharp lower bound for the log canonical threshold of an isolated plurisubharmonic singularity

Jean-Pierre Demailly / Phạm Hoàng Hiệp

Institut Fourier, Université de Grenoble I, France

GAEL XX +

June 28, 2012

Jean-Pierre Demailly / Phạm Hoàng Hiệp

A sharp lower bound for the log canonical threshold

Singularities of plurisubharmonic (psh) functions

Goal: study **local singularities** of a plurisubharmonic function φ on a neighborhood of a point in \mathbb{C}^n .

$\varphi : X \rightarrow [-\infty, +\infty[$ upper semicont. / mean value inequality.

Poles : $\varphi^{-1}(-\infty)$ (not always closed, sometimes fractal)

Algebraic setting:

$$\varphi(z) = \frac{1}{2} \log(|g_1|^2 + \dots + |g_N|^2)$$

associated to some ideal $\mathcal{I} = (g_1, \dots, g_N) \subset \mathcal{O}_{X,p}$ of holomorphic (algebraic) functions on some complex variety X .

More generally: consider a sequence $(\mathcal{I}_k)_{k \in \mathbb{N}}$ of such ideals, with

$$\mathcal{I}_k \mathcal{I}_\ell \subset \mathcal{I}_{k+\ell}$$

Try to understand “ $\lim(\mathcal{I}_k)^{1/k}$ ” (Lazarsfeld, Ein, Mustață...)

Jean-Pierre Demailly / Phạm Hoàng Hiệp

A sharp lower bound for the log canonical threshold

Lelong numbers and log canonical thresholds

The easiest way of measuring singularities of psh functions is by using **Lelong numbers**:

$$\nu(\varphi, p) = \liminf_{z \rightarrow p} \frac{\varphi(z)}{\log |z - p|}.$$

Example :

$$\varphi(z) = \frac{1}{2} \log(|g_1|^2 + \dots + |g_N|^2) \Rightarrow \nu(\varphi, p) = \min \text{ord}_p(g_j) \in \mathbb{N}.$$

Another useful invariant is the **log canonical threshold**.

Definition

Let X be a complex manifold, $p \in X$, and φ be a plurisubharmonic function defined on X . The **log canonical threshold** or *complex singularity exponent* of φ at p is defined by

$$c_p(\varphi) = \sup \{c \geq 0 : e^{-2c\varphi} \text{ is } L^1 \text{ on a neighborhood of } p\}.$$

log canonical threshold of coherent ideals

For simplicity we will take here $p = 0$ and denote

$$c(\varphi) = c_0(\varphi).$$

The log canonical threshold is a subtle invariant !

Calculation in the case of analytic singularities : take

$$\varphi(z) = \frac{1}{2} \log(|g_1|^2 + \dots + |g_N|^2), \quad \mathcal{J} = (g_1, \dots, g_N).$$

Then by Hironaka, \exists modification $\mu : \tilde{X} \rightarrow X$ such that

$$\mu^* \mathcal{J} = (g_1 \circ \mu, \dots, g_N \circ \mu) = \mathcal{O}(-\sum a_j E_j)$$

for some normal crossing divisor. Let $\mathcal{O}(\sum b_j E_j)$ be the divisor of $\text{Jac}(\mu)$. We have

$$c(\varphi) = \min_{E_j, \mu(E_j) \ni 0} \frac{1 + b_j}{a_j} \in \mathbb{Q}_+^*.$$

Proof of the formula for the log canonical threshold

In fact, we have to find the supremum of $c > 0$ such that

$$I = \int_{V \ni 0} \frac{d\lambda(z)}{(|g_1|^2 + \dots + |g_N|^2)^c} < +\infty.$$

Let us perform the change of variable $z = \mu(w)$. Then

$$d\lambda(z) = |\text{Jac}(\mu)(w)|^2 \sim \left| \prod w_j^{b_j} \right|^2 d\lambda(w)$$

with respect to coordinates on the blow-up \tilde{V} of V , and

$$I \sim \int_{\tilde{V}} \frac{|\prod w_j^{b_j}|^2 d\lambda(w)}{|\prod w_j^{a_j}|^{2c}}$$

so convergence occurs if $ca_j - b_j < 1$ for all j .

Notations and basic facts

- A domain $\Omega \subset \mathbb{C}^n$ is called *hyperconvex* if $\exists \psi \in \mathcal{P}SH(\Omega)$, $\psi \leq 0$, such that $\{z : \psi(z) < c\} \Subset \Omega$ for all $c < 0$.
- $\mathcal{E}_0(\Omega) = \left\{ \varphi \in \mathcal{P}SH \cap L^\infty(\Omega) : \lim_{z \rightarrow \partial\Omega} \varphi(z) = 0, \int_{\Omega} (dd^c \varphi)^n < +\infty \right\}$
- $\mathcal{F}(\Omega) = \left\{ \varphi \in \mathcal{P}SH(\Omega) : \exists \mathcal{E}_0(\Omega) \ni \varphi_p \searrow \varphi, \text{ and } \sup_{p \geq 1} \int_{\Omega} (dd^c \varphi_p)^n < +\infty \right\}$,
- $\tilde{\mathcal{E}}(X) = \{ \varphi \in \mathcal{P}SH(X) \text{ locally in } \mathcal{F}(\Omega) \text{ mod } C^\infty(\Omega) \}$

Theorem (U. Cegrell)

$\tilde{\mathcal{E}}(X)$ is the largest subclass of psh functions defined on a complex manifold X for which the complex Monge-Ampère operator is **locally well-defined**.

Intermediate Lelong numbers

Set here $d^c = \frac{i}{2\pi}(\bar{\partial} - \partial)$ so that $dd^c = \frac{i}{\pi}\partial\bar{\partial}$. If $\varphi \in \tilde{\mathcal{E}}(\Omega)$ and $0 \in \Omega$, the products $(dd^c\varphi)^j$ are well defined and one can consider the Lelong numbers

$$e_j(\varphi) = \nu((dd^c\varphi)^j, 0).$$

In other words

$$e_j(\varphi) = \int_{\{0\}} (dd^c\varphi)^j \wedge (dd^c \log \|z\|)^{n-j}.$$

One has $e_0(\varphi) = 1$ and $e_1(\varphi) = \nu(\varphi, 0)$ (usual Lelong number). When

$$\varphi(z) = \frac{1}{2} \log(|g_1|^2 + \dots + |g_N|^2),$$

one has $e_j(\varphi) \in \mathbb{N}$.

The main result

Main Theorem (Demailly & Phạm)

Let $\varphi \in \tilde{\mathcal{E}}(\Omega)$. If $e_1(\varphi) = 0$, then $c(\varphi) = \infty$.
Otherwise, we have

$$c(\varphi) \geq \sum_{j=0}^{n-1} \frac{e_j(\varphi)}{e_{j+1}(\varphi)}.$$

The lower bound improves a classical result of H. Skoda (1972), according to which

$$\frac{1}{e_1(\varphi)} \leq c(\varphi) \leq \frac{n}{e_1(\varphi)}.$$

Remark: The above theorem is optimal, with equality for

$$\varphi(z) = \log(|z_1|^{a_1} + \dots + |z_n|^{a_n}), \quad 0 < a_1 \leq a_2 \leq \dots \leq a_n.$$

Then $e_j(\varphi) = a_1 \dots a_j$, $c(\varphi) = \frac{1}{a_1} + \dots + \frac{1}{a_n}$.

Geometric applications

The log canonical threshold has a lot of applications. It is essentially a local version of Tian's invariant, which determines a sufficient condition for the **existence of Kähler-Einstein metrics**.

Another important application is to **birational rigidity**.

Theorem (Pukhlikov 1998, Corti 2000, de Fernex 2011)

Let X be a smooth hypersurface of degree d in $\mathbb{C}\mathbb{P}^{n+1}$. Then if $d = n + 1$, **$\text{Bir}(X) \simeq \text{Aut}(X)$**

It was first shown by Manin-Iskovskih in the early 70's that a 3-dim quartic in $\mathbb{C}\mathbb{P}^4$ ($n = 3, d = 4$) is not rational.

Question

For $3 \leq d \leq n + 1$, when is it true that $\text{Bir}(X) \simeq \text{Aut}(X)$ (birational rigidity) ?

Lemma 1

Lemma 1

Let $\varphi \in \tilde{\mathcal{E}}(\Omega)$ and $0 \in \Omega$. Then we have that

$$e_j(\varphi)^2 \leq e_{j-1}(\varphi)e_{j+1}(\varphi),$$

for all $j = 1, \dots, n - 1$.

In other words $j \mapsto \log e_j(\varphi)$ is convex, thus we have $e_j(\varphi) \geq e_1(\varphi)^j$ and the ratios $e_{j+1}(\varphi)/e_j(\varphi)$ are increasing.

Corollary

If $e_1(\varphi) = \nu(\varphi, 0) = 0$, then $e_j(\varphi) = 0$ for $j = 1, 2, \dots, n - 1$.

A hard conjecture by V. Guedj and A. Rashkovskii (~ 1998) states that $\varphi \in \tilde{\mathcal{E}}(\Omega)$, $e_1(\varphi) = 0$ also implies $e_n(\varphi) = 0$.

Proof of Lemma 1

Without loss generality we can assume that Ω is the unit ball and $\varphi \in \mathcal{E}_0(\Omega)$. For $h, \psi \in \mathcal{E}_0(\Omega)$ an integration by parts and the Cauchy-Schwarz inequality yield

$$\begin{aligned}
 & \left[\int_{\Omega} -h(dd^c\varphi)^j \wedge (dd^c\psi)^{n-j} \right]^2 \\
 &= \left[\int_{\Omega} d\varphi \wedge d^c\psi \wedge (dd^c\varphi)^{j-1} \wedge (dd^c\psi)^{n-j-1} \wedge dd^c h \right]^2 \\
 &\leq \int_{\Omega} d\psi \wedge d^c\psi \wedge (dd^c\varphi)^{j-1} \wedge (dd^c\psi)^{n-j-1} \wedge dd^c h \\
 &\quad \int_{\Omega} d\varphi \wedge d^c\varphi \wedge (dd^c\varphi)^{j-1} \wedge (dd^c\psi)^{n-j-1} \wedge dd^c h \\
 &= \int_{\Omega} -h(dd^c\varphi)^{j-1} \wedge (dd^c\psi)^{n-j+1} \int_{\Omega} -h(dd^c\varphi)^{j+1} \wedge (dd^c\psi)^{n-j-1},
 \end{aligned}$$

Proof of Lemma 1, continued

Now, as $p \rightarrow +\infty$, take

$$h(z) = h_p(z) = \max \left(-1, \frac{1}{p} \log \|z\| \right) \nearrow \begin{cases} 0 & \text{if } z \in \Omega \setminus \{0\} \\ -1 & \text{if } z = 0. \end{cases}$$

By the monotone convergence theorem we get in the limit that

$$\begin{aligned}
 \left[\int_{\{0\}} (dd^c\varphi)^j \wedge (dd^c\psi)^{n-j} \right]^2 &\leq \int_{\{0\}} (dd^c\varphi)^{j-1} \wedge (dd^c\psi)^{n-j+1} \\
 &\quad \int_{\{0\}} (dd^c\varphi)^{j+1} \wedge (dd^c\psi)^{n-j-1}.
 \end{aligned}$$

For $\psi(z) = \ln \|z\|$, this is the desired estimate. □

Lemma 2

Let $\varphi, \psi \in \tilde{\mathcal{E}}(\Omega)$ be such that $\varphi \leq \psi$ (i.e φ is "more singular" than ψ). Then we have

$$\sum_{j=0}^{n-1} \frac{e_j(\varphi)}{e_{j+1}(\varphi)} \leq \sum_{j=0}^{n-1} \frac{e_j(\psi)}{e_{j+1}(\psi)}.$$

The argument is based on the monotonicity of Lelong numbers with respect to the relation $\varphi \leq \psi$, and on the monotonicity of the right hand side in the relevant range of values.

Proof of Lemma 2

Set

$$D = \{t = (t_1, \dots, t_n) \in [0, +\infty)^n : t_1^2 \leq t_2, t_j^2 \leq t_{j-1}t_{j+1}, \forall j = 2, \dots, n-1\}.$$

Then D is a convex set in \mathbb{R}^n , as can be checked by a straightforward application of the Cauchy-Schwarz inequality. Next, consider the function $f : \text{int } D \rightarrow [0, +\infty)$ defined by

$$f(t_1, \dots, t_n) = \frac{1}{t_1} + \frac{t_1}{t_2} \dots + \frac{t_{n-1}}{t_n}.$$

We have

$$\frac{\partial f}{\partial t_j}(t) = -\frac{t_{j-1}}{t_j^2} + \frac{1}{t_{j+1}} \leq 0, \quad \forall t \in D.$$

Proof of Lemma 2, continued

For $a, b \in \text{int } D$ such that $a_j \geq b_j, j = 1, \dots, n$, the function

$$[0, 1] \ni \lambda \rightarrow f(b + \lambda(a - b))$$

is decreasing. Hence,

$$f(a) \leq f(b) \quad \text{for all } a, b \in \text{int } D, a_j \geq b_j, j = 1, \dots, n.$$

On the other hand, the hypothesis $\varphi \leq \psi$ implies that $e_j(\varphi) \geq e_j(\psi), j = 1, \dots, n$, by the comparison principle. Therefore we have that

$$f(e_1(\varphi), \dots, e_n(\varphi)) \leq f(e_1(\psi), \dots, e_n(\psi)).$$

□

Proof of the Main Theorem

It will be convenient here to introduce Kiselman's refined Lelong number.

Definition

Let $\varphi \in \mathcal{PSH}(\Omega)$. Then the function defined by

$$\nu_\varphi(x) = \lim_{t \rightarrow -\infty} \frac{\max \{ \varphi(z) : |z_1| = e^{x_1 t}, \dots, |z_n| = e^{x_n t} \}}{t}$$

is called the **refined Lelong number** of φ at 0.

The refined Lelong number of φ at 0 is increasing in each variable x_j , and concave on \mathbb{R}^n_+ .

Proof of the Main Theorem

The proof is divided into the following steps:

- **Proof of the theorem in the toric case**, i.e. $\varphi(z_1, \dots, z_n) = \varphi(|z_1|, \dots, |z_n|)$ depends only on $|z_j|$ and therefore we can without loss of generality assume that $\Omega = \Delta^n$ is the unit polydisk.
- **Reduction to the case of plurisubharmonic functions with analytic singularity**, i.e. $\varphi = \log(|f_1|^2 + \dots + |f_N|^2)$, where f_1, \dots, f_N are germs of holomorphic functions at 0.
- **Reduction to the case of monomial ideals**, i.e. for $\varphi = \log(|f_1|^2 + \dots + |f_N|^2)$, where f_1, \dots, f_N are germs of monomial elements at 0.

Proof of the theorem in the toric case

Set

$$\Sigma = \left\{ x = (x_1, \dots, x_n) \in \mathbb{R}_+^n : \sum_{j=1}^n x_j = 1 \right\}.$$

We choose $x^0 = (x_1^0, \dots, x_n^0) \in \Sigma$ such that

$$\nu_\varphi(x^0) = \max\{\nu_\varphi(x) : x \in \Sigma\}.$$

By Theorem 5.8 in [Kis94] we have the following formula

$$c(\varphi) = \frac{1}{\nu_\varphi(x^0)}.$$

Proof of the theorem in the toric case, continued

Set $\zeta(x) = \nu_\varphi(x^0) \min \left(\frac{x_1}{x_1^0}, \dots, \frac{x_n}{x_n^0} \right)$, $\forall x \in \Sigma$.

Then ζ is the smallest nonnegative concave increasing function on Σ such that $\zeta(x^0) = \nu_\varphi(x^0)$, hence $\zeta \leq \nu_\varphi$. This implies that

$$\begin{aligned} \varphi(z_1, \dots, z_n) &\leq -\nu_\varphi(-\ln |z_1|, \dots, -\ln |z_n|) \\ &\leq -\zeta(-\ln |z_1|, \dots, -\ln |z_n|) \\ &\leq \nu_\varphi(x^0) \max \left(\frac{\ln |z_1|}{x_1^0}, \dots, \frac{\ln |z_n|}{x_n^0} \right) := \psi(z_1, \dots, z_n). \end{aligned}$$

By Lemma 2 we get that

$$f(\mathbf{e}_1(\varphi), \dots, \mathbf{e}_n(\varphi)) \leq f(\mathbf{e}_1(\psi), \dots, \mathbf{e}_n(\psi)) = \mathbf{c}(\psi) = \frac{1}{\nu_\varphi(x^0)} = \mathbf{c}(\varphi).$$

Reduction to the case of plurisubharmonic functions with analytic singularity

Let $\mathcal{H}_{m\varphi}(\Omega)$ be the Hilbert space of holomorphic functions f on Ω such that

$$\int_{\Omega} |f|^2 e^{-2m\varphi} dV < +\infty,$$

and let $\psi_m = \frac{1}{2m} \log \sum |g_{m,k}|^2$ where $\{g_{m,k}\}_{k \geq 1}$ be an orthonormal basis for $\mathcal{H}_{m\varphi}(\Omega)$. Using $\bar{\partial}$ -equation with L^2 -estimates (D-Kollár), there are constants $C_1, C_2 > 0$ independent of m such that

$$\varphi(z) - \frac{C_1}{m} \leq \psi_m(z) \leq \sup_{|\zeta-z|<r} \varphi(\zeta) + \frac{1}{m} \log \frac{C_2}{r^n}$$

for every $z \in \Omega$ and $r < d(z, \partial\Omega)$.

Reduction to the case of plurisubharmonic functions with analytic singularity, continued

and

$$\nu(\varphi) - \frac{n}{m} \leq \nu(\psi_m) \leq \nu(\varphi), \quad \frac{1}{c(\varphi)} - \frac{1}{m} \leq \frac{1}{c(\psi_m)} \leq \frac{1}{c(\varphi)}.$$

By Lemma 2, we have that

$$f(\mathbf{e}_1(\varphi), \dots, \mathbf{e}_n(\varphi)) \leq f(\mathbf{e}_1(\psi_m), \dots, \mathbf{e}_n(\psi_m)), \quad \forall m \geq 1.$$

The above inequalities show that in order to prove the lower bound of $c(\varphi)$ in the Main Theorem, we only need prove it for $c(\psi_m)$ and then let $m \rightarrow \infty$.

Reduction to the case of monomial ideals

For $j = 0, \dots, n$ set

$$\mathcal{J} = (f_1, \dots, f_N), \quad c(\mathcal{J}) = c(\varphi), \quad \text{and } e_j(\mathcal{J}) = e_j(\varphi).$$

Now, by fixing a multiplicative order on the monomials

$$z^\alpha = z_1^{\alpha_1} \dots z_n^{\alpha_n}$$

it is well known that one can construct a flat family $(\mathcal{J}_s)_{s \in \mathbb{C}}$ of ideals of $\mathcal{O}_{\mathbb{C}^n, 0}$ depending on a complex parameter $s \in \mathbb{C}$, such that \mathcal{J}_0 is a monomial ideal, $\mathcal{J}_1 = \mathcal{J}$ and

$$\dim(\mathcal{O}_{\mathbb{C}^n, 0} / \mathcal{J}_s^t) = \dim(\mathcal{O}_{\mathbb{C}^n, 0} / \mathcal{J}^t) \quad \text{for all } s, t \in \mathbb{N}.$$

In fact \mathcal{J}_0 is just the initial ideal associated to \mathcal{J} with respect to the monomial order.

Reduction to the case of monomial ideals, continued

Moreover, we can arrange by a generic rotation of coordinates $\mathbb{C}^p \subset \mathbb{C}^n$ that the family of ideals $\mathcal{J}_{s|\mathbb{C}^p}$ is also flat, and that the dimensions

$$\dim (\mathcal{O}_{\mathbb{C}^p,0}/(\mathcal{J}_{s|\mathbb{C}^p})^t) = \dim (\mathcal{O}_{\mathbb{C}^p,0}/(\mathcal{J}_{|\mathbb{C}^p})^t)$$

compute the intermediate multiplicities

$$e_p(\mathcal{J}_s) = \lim_{t \rightarrow +\infty} \frac{p!}{t^p} \dim (\mathcal{O}_{\mathbb{C}^p,0}/(\mathcal{J}_{s|\mathbb{C}^p})^t) = e_p(\mathcal{J}),$$

in particular, $e_p(\mathcal{J}_0) = e_p(\mathcal{J})$ for all p . The semicontinuity property of the log canonical threshold implies that $c(\mathcal{J}_0) \leq c(\mathcal{J}_s) = c(\mathcal{J})$ for all s , so the lower bound is valid for $c(\mathcal{J})$ if it is valid for $c(\mathcal{J}_0)$.

About the continuity of Monge-Ampère operators

Conjecture

Let $\varphi \in \tilde{\mathcal{E}}(\Omega)$ and $\Omega \ni 0$. Then the analytic approximations ψ_m satisfy $e_j(\psi_m) \rightarrow e_j(\varphi)$ as $m \rightarrow +\infty$, in other words, we have “strong continuity” of Monge-Ampère operators and higher Lelong numbers with respect to Bergman kernel approximation.

In the 2-dimensional case, $e_2(\varphi)$ can be computed as follows (at least when $\varphi \in \tilde{\mathcal{E}}(\omega)$ has analytic singularities).

Let $\mu : \tilde{\Omega} \rightarrow \Omega$ be the blow-up of Ω at 0. Take local coordinates (w_1, w_2) on $\tilde{\Omega}$ so that the exceptional divisor E can be written $w_1 = 0$.

About the continuity of Monge-Ampère operators (II)

With $\gamma = \nu(\varphi, 0)$, we get that

$$\tilde{\varphi}(w) = \varphi \circ \mu(w) - \gamma \log |w_1|$$

is psh with generic Lelong numbers equal to 0 along E , and therefore there are at most countably many points $x_\ell \in E$ at which $\gamma_\ell = \nu(\tilde{\varphi}, x_\ell) > 0$. Set $\Theta = dd^c \varphi$, $\tilde{\Theta} = dd^c \tilde{\varphi} = \mu^* \Theta - \gamma[E]$. Since $E^2 = -1$ in cohomology, we have $\{\tilde{\Theta}\}^2 = \{\mu^* \Theta\}^2 - \gamma^2$ in $H^2(E, \mathbb{R})$, hence

$$(*) \quad \int_{\{0\}} (dd^c \varphi)^2 = \gamma^2 + \int_E (dd^c \tilde{\varphi})^2.$$

If $\tilde{\varphi}$ only has ordinary logarithmic poles at the x_ℓ 's, then $\int_E (dd^c \tilde{\varphi})^2 = \sum \gamma_\ell^2$, but in general the situation is more complicated. Let us blow-up any of the points x_ℓ and repeat the process k times.

About the continuity of Monge-Ampère operators (III)

We set $\ell = \ell_1$ in what follows, as this was the first step, and at step $k = 0$ we omit any indices as 0 is the only point we have to blow-up to start with. We then get inductively $(k + 1)$ -iterated blow-ups depending on multi-indices $\ell = (\ell_1, \dots, \ell_k) = (\ell', \ell_k)$ with $\ell' = (\ell_1, \dots, \ell_{k-1})$,

$$\mu_\ell : \tilde{\Omega}_\ell \rightarrow \tilde{\Omega}_{\ell'}, \quad k \geq 1, \quad \mu_\emptyset = \mu : \tilde{\Omega}_\emptyset = \tilde{\Omega} \rightarrow \Omega, \quad \gamma_\emptyset = \gamma$$

and exceptional divisors $E_\ell \subset \tilde{\Omega}_\ell$ lying over points $x_\ell \in E_{\ell'} \subset \tilde{\Omega}_{\ell'}$, where

$$\gamma_\ell = \nu(\tilde{\varphi}_{\ell'}, x_\ell) > 0,$$

$$\tilde{\varphi}_\ell(w) = \tilde{\varphi}_{\ell'} \circ \mu_\ell(w) - \gamma_\ell \log |w_1^{(\ell)}|,$$

$$(w_1^{(\ell)})^2 = 0 \text{ an equation of } E_\ell \text{ in the relevant chart.}$$

Formula (*) implies

$$(**) \quad e_2(\varphi) \geq \sum_{k=0}^{+\infty} \sum_{\ell \in \mathbb{N}^k} \gamma_\ell^2$$

with equality when φ has an analytic singularity at 0. We conjecture that (**) is always an equality whenever $\varphi \in \tilde{\mathcal{E}}(\Omega)$.

This would imply the Guedj-Rashkovskii conjecture.

Notice that the currents $\Theta_\ell = dd^c \tilde{\varphi}_\ell$ satisfy inductively $\Theta_\ell = \mu_\ell^* \Theta_{\ell'} - \gamma_\ell [E_\ell]$, hence the cohomology class of Θ_ℓ restricted to E_ℓ is equal to γ_ℓ times the fundamental generator of E_ℓ . As a consequence we have

$$\sum_{\ell_{k+1} \in \mathbb{N}} \gamma_{\ell, \ell_{k+1}} \leq \gamma_\ell,$$

in particular $\gamma_\ell = 0$ for all $\ell \in \mathbb{N}^k$ if $\gamma = \nu(\varphi, 0) = 0$.

References I

[ÅCCP] Åhag P., Cegrell U., Czyż R. and Phạm H. H., *Monge-Ampère measures on pluripolar sets*, J. Math. Anal. Appl. 92 (2009), 613-627.

[ÅCKPZ] Åhag P., Cegrell U., Kołodziej S., Phạm H.H. and Zeriahi A., *Partial pluricomplex energy and integrability exponents of plurisubharmonic functions*, Adv. Math. 222 (2009), 2036-2058.

[AGZV85] Arnol'd V. I., Guseĭn-Zade S. M. and Varchenko A. N., *Singularities of differentiable maps. Vols. I–II*, Monographs in Mathematics, vol. 82, Birkhäuser Boston Inc., Boston, MA, 1985, Translated from the Russian by Ian Porteous and Mark Reynolds.

[Ati70] Atiyah M. F., *Resolution of singularities and division of distributions*, Comm. Pure Appl. Math. 23 (1970), 145-150.

References II

[Ber71] Bernšteĭn I. N., *Modules over a ring of differential operators. An investigation of the fundamental solutions of equations with constant coefficients*, Funkcional. Anal. i Priložen. 5 (1971), 1-16.

[Ceg04] Cegrell U., *The general definition of the complex Monge-Ampère operator*, Ann. Inst. Fourier (Grenoble) 54 (2004), 159-179.

[DK01] Demailly J.-P. and Kollár J., *Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds*, Ann. Sci. École Norm. Sup. (4) 34 (2001), 525-556.

[dFEM03] de Fernex T, Ein L. and Mustața, *Bounds for log canonical thresholds with applications to birational rigidity*, Math. Res. Lett. 10 (2003) 219-236.

References III

[dFEM04] de Fernex T., Ein L. and Mustața M., *Multiplicities and log canonical threshold*, J. Algebraic Geom. 13 (2004), 603-615.

[dFEM09] de Fernex T., Ein L. and Mustața M., *Shokurov's ACC Conjecture for log canonical thresholds on smooth varieties*, Duke Math. J., 152 (2010) 93-114.

[Eis95] Eisenbud D., *Commutative algebra with a view toward algebraic geometry*, Grad. Texts in Math. 150, Springer, New York, 1995.

[GŠ58] Gel'fand I. M. and Šilov G. E., *Obobshchennye funktsii i deistviya nad nimi*, Obobščennye funktsii, Vypusk 1. Generalized functions, part 1, Gosudarstv. Izdat. Fiz.-Mat. Lit., Moscow, 1958.

References IV

[Hör58] Hörmander L., *On the division of distributions by polynomials*, Ark. Mat. 3 (1958), 555-568.

[Kis94] Kiselman C.O., *Attenuating the singularities of plurisubharmonic functions*, Ann. Polon. Math. 60 (1994) 173-197.

[Łoj58] Łojasiewicz S., *Division d'une distribution par une fonction analytique de variables réelles*, C. R. Acad. Sci. Paris 246 (1958), 683-686.

[PS00] Phong D. H. and Sturm J., *On a conjecture of Demailly and Kollár. Kodaira's issue*, Asian J. Math. 4 (2000), 221-226.

[Sch50] Schwartz L., *Théorie des distributions. Tome I*, Actualités Sci. Ind., no. 1091 = Publ. Inst. Math. Univ. Strasbourg 9, Hermann & Cie., Paris, 1950.

References V

[Sko72] Skoda H., *Sous-ensembles analytiques d'ordre fini ou infini dans \mathbb{C}^n* , Bull. Soc. Math. France 100 (1972), 353-408.