



# Towards the Green-Griffiths-Lang conjecture

Jean-Pierre Demailly

Institut Fourier, Université de Grenoble I, France  
& Académie des Sciences de Paris

June 19, 2014

Complex Geometry & Lie Groups Conference  
Università degli Studi di Torino

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.  
 $X$  is said to be **(Brody) hyperbolic** if  $\nexists$  such  $f : \mathbb{C} \rightarrow X$ .

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.  
 $X$  is said to be **(Brody) hyperbolic** if  $\nexists$  such  $f : \mathbb{C} \rightarrow X$ .
- If  $X$  is a **bounded** open subset  $\Omega \subset \mathbb{C}^n$ , then there are no entire curves  $f : \mathbb{C} \rightarrow \Omega$  (**Liouville's theorem**),  
 $\Rightarrow$  every bounded open set  $\Omega \subset \mathbb{C}^n$  is hyperbolic

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.  
 $X$  is said to be **(Brody) hyperbolic** if  $\nexists$  such  $f : \mathbb{C} \rightarrow X$ .
- If  $X$  is a **bounded** open subset  $\Omega \subset \mathbb{C}^n$ , then there are no entire curves  $f : \mathbb{C} \rightarrow \Omega$  (**Liouville's theorem**),  
 $\Rightarrow$  every bounded open set  $\Omega \subset \mathbb{C}^n$  is hyperbolic
- $X = \overline{\mathbb{C}} \setminus \{0, 1, \infty\} = \mathbb{C} \setminus \{0, 1\}$  has no entire curves, so it is hyperbolic (**Picard's theorem**)

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.  
 $X$  is said to be **(Brody) hyperbolic** if  $\nexists$  such  $f : \mathbb{C} \rightarrow X$ .
- If  $X$  is a **bounded** open subset  $\Omega \subset \mathbb{C}^n$ , then there are no entire curves  $f : \mathbb{C} \rightarrow \Omega$  (**Liouville's theorem**),  
 $\Rightarrow$  every bounded open set  $\Omega \subset \mathbb{C}^n$  is hyperbolic
- $X = \overline{\mathbb{C}} \setminus \{0, 1, \infty\} = \mathbb{C} \setminus \{0, 1\}$  has no entire curves, so it is hyperbolic (**Picard's theorem**)
- A complex torus  $X = \mathbb{C}^n / \Lambda$  ( $\Lambda$  lattice) has a lot of entire curves. As  $\mathbb{C}$  simply connected, every  $f : \mathbb{C} \rightarrow X = \mathbb{C}^n / \Lambda$  lifts as  $\tilde{f} : \mathbb{C} \rightarrow \mathbb{C}^n$ ,  $\tilde{f}(t) = (\tilde{f}_1(t), \dots, \tilde{f}_n(t))$ , and  $\tilde{f}_j : \mathbb{C} \rightarrow \mathbb{C}$  can be arbitrary entire functions.

- Consider now the complex projective  $n$ -space

$$\mathbb{P}^n = \mathbb{P}_{\mathbb{C}}^n = (\mathbb{C}^{n+1} \setminus \{0\})/\mathbb{C}^*, \quad [z] = [z_0 : z_1 : \dots : z_n].$$

- Consider now the complex projective  $n$ -space

$$\mathbb{P}^n = \mathbb{P}_{\mathbb{C}}^n = (\mathbb{C}^{n+1} \setminus \{0\})/\mathbb{C}^*, \quad [z] = [z_0 : z_1 : \dots : z_n].$$

- An entire curve  $f : \mathbb{C} \rightarrow \mathbb{P}^n$  is given by a map

$$t \longmapsto [f_0(t) : f_1(t) : \dots : f_n(t)]$$

where  $f_j : \mathbb{C} \rightarrow \mathbb{C}$  are holomorphic functions without common zeroes (so there are a lot of them).



- Consider now the complex projective  $n$ -space

$$\mathbb{P}^n = \mathbb{P}_{\mathbb{C}}^n = (\mathbb{C}^{n+1} \setminus \{0\})/\mathbb{C}^*, \quad [z] = [z_0 : z_1 : \dots : z_n].$$

- An entire curve  $f : \mathbb{C} \rightarrow \mathbb{P}^n$  is given by a map


$$t \longmapsto [f_0(t) : f_1(t) : \dots : f_n(t)]$$

where  $f_j : \mathbb{C} \rightarrow \mathbb{C}$  are holomorphic functions without common zeroes (so there are a lot of them).


- More generally, look at a (complex) **projective manifold**, i.e.

$$X^n \subset \mathbb{P}^N, \quad X = \{[z]; P_1(z) = \dots = P_k(z) = 0\}$$

where  $P_j(z) = P_j(z_0, z_1, \dots, z_N)$  are homogeneous polynomials (of some degree  $d_j$ ), such that  $X$  is **non singular**.




$g = 0, K_X < 0$   
(positive curvature)




$g = 1, K_X = 0$   
(zero curvature)

Canonical bundle  $K_X = \Lambda^n T_X^*$  (here  $K_X = T_X^*$ )




$g = 0, K_X < 0$   
(positive curvature)




$g = 1, K_X = 0$   
(zero curvature)

Canonical bundle  $K_X = \Lambda^n T_X^*$  (here  $K_X = T_X^*$ )

- $g = 0 : X = \mathbb{P}^1$  curvature  $T_X > 0$  : not hyperbolic




$g = 0, K_X < 0$   
(positive curvature)



$g = 1, K_X = 0$   
(zero curvature)

Canonical bundle  $K_X = \Lambda^n T_X^*$  (here  $K_X = T_X^*$ )


- $g = 0 : X = \mathbb{P}^1$  curvature  $T_X > 0$  : not hyperbolic
- $g = 1 : X = \mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$  curvature  $T_X = 0$  : not hyperbolic



$g > 1, K_X > 0$   
(negative curvature)

$$\deg K_X = 2g - 2$$

If  $g \geq 2$ ,  $X \simeq \mathbb{D}/\Gamma$  ( $T_X < 0$ )  $\Rightarrow$   $X$  is hyperbolic.



$g > 1, K_X > 0$   
(negative curvature)

$$\deg K_X = 2g - 2$$

If  $g \geq 2$ ,  $X \simeq \mathbb{D}/\Gamma$  ( $T_X < 0$ )  $\Rightarrow$   $X$  is hyperbolic.

In fact every curve  $f : \mathbb{C} \rightarrow X \simeq \mathbb{D}/\Gamma$  lifts to  $\tilde{f} : \mathbb{C} \rightarrow \mathbb{D}$ ,  
and so  $\tilde{f}$  and  $f$  must be constant by Liouville.

**Definition** Let  $L \rightarrow X$  be a line bundle on a nonsingular complex projective variety  $X$ .

- $L$  is said to be **ample** if for  $m \gg 1$  the space of sections  $S_m = H^0(X, K_X^{\otimes m})$  gives an embedding

$$\Phi_m : X \hookrightarrow P(S_m^*) = \mathbb{P}^{N_m-1}, \quad N_m = \dim S_m.$$

**Definition** Let  $L \rightarrow X$  be a line bundle on a nonsingular complex projective variety  $X$ .

- $L$  is said to be **ample** if for  $m \gg 1$  the space of sections  $S_m = H^0(X, K_X^{\otimes m})$  gives an embedding

$$\Phi_m : X \hookrightarrow P(S_m^*) = \mathbb{P}^{N_m-1}, \quad N_m = \dim S_m.$$

- $L$  is said to be **big** if the dimensions of sections

$$\dim H^0(X, L^{\otimes m}) \sim cm^n$$

have maximal growth  $\kappa(L) = n = \dim X$ .



**Definition** Let  $L \rightarrow X$  be a line bundle on a nonsingular complex projective variety  $X$ .

- $L$  is said to be **ample** if for  $m \gg 1$  the space of sections  $S_m = H^0(X, K_X^{\otimes m})$  gives an embedding

$$\Phi_m : X \hookrightarrow P(S_m^*) = \mathbb{P}^{N_m-1}, \quad N_m = \dim S_m.$$

- $L$  is said to be **big** if the dimensions of sections

$$\dim H^0(X, L^{\otimes m}) \sim cm^n$$

have maximal growth  $\kappa(L) = n = \dim X$ .

- $X$  is said to be of **general type** if  $K_X$  is big.

**Definition** Let  $L \rightarrow X$  be a line bundle on a nonsingular complex projective variety  $X$ .

- $L$  is said to be **ample** if for  $m \gg 1$  the space of sections  $S_m = H^0(X, K_X^{\otimes m})$  gives an embedding

$$\Phi_m : X \hookrightarrow P(S_m^*) = \mathbb{P}^{N_m-1}, \quad N_m = \dim S_m.$$

- $L$  is said to be **big** if the dimensions of sections

$$\dim H^0(X, L^{\otimes m}) \sim cm^n$$

have maximal growth  $\kappa(L) = n = \dim X$ .

- $X$  is said to be of **general type** if  $K_X$  is big.

**Example:** A non singular hypersurface  $X^n \subset \mathbb{P}^{n+1}$  of degree  $d$  has  $K_X = \mathcal{O}(d - n - 2)$ ,  $X$  is of general type iff  $d > n + 2$ .

**Definition** Let  $L \rightarrow X$  be a line bundle on a nonsingular complex projective variety  $X$ .

- $L$  is said to be **ample** if for  $m \gg 1$  the space of sections  $S_m = H^0(X, K_X^{\otimes m})$  gives an embedding

$$\Phi_m : X \hookrightarrow P(S_m^*) = \mathbb{P}^{N_m-1}, \quad N_m = \dim S_m.$$

- $L$  is said to be **big** if the dimensions of sections

$$\dim H^0(X, L^{\otimes m}) \sim cm^n$$

have maximal growth  $\kappa(L) = n = \dim X$ .

- $X$  is said to be of **general type** if  $K_X$  is big.

**Example:** A non singular hypersurface  $X^n \subset \mathbb{P}^{n+1}$  of degree  $d$  has  $K_X = \mathcal{O}(d - n - 2)$ ,  $X$  is of general type iff  $d > n + 2$ .

**Conjecture CGT.** If a compact variety  $X$  is hyperbolic, then it should be of general type, and if  $X$  is non singular, then  $K_X = \Lambda^n T_X^*$  should be ample, i.e.  $K_X > 0$  (Kodaira).

- **Conjecture (Green-Griffiths-Lang = GGL)**

*Let  $X$  be a projective variety of general type. Then*

*$\exists Y \subsetneq X$  algebraic such that all entire curves  $f : \mathbb{C} \rightarrow X$  satisfy  $f(\mathbb{C}) \subset Y$ .*

- **Conjecture (Green-Griffiths-Lang = GGL)**

*Let  $X$  be a projective variety of general type. Then*

*$\exists Y \subsetneq X$  algebraic such that all entire curves  $f : \mathbb{C} \rightarrow X$  satisfy  $f(\mathbb{C}) \subset Y$ .*

- **Expected consequence of GGL:**

*If every subvariety  $Y$  of  $X$  is of **general type**, then  $X$  is hyperbolic.*

- **Conjecture (Green-Griffiths-Lang = GGL)**

*Let  $X$  be a projective variety of general type. Then*

*$\exists Y \subsetneq X$  algebraic such that all entire curves  $f : \mathbb{C} \rightarrow X$  satisfy  $f(\mathbb{C}) \subset Y$ .*

- **Expected consequence of GGL:**

*If every subvariety  $Y$  of  $X$  is of **general type**, then  $X$  is hyperbolic.*

By CGT conjecture, this should be a necessary and sufficient characterization of hyperbolicity for projective varieties.

- **Conjecture (Green-Griffiths-Lang = GGL)**

*Let  $X$  be a projective variety of general type. Then*

*$\exists Y \subsetneq X$  algebraic such that all entire curves  $f : \mathbb{C} \rightarrow X$  satisfy  $f(\mathbb{C}) \subset Y$ .*

- **Expected consequence of GGL:**

*If every subvariety  $Y$  of  $X$  is of **general type**, then  $X$  is hyperbolic.*

By CGT conjecture, this should be a necessary and sufficient characterization of hyperbolicity for projective varieties.

- **Arithmetic counterpart (Lang 1987).** *If  $X$  is a variety of general type defined over a number field and  $Y$  is the Green-Griffiths locus (Zariski closure of  $\bigcup f(\mathbb{C})$ ), then  $X(\mathbb{K}) \setminus Y$  is finite for every number field  $\mathbb{K}$ .*

- Using “jet technology” and **deep results of McQuillan** for curve foliations on surfaces, D. – El Goul proved **Theorem** (solution of Kobayashi conjecture, 1998).  
*A very generic surface  $X \subset \mathbb{P}^3$  of **degree**  $\geq 21$  is hyperbolic.*  
Independently McQuillan got  $\text{degree} \geq 35$ .  
Recently improved to **degree**  $\geq 18$  (Păun, 2008).



- Using “jet technology” and **deep results of McQuillan** for curve foliations on surfaces, D. – El Goul proved **Theorem** (solution of Kobayashi conjecture, 1998).  
*A very generic surface  $X \subset \mathbb{P}^3$  of **degree  $\geq 21$**  is hyperbolic.*  
Independently McQuillan got  $\text{degree} \geq 35$ .  
Recently improved to **degree  $\geq 18$**  (Păun, 2008).  
Still far : for  $X \subset \mathbb{P}^{n+1}$ , the optimal bound should be **degree  $\geq 2n + 1$  for  $n \geq 2$  (Zaidenberg).**

- Using “jet technology” and **deep results of McQuillan** for curve foliations on surfaces, D. – El Goul proved **Theorem** (solution of Kobayashi conjecture, 1998).  
*A very generic surface  $X \subset \mathbb{P}^3$  of **degree  $\geq 21$**  is hyperbolic.*  
Independently McQuillan got  $\text{degree} \geq 35$ .  
Recently improved to **degree  $\geq 18$**  (Păun, 2008).  
Still far : for  $X \subset \mathbb{P}^{n+1}$ , the optimal bound should be **degree  $\geq 2n + 1$  for  $n \geq 2$  (Zaidenberg)**.
- **Generic GGL conjecture** (S. Diverio, J. Merker, E. Rousseau, 2009). *A generic hypersurface  $X \subset \mathbb{P}^{n+1}$  of degree  $d \geq d_n := 2^{n^5}$  satisfies GGL (no entire curves).*

- Using “jet technology” and **deep results of McQuillan** for curve foliations on surfaces, D. – El Goul proved **Theorem** (solution of Kobayashi conjecture, 1998).  
*A very generic surface  $X \subset \mathbb{P}^3$  of **degree  $\geq 21$**  is hyperbolic.*  
 Independently McQuillan got  $\text{degree} \geq 35$ .  
 Recently improved to **degree  $\geq 18$**  (Păun, 2008).  
 Still far : for  $X \subset \mathbb{P}^{n+1}$ , the optimal bound should be **degree  $\geq 2n + 1$  for  $n \geq 2$  (Zaidenberg)**.
- **Generic GGL conjecture** (S. Diverio, J. Merker, E. Rousseau, 2009). *A generic hypersurface  $X \subset \mathbb{P}^{n+1}$  of degree  $d \geq d_n := 2^{n^5}$  satisfies GGL (no entire curves).*  
 (D-, 2012) *bound improved to  $d_n = \left\lfloor \frac{n^4}{3} (n \log(n \log(24n)))^n \right\rfloor$ .*

- Using “jet technology” and **deep results of McQuillan** for curve foliations on surfaces, D. – El Goul proved **Theorem** (solution of Kobayashi conjecture, 1998).  
*A very generic surface  $X \subset \mathbb{P}^3$  of **degree**  $\geq 21$  is hyperbolic.*  
 Independently McQuillan got  $\text{degree} \geq 35$ .  
 Recently improved to **degree**  $\geq 18$  (Păun, 2008).  
 Still far : for  $X \subset \mathbb{P}^{n+1}$ , the optimal bound should be **degree**  $\geq 2n + 1$  for  $n \geq 2$  (**Zaidenberg**).
- **Generic GGL conjecture** (S. Diverio, J. Merker, E. Rousseau, 2009). *A generic hypersurface  $X \subset \mathbb{P}^{n+1}$  of degree  $d \geq d_n := 2^{n^5}$  satisfies GGL (no entire curves).*  
 (D-, 2012) *bound improved to  $d_n = \left\lfloor \frac{n^4}{3} (n \log(n \log(24n)))^n \right\rfloor$ .*  
**Moreover** (S. Diverio, S. Trapani, 2009) A generic hypersurface  $X \subset \mathbb{P}^4$  of degree  $\geq 593$  is **hyperbolic**.

- **Goal.** More generally, we are interested in curves  $f : \mathbb{C} \rightarrow X$  such that  $f'(\mathbb{C}) \subset V$  where  $V$  is a subbundle of  $T_X$  (or singular linear subspace, i.e. a closed irreducible analytic subspace such that  $\forall x \in X, V_x := V \cap T_{X,x}$  linear).

- **Goal.** More generally, we are interested in curves  $f : \mathbb{C} \rightarrow X$  such that  $f'(\mathbb{C}) \subset V$  where  $V$  is a subbundle of  $T_X$  (or singular linear subspace, i.e. a closed irreducible analytic subspace such that  $\forall x \in X, V_x := V \cap T_{X,x}$  linear).
- **Definition.** *Category of directed manifolds :*
  - **Objects** : pairs  $(X, V)$ ,  $X$  manifold/ $\mathbb{C}$  and  $V \subset T_X$
  - **Arrows**  $\psi : (X, V) \rightarrow (Y, W)$  holomorphic s.t.  $\psi_* V \subset W$

- **Goal.** More generally, we are interested in curves  $f : \mathbb{C} \rightarrow X$  such that  $f'(\mathbb{C}) \subset V$  where  $V$  is a subbundle of  $T_X$  (or singular linear subspace, i.e. a closed irreducible analytic subspace such that  $\forall x \in X, V_x := V \cap T_{X,x}$  linear).
- **Definition.** *Category of directed manifolds :*
  - **Objects** : pairs  $(X, V)$ ,  $X$  manifold/ $\mathbb{C}$  and  $V \subset T_X$
  - **Arrows**  $\psi : (X, V) \rightarrow (Y, W)$  holomorphic s.t.  $\psi_* V \subset W$
  - “**Absolute case**”  $(X, T_X)$ , i.e.  $V = T_X$
  - “**Relative case**”  $(X, T_{X/S})$  where  $X \rightarrow S$
  - “**Integrable case**” when  $[V, V] \subset V$  (foliations)

- Goal.** More generally, we are interested in curves  $f : \mathbb{C} \rightarrow X$  such that  $f'(\mathbb{C}) \subset V$  where  $V$  is a subbundle of  $T_X$  (or singular linear subspace, i.e. a closed irreducible analytic subspace such that  $\forall x \in X, V_x := V \cap T_{X,x}$  linear).
- Definition.** *Category of directed manifolds :*
  - **Objects** : pairs  $(X, V)$ ,  $X$  manifold/ $\mathbb{C}$  and  $V \subset T_X$
  - **Arrows**  $\psi : (X, V) \rightarrow (Y, W)$  holomorphic s.t.  $\psi_* V \subset W$
  - “**Absolute case**”  $(X, T_X)$ , i.e.  $V = T_X$
  - “**Relative case**”  $(X, T_{X/S})$  where  $X \rightarrow S$
  - “**Integrable case**” when  $[V, V] \subset V$  (foliations)
- Fonctor “1-jet”** :  $(X, V) \mapsto (\tilde{X}, \tilde{V})$  where :
  - $\tilde{X} = P(V)$  = bundle of projective spaces of lines in  $V$
  - $\pi : \tilde{X} = P(V) \rightarrow X, \quad (x, [v]) \mapsto x, \quad v \in V_x$
  - $\tilde{V}_{(x,[v])} = \{ \xi \in T_{\tilde{X},(x,[v])} ; \pi_* \xi \in \mathbb{C}v \subset T_{X,x} \}$



- For every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  tangent to  $V$

$$f_{[1]}(t) := (f(t), [f'(t)]) \in P(V_{f(t)}) \subset \tilde{X}$$

$$f_{[1]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (\tilde{X}, \tilde{V}) \quad (\text{projectivized 1st-jet})$$

- For every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  tangent to  $V$

$$f_{[1]}(t) := (f(t), [f'(t)]) \in P(V_{f(t)}) \subset \tilde{X}$$

$$f_{[1]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (\tilde{X}, \tilde{V}) \quad (\text{projectivized 1st-jet})$$

- **Definition.** *Simple jet bundles :*

- $(X_k, V_k) = k$ -th iteration of functor  $(X, V) \mapsto (\tilde{X}, \tilde{V})$
- $f_{[k]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X_k, V_k)$  is the **projectivized  $k$ -jet of  $f$** .

- For every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  tangent to  $V$

$$f_{[1]}(t) := (f(t), [f'(t)]) \in P(V_{f(t)}) \subset \tilde{X}$$

$$f_{[1]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (\tilde{X}, \tilde{V}) \quad (\text{projectivized 1st-jet})$$

- **Definition.** *Simple jet bundles :*

–  $(X_k, V_k) = k$ -th iteration of functor  $(X, V) \mapsto (\tilde{X}, \tilde{V})$

–  $f_{[k]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X_k, V_k)$  is the **projectivized  $k$ -jet of  $f$** .

- **Basic exact sequences**

$$0 \rightarrow T_{\tilde{X}/X} \rightarrow \tilde{V} \xrightarrow{\pi^*} \mathcal{O}_{\tilde{X}}(-1) \rightarrow 0 \quad \Rightarrow \text{rk } \tilde{V} = r = \text{rk } V$$

$$0 \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow \pi^* V \otimes \mathcal{O}_{\tilde{X}}(1) \rightarrow T_{\tilde{X}/X} \rightarrow 0 \quad (\text{Euler})$$

- For every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  tangent to  $V$

$$f_{[1]}(t) := (f(t), [f'(t)]) \in P(V_{f(t)}) \subset \tilde{X}$$

$$f_{[1]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (\tilde{X}, \tilde{V}) \quad (\text{projectivized 1st-jet})$$

- **Definition.** *Simple jet bundles :*

–  $(X_k, V_k) = k$ -th iteration of functor  $(X, V) \mapsto (\tilde{X}, \tilde{V})$

–  $f_{[k]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X_k, V_k)$  is the **projectivized  $k$ -jet of  $f$** .

- **Basic exact sequences**

$$0 \rightarrow T_{\tilde{X}/X} \rightarrow \tilde{V} \xrightarrow{\pi^*} \mathcal{O}_{\tilde{X}}(-1) \rightarrow 0 \quad \Rightarrow \text{rk } \tilde{V} = r = \text{rk } V$$

$$0 \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow \pi^* V \otimes \mathcal{O}_{\tilde{X}}(1) \rightarrow T_{\tilde{X}/X} \rightarrow 0 \quad (\text{Euler})$$

$$0 \rightarrow T_{X_k/X_{k-1}} \rightarrow V_k \xrightarrow{(\pi_k)^*} \mathcal{O}_{X_k}(-1) \rightarrow 0 \quad \Rightarrow \text{rk } V_k = r$$

$$0 \rightarrow \mathcal{O}_{X_k} \rightarrow \pi_k^* V_{k-1} \otimes \mathcal{O}_{X_k}(1) \rightarrow T_{X_k/X_{k-1}} \rightarrow 0 \quad (\text{Euler})$$

- For  $n = \dim X$  and  $r = \operatorname{rk} V$ , get a **tower of  $\mathbb{P}^{r-1}$ -bundles**

$$\pi_{k,0} : X_k \xrightarrow{\pi_k} X_{k-1} \rightarrow \cdots \rightarrow X_1 \xrightarrow{\pi_1} X_0 = X$$

with  $\dim X_k = n + k(r - 1)$ ,  $\operatorname{rk} V_k = r$ ,  
and **tautological line bundles  $\mathcal{O}_{X_k}(1)$  on  $X_k = P(V_{k-1})$ .**

- For  $n = \dim X$  and  $r = \operatorname{rk} V$ , get a **tower of  $\mathbb{P}^{r-1}$ -bundles**

$$\pi_{k,0} : X_k \xrightarrow{\pi_k} X_{k-1} \rightarrow \cdots \rightarrow X_1 \xrightarrow{\pi_1} X_0 = X$$

with  $\dim X_k = n + k(r - 1)$ ,  $\operatorname{rk} V_k = r$ ,  
and **tautological line bundles  $\mathcal{O}_{X_k}(1)$**  on  $X_k = P(V_{k-1})$ .

- **Theorem.**  $X_k$  is a smooth compactification of

$$X_k^{\operatorname{GG}, \operatorname{reg}} / \mathbb{G}_k = J_k^{\operatorname{GG}, \operatorname{reg}} / \mathbb{G}_k$$

where  $\mathbb{G}_k$  is the group of  $k$ -jets of germs of biholomorphisms of  $(\mathbb{C}, 0)$ , acting on the right by reparametrization:  $(f, \varphi) \mapsto f \circ \varphi$ , and  $J_k^{\operatorname{reg}}$  is the space of  $k$ -jets of regular curves.

- For  $n = \dim X$  and  $r = \text{rk } V$ , get a **tower of  $\mathbb{P}^{r-1}$ -bundles**

$$\pi_{k,0} : X_k \xrightarrow{\pi_k} X_{k-1} \rightarrow \cdots \rightarrow X_1 \xrightarrow{\pi_1} X_0 = X$$

with  $\dim X_k = n + k(r - 1)$ ,  $\text{rk } V_k = r$ ,  
and **tautological line bundles  $\mathcal{O}_{X_k}(1)$**  on  $X_k = P(V_{k-1})$ .

- **Theorem.**  $X_k$  is a smooth compactification of

$$X_k^{\text{GG,reg}} / \mathbb{G}_k = J_k^{\text{GG,reg}} / \mathbb{G}_k$$

where  $\mathbb{G}_k$  is the group of  $k$ -jets of germs of biholomorphisms of  $(\mathbb{C}, 0)$ , acting on the right by reparametrization:  $(f, \varphi) \mapsto f \circ \varphi$ , and  $J_k^{\text{reg}}$  is the space of  $k$ -jets of regular curves.

- **Direct image formula.**  $(\pi_{k,0})_* \mathcal{O}_{X_k}(m) = E_{k,m} V^* =$   
invariant algebraic differential operators  $f \mapsto P(f_{[k]})$   
acting on germs of curves  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ .

Let  $(\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ ,  $t \mapsto f(t) = (f_1(t), \dots, f_n(t))$  be a curve written in some local holomorphic coordinates  $(z_1, \dots, z_n)$  on  $X$ . It has a local Taylor expansion

$$f(t) = x + t\xi_1 + \dots + t^k\xi_k + O(t^{k+1}), \quad \xi_s = \frac{1}{s!} \nabla^s f(0)$$

for some connection  $\nabla$  on  $V$ .



Let  $(\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ ,  $t \mapsto f(t) = (f_1(t), \dots, f_n(t))$  be a curve written in some local holomorphic coordinates  $(z_1, \dots, z_n)$  on  $X$ . It has a local Taylor expansion

$$f(t) = x + t\xi_1 + \dots + t^k\xi_k + O(t^{k+1}), \quad \xi_s = \frac{1}{s!} \nabla^s f(0)$$

for some connection  $\nabla$  on  $V$ .

One considers the **Green-Griffiths bundle**  $E_{k,m}^{\text{GG}} V^*$  of polynomials of weighted degree  $m$  written locally in coordinate charts as

$$P(x; \xi_1, \dots, \xi_k) = \sum a_{\alpha_1 \alpha_2 \dots \alpha_k}(x) \xi_1^{\alpha_1} \dots \xi_k^{\alpha_k}, \quad \xi_s \in V,$$

also viewed as **algebraic differential operators**

$$\begin{aligned} P(f_{[k]}) &= P(f', f'', \dots, f^{(k)}) \\ &= \sum a_{\alpha_1 \alpha_2 \dots \alpha_k}(f(t)) f'(t)^{\alpha_1} f''(t)^{\alpha_2} \dots f^{(k)}(t)^{\alpha_k}. \end{aligned}$$

# Definition of algebraic differential operators (2) 42/69

Here  $t \mapsto z = f(t)$  is a curve,  $f_{[k]} = (f', f'', \dots, f^{(k)})$  its  $k$ -jet, and  $a_{\alpha_1 \alpha_2 \dots \alpha_k}(z)$  are supposed to be holomorphic functions on  $X$ .

Here  $t \mapsto z = f(t)$  is a curve,  $f_{[k]} = (f', f'', \dots, f^{(k)})$  its  $k$ -jet, and  $a_{\alpha_1 \alpha_2 \dots \alpha_k}(z)$  are supposed to be holomorphic functions on  $X$ .

The  $\mathbb{G}_k$ -action:  $(f, \varphi) \mapsto f \circ \varphi$ , yields in particular,  $\varphi_\lambda(t) = \lambda t \Rightarrow (f \circ \varphi_\lambda)^{(k)}(t) = \lambda^k f^{(k)}(\lambda t)$ , whence a  $\mathbb{C}^*$ -action

$$\lambda \cdot (\xi_1, \xi_1, \dots, \xi_k) = (\lambda \xi_1, \lambda^2 \xi_2, \dots, \lambda^k \xi_k).$$

Here  $t \mapsto z = f(t)$  is a curve,  $f_{[k]} = (f', f'', \dots, f^{(k)})$  its  $k$ -jet, and  $a_{\alpha_1 \alpha_2 \dots \alpha_k}(z)$  are supposed to be holomorphic functions on  $X$ .

The  $\mathbb{G}_k$ -action:  $(f, \varphi) \mapsto f \circ \varphi$ , yields in particular,  $\varphi_\lambda(t) = \lambda t \Rightarrow (f \circ \varphi_\lambda)^{(k)}(t) = \lambda^k f^{(k)}(\lambda t)$ , whence a  $\mathbb{C}^*$ -action

$$\lambda \cdot (\xi_1, \xi_2, \dots, \xi_k) = (\lambda \xi_1, \lambda^2 \xi_2, \dots, \lambda^k \xi_k).$$

$E_{k,m}^{\text{GG}}$  is precisely the set of polynomials of weighted degree  $m$ , corresponding to coefficients  $a_{\alpha_1 \dots \alpha_k}$  with  $m = |\alpha_1| + 2|\alpha_2| + \dots + k|\alpha_k|$ .

Here  $t \mapsto z = f(t)$  is a curve,  $f_{[k]} = (f', f'', \dots, f^{(k)})$  its  $k$ -jet, and  $a_{\alpha_1 \alpha_2 \dots \alpha_k}(z)$  are supposed to holomorphic functions on  $X$ .

The  $\mathbb{G}_k$ -action :  $(f, \varphi) \mapsto f \circ \varphi$ , yields in particular,  $\varphi_\lambda(t) = \lambda t \Rightarrow (f \circ \varphi_\lambda)^{(k)}(t) = \lambda^k f^{(k)}(\lambda t)$ , whence a  $\mathbb{C}^*$ -action

$$\lambda \cdot (\xi_1, \xi_2, \dots, \xi_k) = (\lambda \xi_1, \lambda^2 \xi_2, \dots, \lambda^k \xi_k).$$

$E_{k,m}^{\text{GG}}$  is precisely the set of polynomials of weighted degree  $m$ , corresponding to coefficients  $a_{\alpha_1 \dots \alpha_k}$  with  $m = |\alpha_1| + 2|\alpha_2| + \dots + k|\alpha_k|$ .

$E_{k,m} V^* \subset E_{k,m}^{\text{GG}} V^*$  is the bundle of  $\mathbb{G}_k$ -"invariant" operators, i.e. such that

$$P((f \circ \varphi)_{[k]}) = \varphi'^m P(f_{[k]}) \circ \varphi, \quad \forall \varphi \in \mathbb{G}_k.$$

When  $V$  is nonsingular, we simply set  $K_V = \det(V^*)$ .  
When  $V$  is singular, the canonical sheaf  $K_V$  is the rank 1 analytic sheaf defined as the integral closure of the image of the natural morphism

$$\Lambda^r T_X^* \rightarrow \Lambda^r V^* \rightarrow \mathcal{L}_V := \text{invert. sheaf } (\Lambda^r V^*)^{**}$$

that is, if the image is  $\mathcal{L}_V \otimes \mathcal{I}_V$ ,  $\mathcal{I}_V \subset \mathcal{O}_X$ , one sets

$$K_V = \mathcal{L}_V \otimes \overline{\mathcal{I}_V}.$$

When  $V$  is nonsingular, we simply set  $K_V = \det(V^*)$ .  
 When  $V$  is singular, the canonical sheaf  $K_V$  is the rank 1 analytic sheaf defined as the integral closure of the image of the natural morphism

$$\Lambda^r T_X^* \rightarrow \Lambda^r V^* \rightarrow \mathcal{L}_V := \text{invert. sheaf } (\Lambda^r V^*)^{**}$$

that is, if the image is  $\mathcal{L}_V \otimes \mathcal{I}_V$ ,  $\mathcal{I}_V \subset \mathcal{O}_X$ , one sets

$$K_V = \mathcal{L}_V \otimes \overline{\mathcal{I}_V}.$$

**Definition.** We say that  $(X, V)$  is of general type if there exist proper modifications  $\mu = \hat{\mu} \circ \tilde{\mu} : \hat{X} \rightarrow \tilde{X} \rightarrow X$  such that  $\hat{\mu}^* K_{\tilde{V}}$  is a **big invertible sheaf** on  $\hat{X}$ , where  $\tilde{X}$  is equipped with the pull-back directed structure  $\tilde{V} = \overline{\tilde{\mu}^{-1}(V)}$ .

- **Generalized GGL conjecture.** *If  $(X, V)$  is directed manifold of general type, i.e.  $K_V$  big, then  $\exists Y \subsetneq X$  such that  $\forall f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  non const., one has  $f(\mathbb{C}) \subset Y$ .*



- **Generalized GGL conjecture.** *If  $(X, V)$  is directed manifold of general type, i.e.  $K_V$  big, then  $\exists Y \subsetneq X$  such that  $\forall f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  non const., one has  $f(\mathbb{C}) \subset Y$ .*
- **Remark.** Elementary by Ahlfors-Schwarz if  $r = \text{rk } V = 1$ .  $t \mapsto \log \|f'(t)\|_{V,h}$  is strictly subharmonic if  $r = 1$  and  $(V^*, h^*)$  has  $> 0$  curvature in the sense of currents.

- Generalized GGL conjecture.** *If  $(X, V)$  is directed manifold of general type, i.e.  $K_V$  big, then  $\exists Y \subsetneq X$  such that  $\forall f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  non const., one has  $f(\mathbb{C}) \subset Y$ .*
- Remark.** Elementary by Ahlfors-Schwarz if  $r = \text{rk } V = 1$ .  $t \mapsto \log \|f'(t)\|_{V,h}$  is strictly subharmonic if  $r = 1$  and  $(V^*, h^*)$  has  $> 0$  curvature in the sense of currents.
- Strategy : fundamental vanishing theorem.**  
 [Green-Griffiths 1979], [Demailly 1995], [Siu-Yeung 1996]  
*Let  $P \in H^0(X, E_{k,m}^{\text{GG}} V^* \otimes \mathcal{O}(-A))$  be a global algebraic differential operator whose coefficients vanish on some ample divisor  $A$ . Then  $\forall f : \mathbb{C} \rightarrow (X, V)$ , one has  $P(f_{[k]}) \equiv 0$ .*

- Generalized GGL conjecture.** *If  $(X, V)$  is directed manifold of general type, i.e.  $K_V$  big, then  $\exists Y \subsetneq X$  such that  $\forall f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  non const., one has  $f(\mathbb{C}) \subset Y$ .*
- Remark.** Elementary by Ahlfors-Schwarz if  $r = \text{rk } V = 1$ .  $t \mapsto \log \|f'(t)\|_{V,h}$  is strictly subharmonic if  $r = 1$  and  $(V^*, h^*)$  has  $> 0$  curvature in the sense of currents.
- Strategy : fundamental vanishing theorem.**  
 [Green-Griffiths 1979], [Demailly 1995], [Siu-Yeung 1996]  
*Let  $P \in H^0(X, E_{k,m}^{\text{GG}} V^* \otimes \mathcal{O}(-A))$  be a global algebraic differential operator whose coefficients vanish on some ample divisor  $A$ . Then  $\forall f : \mathbb{C} \rightarrow (X, V)$ , one has  $P(f_{[k]}) \equiv 0$ .*
- Theorem (D-, 2010).** Let  $(X, V)$  be of general type, i.e. s.t.  $K_V$  is a **big** rank 1 sheaf. Then  $\exists k \geq 1$  and  $\exists$  algebraic hypersurface  $\Sigma \subsetneq X_k$  such that every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  satisfies  $f_{[k]}(\mathbb{C}) \subset \Sigma$ .

Let  $J_k V$  be the bundle of  $k$ -jets of curves  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$

Let  $J_k V$  be the bundle of  $k$ -jets of curves  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$   
Assuming that  $V$  is equipped with a hermitian metric  $h$ , one defines a "weighted Finsler metric" on  $J^k V$  by taking  $p = k!$  and

$$\Psi_{h_k}(f) := \left( \sum_{1 \leq s \leq k} \varepsilon_s \|\nabla^s f(0)\|_{h(x)}^{2p/s} \right)^{1/p}, \quad 1 = \varepsilon_1 \gg \varepsilon_2 \gg \cdots \gg \varepsilon_k.$$

Let  $J_k V$  be the bundle of  $k$ -jets of curves  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ . Assuming that  $V$  is equipped with a hermitian metric  $h$ , one defines a "weighted Finsler metric" on  $J^k V$  by taking  $p = k!$  and

$$\Psi_{h_k}(f) := \left( \sum_{1 \leq s \leq k} \varepsilon_s \|\nabla^s f(0)\|_{h(x)}^{2p/s} \right)^{1/p}, \quad 1 = \varepsilon_1 \gg \varepsilon_2 \gg \dots \gg \varepsilon_k.$$

Letting  $\xi_s = \nabla^s f(0)$ , this can actually be viewed as a metric  $h_k$  on  $L_k := \mathcal{O}_{X_k^{\text{GG}}}(1)$ , with curvature form  $(x, \xi_1, \dots, \xi_k) \mapsto$

$$\Theta_{L_k, h_k} = \omega_{\text{FS}, k}(\xi) + \frac{i}{2\pi} \sum_{1 \leq s \leq k} \frac{1}{s} \frac{|\xi_s|^{2p/s}}{\sum_t |\xi_t|^{2p/t}} \sum_{i, j, \alpha, \beta} c_{ij\alpha\beta} \frac{\xi_{s\alpha} \bar{\xi}_{s\beta}}{|\xi_s|^2} dz_i \wedge d\bar{z}_j$$

where  $(c_{ij\alpha\beta})$  are the coefficients of the curvature tensor  $\Theta_{V^*, h^*}$  and  $\omega_{\text{FS}, k}$  is the vertical Fubini-Study metric on the fibers of  $X_k^{\text{GG}} \rightarrow X$ . The expression gets simpler by using polar coordinates  $x_s = |\xi_s|_h^{2p/s}$ ,  $u_s = \xi_s / |\xi_s|_h = \nabla^s f(0) / |\nabla^s f(0)|$ .

In such polar coordinates, one gets the formula

$$\Theta_{L_k, h_k} = \omega_{\text{FS}, p, k}(\xi) + \frac{i}{2\pi} \sum_{1 \leq s \leq k} \frac{1}{s} x_s \sum_{i, j, \alpha, \beta} c_{ij\alpha\beta}(z) u_{s\alpha} \bar{u}_{s\beta} dz_i \wedge d\bar{z}_j$$

where  $\omega_{\text{FS}, k}(\xi)$  is positive definite in  $\xi$ . The other terms are a weighted average of the values of the curvature tensor  $\Theta_{V, h}$  on vectors  $u_s$  in the unit sphere bundle  $SV \subset V$ .

In such polar coordinates, one gets the formula

$$\Theta_{L_k, h_k} = \omega_{\text{FS}, p, k}(\xi) + \frac{i}{2\pi} \sum_{1 \leq s \leq k} \frac{1}{s} x_s \sum_{i, j, \alpha, \beta} c_{ij\alpha\beta}(z) u_{s\alpha} \bar{u}_{s\beta} dz_i \wedge d\bar{z}_j$$

where  $\omega_{\text{FS}, k}(\xi)$  is positive definite in  $\xi$ . The other terms are a weighted average of the values of the curvature tensor  $\Theta_{V, h}$  on vectors  $u_s$  in the unit sphere bundle  $SV \subset V$ .

The weighted projective space can be viewed as a circle quotient of the pseudosphere  $\sum |\xi_s|^{2p/s} = 1$ , so we can take here  $x_s \geq 0$ ,  $\sum x_s = 1$ . This is essentially a sum of the form  $\sum \frac{1}{s} \gamma(u_s)$  where  $u_s$  are random points of the sphere, and so as  $k \rightarrow +\infty$  this can be estimated by a **“Monte-Carlo” integral**

$$\left(1 + \frac{1}{2} + \dots + \frac{1}{k}\right) \int_{u \in SV} \gamma(u) du.$$

As  $\gamma$  is quadratic here,  $\int_{u \in SV} \gamma(u) du = \frac{1}{r} \text{Tr}(\gamma)$ .



It follows that the leading term in the estimate only involves the trace of  $\Theta_{V^*, h^*}$ , i.e. the curvature of  $(\det V^*, \det h^*)$ , which can be taken to be  $> 0$  if  $\det V^*$  is big.

**Corollary (D-, 2010).** Let  $(X, V)$  be a directed manifold,  $F \rightarrow X$  a  $\mathbb{Q}$ -line bundle,  $(V, h)$  and  $(F, h_F)$  hermitian. Define

$$L_k = \mathcal{O}_{X_k^{\text{GG}}}(1) \otimes \pi_k^* \mathcal{O}\left(\frac{1}{kr} \left(1 + \frac{1}{2} + \dots + \frac{1}{k}\right) F\right),$$
$$\eta = \Theta_{\det V^*, \det h^*} + \Theta_{F, h_F}.$$

Then for all  $q \geq 0$  and all  $m \gg k \gg 1$  such that  $m$  is sufficiently divisible, we have

$$h^q(X_k^{\text{GG}}, \mathcal{O}(L_k^{\otimes m})) \leq \frac{m^{n+kr-1}}{(n+kr-1)!} \frac{(\log k)^n}{n! (k!)^r} \left( \int_{X(\eta, q)} (-1)^q \eta^n + \frac{C}{\log k} \right)$$

It follows that the leading term in the estimate only involves the trace of  $\Theta_{V^*, h^*}$ , i.e. the curvature of  $(\det V^*, \det h^*)$ , which can be taken to be  $> 0$  if  $\det V^*$  is big.

**Corollary (D-, 2010).** Let  $(X, V)$  be a directed manifold,  $F \rightarrow X$  a  $\mathbb{Q}$ -line bundle,  $(V, h)$  and  $(F, h_F)$  hermitian. Define

$$L_k = \mathcal{O}_{X_k^{\text{GG}}}(1) \otimes \pi_k^* \mathcal{O}\left(\frac{1}{kr}\left(1 + \frac{1}{2} + \dots + \frac{1}{k}\right)F\right),$$

$$\eta = \Theta_{\det V^*, \det h^*} + \Theta_{F, h_F}.$$

Then for all  $q \geq 0$  and all  $m \gg k \gg 1$  such that  $m$  is sufficiently divisible, we have

$$h^q(X_k^{\text{GG}}, \mathcal{O}(L_k^{\otimes m})) \leq \frac{m^{n+kr-1}}{(n+kr-1)!} \frac{(\log k)^n}{n! (k!)^r} \left( \int_{X(\eta, q)} (-1)^q \eta^n + \frac{C}{\log k} \right)$$

$$h^0(X_k^{\text{GG}}, \mathcal{O}(L_k^{\otimes m})) \geq \frac{m^{n+kr-1}}{(n+kr-1)!} \frac{(\log k)^n}{n! (k!)^r} \left( \int_{X(\eta, \leq 1)} \eta^n - \frac{C}{\log k} \right).$$

Let  $Z$  be an irreducible algebraic subset of some  $k$ -jet bundle  $X_k$  over  $X$ , such that  $Z$  projects onto  $X_{k-1}$ , i.e.

$$\pi_{k,k-1}(Z) = X_{k-1}.$$

Let  $Z$  be an irreducible algebraic subset of some  $k$ -jet bundle  $X_k$  over  $X$ , such that  $Z$  projects onto  $X_{k-1}$ , i.e.

$$\pi_{k,k-1}(Z) = X_{k-1}.$$

We define the linear subspace  $W \subset T_Z \subset T_{X_k|Z}$  to be the closure of  $T_{Z'} \cap V_k$  taken on a suitable Zariski open set  $Z' \subset Z_{\text{reg}}$  where the intersection has constant rank and is a subbundle of  $T_{Z'}$ .

Let  $Z$  be an irreducible algebraic subset of some  $k$ -jet bundle  $X_k$  over  $X$ , such that  $Z$  projects onto  $X_{k-1}$ , i.e.

$$\pi_{k,k-1}(Z) = X_{k-1}.$$

We define the linear subspace  $W \subset T_Z \subset T_{X_k|Z}$  to be the closure of  $T_{Z'} \cap V_k$  taken on a suitable Zariski open set  $Z' \subset Z_{\text{reg}}$  where the intersection has constant rank and is a subbundle of  $T_{Z'}$ .

Alternatively, one could also take  $W$  to be the closure of  $T_{Z'} \cap V_k$  in the  $k$ -th stage  $(\mathcal{X}_k, \mathcal{A}_k)$  of the “absolute Semple tower” associated with  $(\mathcal{X}_0, \mathcal{A}_0) = (X, T_X)$ .

Let  $Z$  be an irreducible algebraic subset of some  $k$ -jet bundle  $X_k$  over  $X$ , such that  $Z$  projects onto  $X_{k-1}$ , i.e.

$$\pi_{k,k-1}(Z) = X_{k-1}.$$

We define the linear subspace  $W \subset T_Z \subset T_{X_k|Z}$  to be the closure of  $T_{Z'} \cap V_k$  taken on a suitable Zariski open set  $Z' \subset Z_{\text{reg}}$  where the intersection has constant rank and is a subbundle of  $T_{Z'}$ .

Alternatively, one could also take  $W$  to be the closure of  $T_{Z'} \cap V_k$  in the  $k$ -th stage  $(\mathcal{X}_k, \mathcal{A}_k)$  of the “absolute Semple tower” associated with  $(\mathcal{X}_0, \mathcal{A}_0) = (X, T_X)$ .

This produces an induced directed pair

$$(Z, W) \subset (X_k, V_k),$$

and it is easy to show that  $\text{rk } W < \text{rk } V_k = \text{rk } V$ .

**Definition.** Let  $(X, V)$  be a directed pair where  $X$  is projective algebraic. We say that  $(X, V)$  is “strongly of general type” if it is of general type and for every irreducible component  $Z \subsetneq X_k$  that projects onto  $X_{k-1}$ ,  $X_k \not\subset D_k := P(T_{X_{k-1}/X_{k-2}})$ , the induced directed structure  $(Z, W) \subset (X_k, V_k)$  is of **general type modulo  $X_k \rightarrow X$** , i.e.  $K_W \otimes \mathcal{O}_{X_k}(m)|_Z$  is big for some  $m \in \mathbb{Q}_+$ .

**Definition.** Let  $(X, V)$  be a directed pair where  $X$  is projective algebraic. We say that  $(X, V)$  is “strongly of general type” if it is of general type and for every irreducible component  $Z \subsetneq X_k$  that projects onto  $X_{k-1}$ ,  $X_k \not\subset D_k := P(T_{X_{k-1}/X_{k-2}})$ , the induced directed structure  $(Z, W) \subset (X_k, V_k)$  is of **general type modulo  $X_k \rightarrow X$** , i.e.  $K_W \otimes \mathcal{O}_{X_k}(m)|_Z$  is big for some  $m \in \mathbb{Q}_+$ .

**Theorem (D-, 2014)** If  $(X, V)$  is strongly of general type, the **Green-Griffiths-Lang conjecture holds true** for  $(X, V)$ , namely there  $\exists Y \subsetneq X$  such that every non constant holomorphic curve  $f : (\mathbb{C}, T_{\mathbb{C}} \rightarrow (X, V)$  satisfies  $f(\mathbb{C}) \subset Y$ .



**Definition.** Let  $(X, V)$  be a directed pair where  $X$  is projective algebraic. We say that  $(X, V)$  is “strongly of general type” if it is of general type and for every irreducible component  $Z \subsetneq X_k$  that projects onto  $X_{k-1}$ ,  $X_k \not\subset D_k := P(T_{X_{k-1}/X_{k-2}})$ , the induced directed structure  $(Z, W) \subset (X_k, V_k)$  is of **general type modulo  $X_k \rightarrow X$** , i.e.  $K_W \otimes \mathcal{O}_{X_k}(m)|_Z$  is big for some  $m \in \mathbb{Q}_+$ .

**Theorem (D-, 2014)** If  $(X, V)$  is strongly of general type, the **Green-Griffiths-Lang conjecture holds true** for  $(X, V)$ , namely there  $\exists Y \subsetneq X$  such that every non constant holomorphic curve  $f : (\mathbb{C}, T_{\mathbb{C}} \rightarrow (X, V)$  satisfies  $f(\mathbb{C}) \subset Y$ .

**Proof:** Induction using existence theorem for jet differentials. Unfortunately by contradiction, and thus non constructive.

**Definition.** Fix an ample divisor  $A$  on  $X$ . For every irreducible subvariety  $Z \subset X_k$  that projects onto  $X_{k-1}$  for  $k \geq 1$ ,  $Z \not\subset D_k$ , and  $Z = X = X_0$  for  $k = 0$ , we define the **slope** of the corresponding directed variety  $(Z, W)$  to be  $\mu_A(Z, W) =$

$$\frac{\inf \{ \lambda \in \mathbb{Q}; \exists m \in \mathbb{Q}_+, K_W \otimes (\mathcal{O}_{X_k}(m) \otimes \pi_{k,0}^* \mathcal{O}(\lambda A))|_Z \text{ big on } Z \}}{\text{rank } W}.$$

**Definition.** Fix an ample divisor  $A$  on  $X$ . For every irreducible subvariety  $Z \subset X_k$  that projects onto  $X_{k-1}$  for  $k \geq 1$ ,  $Z \not\subset D_k$ , and  $Z = X = X_0$  for  $k = 0$ , we define the **slope** of the corresponding directed variety  $(Z, W)$  to be  $\mu_A(Z, W) =$

$$\frac{\inf \{ \lambda \in \mathbb{Q}; \exists m \in \mathbb{Q}_+, K_W \otimes (\mathcal{O}_{X_k}(m) \otimes \pi_{k,0}^* \mathcal{O}(\lambda A))|_Z \text{ big on } Z \}}{\text{rank } W}.$$

Notice that  $(X, V)$  is of general type iff  $\mu_A(X, V) < 0$ .

**Definition.** Fix an ample divisor  $A$  on  $X$ . For every irreducible subvariety  $Z \subset X_k$  that projects onto  $X_{k-1}$  for  $k \geq 1$ ,  $Z \not\subset D_k$ , and  $Z = X = X_0$  for  $k = 0$ , we define the **slope** of the corresponding directed variety  $(Z, W)$  to be  $\mu_A(Z, W) =$

$$\frac{\inf \{ \lambda \in \mathbb{Q}; \exists m \in \mathbb{Q}_+, K_W \otimes (\mathcal{O}_{X_k}(m) \otimes \pi_{k,0}^* \mathcal{O}(\lambda A))|_Z \text{ big on } Z \}}{\text{rank } W}.$$

Notice that  $(X, V)$  is of general type iff  $\mu_A(X, V) < 0$ .

We say that  $(X, V)$  is  **$A$ -jet-stable** (resp.  **$A$ -jet-semi-stable**) if  $\mu_A(Z, W) < \mu_A(X, V)$  (resp.  $\mu_A(Z, W) \leq \mu_A(X, V)$ ) for all  $Z \subsetneq X_k$  as above.

**Definition.** Fix an ample divisor  $A$  on  $X$ . For every irreducible subvariety  $Z \subset X_k$  that projects onto  $X_{k-1}$  for  $k \geq 1$ ,  $Z \not\subset D_k$ , and  $Z = X = X_0$  for  $k = 0$ , we define the **slope** of the corresponding directed variety  $(Z, W)$  to be  $\mu_A(Z, W) =$

$$\frac{\inf \{ \lambda \in \mathbb{Q}; \exists m \in \mathbb{Q}_+, K_W \otimes (\mathcal{O}_{X_k}(m) \otimes \pi_{k,0}^* \mathcal{O}(\lambda A))|_Z \text{ big on } Z \}}{\text{rank } W}.$$

Notice that  $(X, V)$  is of general type iff  $\mu_A(X, V) < 0$ .

We say that  $(X, V)$  is **A-jet-stable** (resp. **A-jet-semi-stable**) if  $\mu_A(Z, W) < \mu_A(X, V)$  (resp.  $\mu_A(Z, W) \leq \mu_A(X, V)$ ) for all  $Z \subsetneq X_k$  as above.

**Observation.** If  $(X, V)$  is of general type and A-jet-semi-stable, then  $(X, V)$  is strongly of general type.