



# Hyperbolic algebraic varieties and holomorphic differential equations

Jean-Pierre Demailly

Institut Fourier, Université de Grenoble I, France  
& Académie des Sciences de Paris

August 26, 2012 / VIASM Yearly Meeting, Hanoi

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.  
 $X$  is said to be **(Brody) hyperbolic** if  $\nexists$  such  $f : \mathbb{C} \rightarrow X$ .

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.  
 $X$  is said to be **(Brody) hyperbolic** if  $\nexists$  such  $f : \mathbb{C} \rightarrow X$ .
- If  $X$  is a **bounded** open subset  $\Omega \subset \mathbb{C}^n$ , then there are no entire curves  $f : \mathbb{C} \rightarrow \Omega$  (**Liouville's theorem**),  
 $\Rightarrow$  every bounded open set  $\Omega \subset \mathbb{C}^n$  is hyperbolic

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.  
 $X$  is said to be **(Brody) hyperbolic** if  $\nexists$  such  $f : \mathbb{C} \rightarrow X$ .
- If  $X$  is a **bounded** open subset  $\Omega \subset \mathbb{C}^n$ , then there are no entire curves  $f : \mathbb{C} \rightarrow \Omega$  (**Liouville's theorem**),  
 $\Rightarrow$  every bounded open set  $\Omega \subset \mathbb{C}^n$  is hyperbolic
- $X = \overline{\mathbb{C}} \setminus \{0, 1, \infty\} = \mathbb{C} \setminus \{0, 1\}$  has no entire curves, so it is hyperbolic (**Picard's theorem**)

- **Definition.** By an **entire curve** we mean a non constant holomorphic map  $f : \mathbb{C} \rightarrow X$  into a complex  $n$ -dimensional manifold.  
 $X$  is said to be **(Brody) hyperbolic** if  $\nexists$  such  $f : \mathbb{C} \rightarrow X$ .
- If  $X$  is a **bounded** open subset  $\Omega \subset \mathbb{C}^n$ , then there are no entire curves  $f : \mathbb{C} \rightarrow \Omega$  (**Liouville's theorem**),  
 $\Rightarrow$  every bounded open set  $\Omega \subset \mathbb{C}^n$  is hyperbolic
- $X = \overline{\mathbb{C}} \setminus \{0, 1, \infty\} = \mathbb{C} \setminus \{0, 1\}$  has no entire curves, so it is hyperbolic (**Picard's theorem**)
- A complex torus  $X = \mathbb{C}^n / \Lambda$  ( $\Lambda$  lattice) has a lot of entire curves. As  $\mathbb{C}$  simply connected, every  $f : \mathbb{C} \rightarrow X = \mathbb{C}^n / \Lambda$  lifts as  $\tilde{f} : \mathbb{C} \rightarrow \mathbb{C}^n$ ,  $\tilde{f}(t) = (\tilde{f}_1(t), \dots, \tilde{f}_n(t))$ , and  $\tilde{f}_j : \mathbb{C} \rightarrow \mathbb{C}$  can be arbitrary entire functions.

- Consider now the complex projective  $n$ -space

$$\mathbb{P}^n = \mathbb{P}_{\mathbb{C}}^n = (\mathbb{C}^{n+1} \setminus \{0\})/\mathbb{C}^*, \quad [z] = [z_0 : z_1 : \dots : z_n].$$

- Consider now the complex projective  $n$ -space

$$\mathbb{P}^n = \mathbb{P}_{\mathbb{C}}^n = (\mathbb{C}^{n+1} \setminus \{0\})/\mathbb{C}^*, \quad [z] = [z_0 : z_1 : \dots : z_n].$$

- An entire curve  $f : \mathbb{C} \rightarrow \mathbb{P}^n$  is given by a map

$$t \longmapsto [f_0(t) : f_1(t) : \dots : f_n(t)]$$

where  $f_j : \mathbb{C} \rightarrow \mathbb{C}$  are holomorphic functions without common zeroes (so there are a lot of them).



- Consider now the complex projective  $n$ -space

$$\mathbb{P}^n = \mathbb{P}_{\mathbb{C}}^n = (\mathbb{C}^{n+1} \setminus \{0\})/\mathbb{C}^*, \quad [z] = [z_0 : z_1 : \dots : z_n].$$

- An entire curve  $f : \mathbb{C} \rightarrow \mathbb{P}^n$  is given by a map


$$t \longmapsto [f_0(t) : f_1(t) : \dots : f_n(t)]$$

where  $f_j : \mathbb{C} \rightarrow \mathbb{C}$  are holomorphic functions without common zeroes (so there are a lot of them).


- More generally, look at a (complex) **projective manifold**, i.e.

$$X^n \subset \mathbb{P}^N, \quad X = \{[z]; P_1(z) = \dots = P_k(z) = 0\}$$

where  $P_j(z) = P_j(z_0, z_1, \dots, z_N)$  are homogeneous polynomials (of some degree  $d_j$ ), such that  $X$  is **non singular**.




$g = 0, K_X < 0$   
(positive curvature)




$g = 1, K_X = 0$   
(zero curvature)

Canonical bundle  $K_X = \Lambda^n T_X^*$  (here  $K_X = T_X^*$ )




$g = 0, K_X < 0$   
(positive curvature)




$g = 1, K_X = 0$   
(zero curvature)

Canonical bundle  $K_X = \Lambda^n T_X^*$  (here  $K_X = T_X^*$ )

- $g = 0 : X = \mathbb{P}^1$       courbure  $T_X > 0$  **not hyperbolic**




$g = 0, K_X < 0$   
(positive curvature)



$g = 1, K_X = 0$   
(zero curvature)

Canonical bundle  $K_X = \Lambda^n T_X^*$  (here  $K_X = T_X^*$ )


- $g = 0 : X = \mathbb{P}^1$       courbure  $T_X > 0$  **not hyperbolic**
- $g = 1 : X = \mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$       courbure  $T_X = 0$  **not hyperbolic**



$g > 1, K_X > 0$   
(negative curvature)

$$\deg K_X = 2g - 2$$

If  $g \geq 2$ ,  $X \simeq \mathbb{D}/\Gamma$  ( $T_X < 0$ )  $\Rightarrow$   $X$  is hyperbolic.



$g > 1, K_X > 0$   
(negative curvature)

$$\deg K_X = 2g - 2$$

If  $g \geq 2$ ,  $X \simeq \mathbb{D}/\Gamma$  ( $T_X < 0$ )  $\Rightarrow$   $X$  is hyperbolic.

In fact every curve  $f : \mathbb{C} \rightarrow X \simeq \mathbb{D}/\Gamma$  lifts to  $\tilde{f} : \mathbb{C} \rightarrow \mathbb{D}$ ,  
and so must be constant by Liouville.

- For a complex manifold,  $n = \dim_{\mathbb{C}} X$ , one defines the **Kobayashi pseudo-metric** :  $x \in X$ ,  $\xi \in T_x$

$$\kappa_x(\xi) = \inf\{\lambda > 0; \exists f : \mathbb{D} \rightarrow X, f(0) = x, \lambda f_*(0) = \xi\}$$

On  $\mathbb{C}^n$ ,  $\mathbb{P}^n$  or complex tori  $X = \mathbb{C}^n/\Lambda$ , one has  $\kappa_X \equiv 0$ .

- For a complex manifold,  $n = \dim_{\mathbb{C}} X$ , one defines **the Kobayashi pseudo-metric** :  $x \in X, \xi \in T_x$   
$$\kappa_x(\xi) = \inf\{\lambda > 0; \exists f : \mathbb{D} \rightarrow X, f(0) = x, \lambda f_*(0) = \xi\}$$
On  $\mathbb{C}^n, \mathbb{P}^n$  or complex tori  $X = \mathbb{C}^n/\Lambda$ , one has  $\kappa_X \equiv 0$ .
- $X$  is said to be **hyperbolic in the sense of Kobayashi** if the associated integrated pseudo-distance is a distance (i.e. it separates points – i.e. has Hausdorff topology).



- For a complex manifold,  $n = \dim_{\mathbb{C}} X$ , one defines **the Kobayashi pseudo-metric** :  $x \in X, \xi \in T_x$   

$$\kappa_x(\xi) = \inf\{\lambda > 0; \exists f : \mathbb{D} \rightarrow X, f(0) = x, \lambda f_*(0) = \xi\}$$
 On  $\mathbb{C}^n, \mathbb{P}^n$  or complex tori  $X = \mathbb{C}^n/\Lambda$ , one has  $\kappa_X \equiv 0$ .
- $X$  is said to be **hyperbolic in the sense of Kobayashi** if the associated integrated pseudo-distance is a distance (i.e. it separates points – i.e. has Hausdorff topology).
- **Examples.** \*  $X = \Omega/\Gamma$ ,  $\Omega$  bounded symmetric domain.  
 \* any product  $X = X_1 \times \dots \times X_s$  where  $X_j$  hyperbolic.

- For a complex manifold,  $n = \dim_{\mathbb{C}} X$ , one defines **the Kobayashi pseudo-metric** :  $x \in X, \xi \in T_x$   

$$\kappa_x(\xi) = \inf\{\lambda > 0; \exists f : \mathbb{D} \rightarrow X, f(0) = x, \lambda f_*(0) = \xi\}$$
 On  $\mathbb{C}^n, \mathbb{P}^n$  or complex tori  $X = \mathbb{C}^n/\Lambda$ , one has  $\kappa_X \equiv 0$ .
- $X$  is said to be **hyperbolic in the sense of Kobayashi** if the associated integrated pseudo-distance is a distance (i.e. it separates points – i.e. has Hausdorff topology).
- **Examples.** \*  $X = \Omega/\Gamma$ ,  $\Omega$  bounded symmetric domain.  
 \* any product  $X = X_1 \times \dots \times X_s$  where  $X_j$  hyperbolic.
- **Theorem (dimension  $n$  arbitrary)** (Kobayashi, 1970)  
 $T_x$  *negatively curved* ( $T_x^* > 0$ , i.e. *ample*)  $\Rightarrow X$  *hyperbolic*.  
 Recall that a holomorphic vector bundle  $E$  is **ample** iff its symmetric powers  $S^m E$  have global sections which generate 1-jets of (germs of) sections at any point  $x \in X$ .

The proof of the above Kobayashi result depends crucially on:

**Ahlfors-Schwarz lemma.** Let  $\gamma = i \sum \gamma_{jk} dt_j \wedge d\bar{t}_k$  be an almost everywhere positive hermitian form on the ball  $B(0, R) \subset \mathbb{C}^p$ , such that  $-\text{Ricci}(\gamma) := i \partial \bar{\partial} \log \det \gamma \geq A\gamma$  in the sense of currents, for some constant  $A > 0$  (this means in particular that  $\det \gamma = \det(\gamma_{jk})$  is such that  $\log \det \gamma$  is plurisubharmonic). Then the  $\gamma$ -volume form is controlled by the Poincaré volume form :

$$\det(\gamma) \leq \left( \frac{p+1}{AR^2} \right)^p \frac{1}{(1 - |t|^2/R^2)^{p+1}}.$$

**Brody reparametrization Lemma.** *Assume that  $X$  is compact, let  $\omega$  be a hermitian metric on  $X$  and  $f : \mathbb{D} \rightarrow X$  a holomorphic map. For every  $\varepsilon > 0$ , there exists a radius  $R \geq (1 - \varepsilon)\|f'(0)\|_\omega$  and a homographic transformation  $\psi$  of the disk  $D(0, R)$  onto  $(1 - \varepsilon)\mathbb{D}$  such that  $\|(f \circ \psi)'(0)\|_\omega = 1$  and  $\|(f \circ \psi)'(t)\|_\omega \leq (1 - |t|^2/R^2)^{-1}$  for every  $t \in D(0, R)$ .  
 $\Rightarrow$  if  $f'$  unbounded,  $\exists g = \lim f \circ \psi_\nu : \mathbb{C} \rightarrow X$  with  $\|g'\|_\omega \leq 1$ .*

**Brody reparametrization Lemma.** Assume that  $X$  is *compact*, let  $\omega$  be a hermitian metric on  $X$  and  $f : \mathbb{D} \rightarrow X$  a holomorphic map. For every  $\varepsilon > 0$ , there exists a radius  $R \geq (1 - \varepsilon)\|f'(0)\|_\omega$  and a homographic transformation  $\psi$  of the disk  $D(0, R)$  onto  $(1 - \varepsilon)\mathbb{D}$  such that  $\|(f \circ \psi)'(0)\|_\omega = 1$  and  $\|(f \circ \psi)'(t)\|_\omega \leq (1 - |t|^2/R^2)^{-1}$  for every  $t \in D(0, R)$ .  
 $\Rightarrow$  if  $f'$  unbounded,  $\exists g = \lim f \circ \psi_\nu : \mathbb{C} \rightarrow X$  with  $\|g'\|_\omega \leq 1$ .

**Brody theorem (1978).** If  $X$  is *compact* then  $X$  is Kobayashi hyperbolic if and only if there are no entire holomorphic curves  $f : \mathbb{C} \rightarrow X$  (*Brody hyperbolicity*).

**Brody reparametrization Lemma.** Assume that  $X$  is *compact*, let  $\omega$  be a hermitian metric on  $X$  and  $f : \mathbb{D} \rightarrow X$  a holomorphic map. For every  $\varepsilon > 0$ , there exists a radius  $R \geq (1 - \varepsilon) \|f'(0)\|_\omega$  and a homographic transformation  $\psi$  of the disk  $D(0, R)$  onto  $(1 - \varepsilon)\mathbb{D}$  such that  $\|(f \circ \psi)'(0)\|_\omega = 1$  and  $\|(f \circ \psi)'(t)\|_\omega \leq (1 - |t|^2/R^2)^{-1}$  for every  $t \in D(0, R)$ .  
 $\Rightarrow$  if  $f'$  unbounded,  $\exists g = \lim f \circ \psi_\nu : \mathbb{C} \rightarrow X$  with  $\|g'\|_\omega \leq 1$ .

**Brody theorem (1978).** If  $X$  is *compact* then  $X$  is Kobayashi hyperbolic if and only if there are no entire holomorphic curves  $f : \mathbb{C} \rightarrow X$  (*Brody hyperbolicity*).

Hyperbolic varieties are especially interesting for their expected diophantine properties :

**Conjecture** (S. Lang, 1986) An arithmetic projective variety  $X$  is hyperbolic iff  $X(\mathbb{K})$  is finite for every number field  $\mathbb{K}$ .

- **Definition** A non singular projective variety  $X$  is said to be of *general type* if the growth of pluricanonical sections

$$\dim H^0(X, K_X^{\otimes m}) \sim cm^n, \quad K_X = \Lambda^n T_X^*$$

is maximal.

(sections locally of the form  $f(z) (dz_1 \wedge \dots \wedge dz_n)^{\otimes m}$ )

**Example:** A non singular hypersurface  $X^n \subset \mathbb{P}^{n+1}$  of degree  $d$  satisfies  $K_X = \mathcal{O}(d - n - 2)$ ,  
 $X$  is of general type iff  $d > n + 2$ .

- **Definition** A non singular projective variety  $X$  is said to be of *general type* if the growth of pluricanonical sections

$$\dim H^0(X, K_X^{\otimes m}) \sim cm^n, \quad K_X = \Lambda^n T_X^*$$

is maximal.

(sections locally of the form  $f(z) (dz_1 \wedge \dots \wedge dz_n)^{\otimes m}$ )

**Example:** A non singular hypersurface  $X^n \subset \mathbb{P}^{n+1}$  of degree  $d$  satisfies  $K_X = \mathcal{O}(d - n - 2)$ ,  
 $X$  is of general type iff  $d > n + 2$ .

- **Conjecture CGT.** If a compact variety  $X$  is hyperbolic, then it should be of general type, and if  $X$  is non singular, then  $K_X = \Lambda^n T_X^*$  should be ample, i.e.  $K_X > 0$  (Kodaira) (equivalently  $\exists$  Kähler metric  $\omega$  such that  $\text{Ricci}(\omega) < 0$ ).



- **Theorem.** *Let  $X$  be projective algebraic. Consider the following properties :*

(GT) *Every subvariety  $Y$  of  $X$  is of **general type**.*

(AH)  $\exists \varepsilon > 0, \forall C \subset X$  *algebraic curve*

$$2g(\bar{C}) - 2 \geq \varepsilon \deg(C).$$

( $X$  *“algebraically hyperbolic”*)

(HY)  $X$  is *hyperbolic*

(JC)  $X$  possesses a *jet-metric with negative curvature* on its  $k$ -jet bundle  $X_k$  [to be defined later], for  $k \geq k_0 \gg 1$ .

Then (JC)  $\Rightarrow$  (GT), (AH), (HY),

(HY)  $\Rightarrow$  (AH),

and if Conjecture CGT holds, (HY)  $\Rightarrow$  (GT).

- **Theorem.** Let  $X$  be projective algebraic. Consider the following properties :

(GT) Every subvariety  $Y$  of  $X$  is of *general type*.

(AH)  $\exists \varepsilon > 0, \forall C \subset X$  algebraic curve

$$2g(\bar{C}) - 2 \geq \varepsilon \deg(C).$$

( $X$  “*algebraically hyperbolic*”)

(HY)  $X$  is *hyperbolic*

(JC)  $X$  possesses a *jet-metric with negative curvature* on its  $k$ -jet bundle  $X_k$  [to be defined later], for  $k \geq k_0 \gg 1$ .

Then (JC)  $\Rightarrow$  (GT), (AH), (HY),

(HY)  $\Rightarrow$  (AH),

and if Conjecture CGT holds, (HY)  $\Rightarrow$  (GT).

- It is expected that all 4 properties are in fact equivalent for projective varieties.

- **Conjecture** (Green-Griffiths-Lang = GGL) *Let  $X$  be a projective variety of general type. Then there exists an algebraic variety  $Y \subsetneq X$  such that for all non-constant holomorphic  $f : \mathbb{C} \rightarrow X$  one has  $f(\mathbb{C}) \subset Y$ .*

- **Conjecture** (Green-Griffiths-Lang = GGL) *Let  $X$  be a projective variety of general type. Then there exists an algebraic variety  $Y \subsetneq X$  such that for all non-constant holomorphic  $f : \mathbb{C} \rightarrow X$  one has  $f(\mathbb{C}) \subset Y$ .*
- Combining the above conjectures, we get :  
**Expected consequence** (of CGT + GGL) Properties:  
(HY)  $X$  is *hyperbolic*  
(GT) *Every subvariety  $Y$  of  $X$  is of general type*  
*are equivalent if CGT + GGL hold.*

- **Conjecture** (Green-Griffiths-Lang = GGL) *Let  $X$  be a projective variety of general type. Then there exists an algebraic variety  $Y \subsetneq X$  such that for all non-constant holomorphic  $f : \mathbb{C} \rightarrow X$  one has  $f(\mathbb{C}) \subset Y$ .*
- Combining the above conjectures, we get :  
**Expected consequence** (of CGT + GGL) Properties:  
(HY)  $X$  is *hyperbolic*  
(GT) Every subvariety  $Y$  of  $X$  is of *general type*  
are equivalent if CGT + GGL hold.
- **Arithmetic counterpart** (Lang 1987). *If  $X$  is a variety of general type defined over a number field and  $Y$  is the Green-Griffiths locus (Zariski closure of  $\bigcup f(\mathbb{C})$ ), then  $X(\mathbb{K}) \setminus Y$  is finite for every number field  $\mathbb{K}$ .*

- Using “jet technology” and **deep results of McQuillan** for curve foliations on surfaces, D. – El Goul proved **Theorem** (solution of Kobayashi conjecture, 1998).  
*A very generic surface  $X \subset \mathbb{P}^3$  of degree  $\geq 21$  is hyperbolic.*  
 Independently McQuillan got degree  $\geq 35$ .  
 Recently improved to **degree  $\geq 18$**  (Păun, 2008).  
 For  $X \subset \mathbb{P}^{n+1}$ , the optimal bound should be **degree  $\geq 2n + 1$  for  $n \geq 2$  (Zaidenberg)**.
- **Generic GGL conjecture for  $\dim_{\mathbb{C}} X = n$**   
 (S. Diverio, J. Merker, E. Rousseau, 2009).  
*If  $X \subset \mathbb{P}^{n+1}$  is a generic  $n$ -fold of degree  $d \geq d_n := 2^{n^5}$ ,*  
**[also  $d_3 = 593, d_4 = 3203, d_5 = 35355, d_6 = 172925$ ]**  
 then  $\exists Y \subsetneq X$  s.t.  $\forall$  non const.  $f: \mathbb{C} \rightarrow X$  satisfies  $f(\mathbb{C}) \subset Y$

- Using “jet technology” and **deep results of McQuillan** for curve foliations on surfaces, D. – El Goul proved **Theorem** (solution of Kobayashi conjecture, 1998).  
*A very generic surface  $X \subset \mathbb{P}^3$  of degree  $\geq 21$  is hyperbolic.*  
 Independently McQuillan got degree  $\geq 35$ .  
 Recently improved to **degree  $\geq 18$**  (Păun, 2008).  
 For  $X \subset \mathbb{P}^{n+1}$ , the optimal bound should be **degree  $\geq 2n + 1$  for  $n \geq 2$  (Zaidenberg)**.
- Generic GGL conjecture for  $\dim_{\mathbb{C}} X = n$**   
 (S. Diverio, J. Merker, E. Rousseau, 2009).  
*If  $X \subset \mathbb{P}^{n+1}$  is a generic  $n$ -fold of degree  $d \geq d_n := 2^{n^5}$ ,*  
*[also  $d_3 = 593, d_4 = 3203, d_5 = 35355, d_6 = 172925$ ]*  
 then  $\exists Y \subsetneq X$  s.t.  $\forall$  non const.  $f: \mathbb{C} \rightarrow X$  satisfies  $f(\mathbb{C}) \subset Y$   
**Moreover** (S. Diverio, S. Trapani, 2009)  **$\text{codim}_{\mathbb{C}} Y \geq 2 \Rightarrow$**   
 generic hypersurface  $X \subset \mathbb{P}^4$  of degree  $\geq 593$  is **hyperbolic**.

The main idea in order to attack GGL is to use differential equations. Let

$$\mathbb{C} \rightarrow X, \quad t \mapsto f(t) = (f_1(t), \dots, f_n(t))$$

be a curve written in some local holomorphic coordinates  $(z_1, \dots, z_n)$  on  $X$ .



The main idea in order to attack GGL is to use differential equations. Let

$$\mathbb{C} \rightarrow X, \quad t \mapsto f(t) = (f_1(t), \dots, f_n(t))$$

be a curve written in some local holomorphic coordinates  $(z_1, \dots, z_n)$  on  $X$ .

Consider **algebraic differential operators** which can be written locally in multi-index notation

$$\begin{aligned} P(f_{[k]}) &= P(f', f'', \dots, f^{(k)}) \\ &= \sum a_{\alpha_1 \alpha_2 \dots \alpha_k} (f(t)) f'(t)^{\alpha_1} f''(t)^{\alpha_2} \dots f^{(k)}(t)^{\alpha_k} \end{aligned}$$

where  $a_{\alpha_1 \alpha_2 \dots \alpha_k}(z)$  are holomorphic coefficients on  $X$  and  $t \mapsto z = f(t)$  is a curve,  $f_{[k]} = (f', f'', \dots, f^{(k)})$  its  **$k$ -jet**.

The main idea in order to attack GGL is to use differential equations. Let

$$\mathbb{C} \rightarrow X, \quad t \mapsto f(t) = (f_1(t), \dots, f_n(t))$$

be a curve written in some local holomorphic coordinates  $(z_1, \dots, z_n)$  on  $X$ .

Consider **algebraic differential operators** which can be written locally in multi-index notation

$$\begin{aligned} P(f_{[k]}) &= P(f', f'', \dots, f^{(k)}) \\ &= \sum a_{\alpha_1 \alpha_2 \dots \alpha_k} (f(t)) f'(t)^{\alpha_1} f''(t)^{\alpha_2} \dots f^{(k)}(t)^{\alpha_k} \end{aligned}$$

where  $a_{\alpha_1 \alpha_2 \dots \alpha_k}(z)$  are holomorphic coefficients on  $X$  and  $t \mapsto z = f(t)$  is a curve,  $f_{[k]} = (f', f'', \dots, f^{(k)})$  its  **$k$ -jet**.

Obvious  $\mathbb{C}^*$ -action :

$$\lambda \cdot f(t) = f(\lambda t), \quad (\lambda \cdot f)^{(k)}(t) = \lambda^k f^{(k)}(\lambda t)$$

- **Definition.**  $E_{k,m}^{\text{GG}}$  is the sheaf (bundle) of algebraic differential operators of order  $k$  and weighted degree  $m$ .

- **Definition.**  $E_{k,m}^{\text{GG}}$  is the sheaf (bundle) of algebraic differential operators of order  $k$  and weighted degree  $m$ .
- **Fundamental vanishing theorem**  
[Green-Griffiths 1979], [Demailly 1995], [Siu-Yeung 1996]  
Let  $P \in H^0(X, E_{k,m}^{\text{GG}} \otimes \mathcal{O}(-A))$  be a global algebraic differential operator whose coefficients vanish on some ample divisor  $A$ . Then  $\forall f : \mathbb{C} \rightarrow X, P(f_{[k]}) \equiv 0$ .

- **Definition.**  $E_{k,m}^{\text{GG}}$  is the sheaf (bundle) of algebraic differential operators of order  $k$  and weighted degree  $m$ .
- **Fundamental vanishing theorem**  
[Green-Griffiths 1979], [Demailly 1995], [Siu-Yeung 1996]  
Let  $P \in H^0(X, E_{k,m}^{\text{GG}} \otimes \mathcal{O}(-A))$  be a global algebraic differential operator whose coefficients vanish on some ample divisor  $A$ . Then  $\forall f : \mathbb{C} \rightarrow X, P(f_{[k]}) \equiv 0$ .
- **Proof.** One can assume that  $A$  is very ample and intersects  $f(\mathbb{C})$ . Also assume  $f'$  bounded (this is not so restrictive by Brody !). Then all  $f^{(k)}$  are bounded by Cauchy inequality. Hence

$$\mathbb{C} \ni t \mapsto P(f', f'', \dots, f^{(k)})(t)$$

is a bounded holomorphic function on  $\mathbb{C}$  which vanishes at some point. Apply Liouville's theorem !



- Let  $X_k^{\text{GG}} = J_k(X)^*/\mathbb{C}^*$  be the **projectivized  $k$ -jet bundle** of  $X =$  quotient of non constant  $k$ -jets by  $\mathbb{C}^*$ -action. Fibers are weighted projective spaces.

**Observation.** If  $\pi_k : X_k^{\text{GG}} \rightarrow X$  is canonical projection and  $\mathcal{O}_{X_k^{\text{GG}}}(1)$  is the **tautological line bundle**, then

$$E_{k,m}^{\text{GG}} = (\pi_k)_* \mathcal{O}_{X_k^{\text{GG}}}(m)$$

- Let  $X_k^{\text{GG}} = J_k(X)^*/\mathbb{C}^*$  be the **projectivized  $k$ -jet bundle** of  $X =$  quotient of non constant  $k$ -jets by  $\mathbb{C}^*$ -action. Fibers are weighted projective spaces.

**Observation.** If  $\pi_k : X_k^{\text{GG}} \rightarrow X$  is canonical projection and  $\mathcal{O}_{X_k^{\text{GG}}}(1)$  is the **tautological line bundle**, then

$$E_{k,m}^{\text{GG}} = (\pi_k)_* \mathcal{O}_{X_k^{\text{GG}}}(m)$$

- Saying that  $f : \mathbb{C} \rightarrow X$  satisfies the differential equation  $P(f_{[k]}) = 0$  means that

$$f_{[k]}(\mathbb{C}) \subset Z_P$$

where  $Z_P$  is the zero divisor of the section

$$\sigma_P \in H^0(X_k^{\text{GG}}, \mathcal{O}_{X_k^{\text{GG}}}(m) \otimes \pi_k^* \mathcal{O}(-A))$$

associated with  $P$ .

- **Consequence of fundamental vanishing theorem.**

*If  $P_j \in H^0(X, E_{k,m}^{GG} \otimes \mathcal{O}(-A))$  is a basis of sections then the image  $f(\mathbb{C})$  lies in  $Y = \pi_k(\bigcap Z_{P_j})$ , hence property asserted by the GGL conjecture holds true if there are “enough independent differential equations” so that*

$$Y = \pi_k\left(\bigcap_j Z_{P_j}\right) \subsetneq X.$$



- **Consequence of fundamental vanishing theorem.**

*If  $P_j \in H^0(X, E_{k,m}^{GG} \otimes \mathcal{O}(-A))$  is a basis of sections then the image  $f(\mathbb{C})$  lies in  $Y = \pi_k(\bigcap Z_{P_j})$ , hence property asserted by the GGL conjecture holds true if there are “enough independent differential equations” so that*

$$Y = \pi_k\left(\bigcap_j Z_{P_j}\right) \subsetneq X.$$

- However, **some differential equations are not very useful.** On a surface with coordinates  $(z_1, z_2)$ , a Wronskian equation  $f_1' f_2'' - f_2' f_1'' = 0$  tells us that  $f(\mathbb{C})$  sits on a line, but  $f_2''(t) = 0$  says that the second component is linear affine in time, an essentially **meaningless information** which is lost by a change of parameter  $t \mapsto \varphi(t)$ .

- The  $k$ -th order Wronskian operator

$$W_k(f) = f' \wedge f'' \wedge \dots \wedge f^{(k)}$$

(locally defined in coordinates) has degree  $m = \frac{k(k+1)}{2}$   
and

$$W_k(f \circ \varphi) = \varphi'^m W_k(f) \circ \varphi.$$

- The  $k$ -th order Wronskian operator

$$W_k(f) = f' \wedge f'' \wedge \dots \wedge f^{(k)}$$

(locally defined in coordinates) has degree  $m = \frac{k(k+1)}{2}$   
and

$$W_k(f \circ \varphi) = \varphi'^m W_k(f) \circ \varphi.$$

- **Definition.** A differential operator  $P$  of order  $k$  and degree  $m$  is said to be invariant by reparametrization if

$$P(f \circ \varphi) = \varphi'^m P(f) \circ \varphi$$

for any parameter change  $t \mapsto \varphi(t)$ . Consider their set

$$E_{k,m} \subset E_{k,m}^{\text{GG}} \quad (\text{a subbundle})$$

(Any polynomial  $Q(W_1, W_2, \dots, W_k)$  is invariant, but for  $k \geq 3$  there are other invariant operators.)

- **Goal.** We are interested in curves  $f : \mathbb{C} \rightarrow X$  such that  $f'(\mathbb{C}) \subset V$  where  $V$  is a subbundle (or subsheaf) of  $T_X$ .

- **Goal.** We are interested in curves  $f : \mathbb{C} \rightarrow X$  such that  $f'(\mathbb{C}) \subset V$  where  $V$  is a subbundle (or subsheaf) of  $T_X$ .
- **Definition.** *Category of directed manifolds :*
  - **Objects** : pairs  $(X, V)$ ,  $X$  manifold/ $\mathbb{C}$  and  $V \subset \mathcal{O}(T_X)$
  - **Arrows**  $\psi : (X, V) \rightarrow (Y, W)$  holomorphic s.t.  $\psi_* V \subset W$

- **Goal.** We are interested in curves  $f : \mathbb{C} \rightarrow X$  such that  $f'(\mathbb{C}) \subset V$  where  $V$  is a subbundle (or subsheaf) of  $T_X$ .
- **Definition.** *Category of directed manifolds :*
  - **Objects** : pairs  $(X, V)$ ,  $X$  manifold/ $\mathbb{C}$  and  $V \subset \mathcal{O}(T_X)$
  - **Arrows**  $\psi : (X, V) \rightarrow (Y, W)$  holomorphic s.t.  $\psi_* V \subset W$
  - “**Absolute case**”  $(X, T_X)$
  - “**Relative case**”  $(X, T_{X/S})$  where  $X \rightarrow S$
  - “**Integrable case**” when  $[V, V] \subset V$  (foliations)

- **Goal.** We are interested in curves  $f : \mathbb{C} \rightarrow X$  such that  $f'(\mathbb{C}) \subset V$  where  $V$  is a subbundle (or subsheaf) of  $T_X$ .
- **Definition.** *Category of directed manifolds :*
  - **Objects** : pairs  $(X, V)$ ,  $X$  manifold/ $\mathbb{C}$  and  $V \subset \mathcal{O}(T_X)$
  - **Arrows**  $\psi : (X, V) \rightarrow (Y, W)$  holomorphic s.t.  $\psi_* V \subset W$
  - “**Absolute case**”  $(X, T_X)$
  - “**Relative case**”  $(X, T_{X/S})$  where  $X \rightarrow S$
  - “**Integrable case**” when  $[V, V] \subset V$  (foliations)
- **Fonctor “1-jet”** :  $(X, V) \mapsto (\tilde{X}, \tilde{V})$  where :

$\tilde{X} = P(V) =$  bundle of projective spaces of lines in  $V$

$\pi : \tilde{X} = P(V) \rightarrow X, \quad (x, [v]) \mapsto x, \quad v \in V_x$

$\tilde{V}_{(x, [v])} = \{ \xi \in T_{\tilde{X}, (x, [v])} ; \pi_* \xi \in \mathbb{C}v \subset T_{X, x} \}$

- For every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  tangent to  $V$

$$f_{[1]}(t) := (f(t), [f'(t)]) \in P(V_{f(t)}) \subset \tilde{X}$$

$$f_{[1]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (\tilde{X}, \tilde{V}) \quad (\text{projectivized 1st-jet})$$



- For every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  tangent to  $V$

$$f_{[1]}(t) := (f(t), [f'(t)]) \in P(V_{f(t)}) \subset \tilde{X}$$

$$f_{[1]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (\tilde{X}, \tilde{V}) \quad (\text{projectivized 1st-jet})$$

- **Definition.** *Simple jet bundles :*

- $(X_k, V_k) = k$ -th iteration of functor  $(X, V) \mapsto (\tilde{X}, \tilde{V})$
- $f_{[k]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X_k, V_k)$  is the **projectivized  $k$ -jet of  $f$** .

- For every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  tangent to  $V$

$$f_{[1]}(t) := (f(t), [f'(t)]) \in P(V_{f(t)}) \subset \tilde{X}$$

$$f_{[1]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (\tilde{X}, \tilde{V}) \quad (\text{projectivized 1st-jet})$$

- **Definition.** *Simple jet bundles :*

–  $(X_k, V_k) = k$ -th iteration of functor  $(X, V) \mapsto (\tilde{X}, \tilde{V})$

–  $f_{[k]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X_k, V_k)$  is the **projectivized  $k$ -jet of  $f$** .

- **Basic exact sequences**

$$0 \rightarrow T_{\tilde{X}/X} \rightarrow \tilde{V} \xrightarrow{\pi^*} \mathcal{O}_{\tilde{X}}(-1) \rightarrow 0 \quad \Rightarrow \text{rk } \tilde{V} = r = \text{rk } V$$

$$0 \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow \pi^* V \otimes \mathcal{O}_{\tilde{X}}(1) \rightarrow T_{\tilde{X}/X} \rightarrow 0 \quad (\text{Euler})$$

- For every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  tangent to  $V$

$$f_{[1]}(t) := (f(t), [f'(t)]) \in P(V_{f(t)}) \subset \tilde{X}$$

$$f_{[1]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (\tilde{X}, \tilde{V}) \quad (\text{projectivized 1st-jet})$$

- **Definition.** *Simple jet bundles :*

–  $(X_k, V_k) = k$ -th iteration of functor  $(X, V) \mapsto (\tilde{X}, \tilde{V})$

–  $f_{[k]} : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X_k, V_k)$  is the **projectivized  $k$ -jet of  $f$** .

- **Basic exact sequences**

$$0 \rightarrow T_{\tilde{X}/X} \rightarrow \tilde{V} \xrightarrow{\pi^*} \mathcal{O}_{\tilde{X}}(-1) \rightarrow 0 \quad \Rightarrow \text{rk } \tilde{V} = r = \text{rk } V$$

$$0 \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow \pi^* V \otimes \mathcal{O}_{\tilde{X}}(1) \rightarrow T_{\tilde{X}/X} \rightarrow 0 \quad (\text{Euler})$$

$$0 \rightarrow T_{X_k/X_{k-1}} \rightarrow V_k \xrightarrow{(\pi_k)^*} \mathcal{O}_{X_k}(-1) \rightarrow 0 \quad \Rightarrow \text{rk } V_k = r$$

$$0 \rightarrow \mathcal{O}_{X_k} \rightarrow \pi_k^* V_{k-1} \otimes \mathcal{O}_{X_k}(1) \rightarrow T_{X_k/X_{k-1}} \rightarrow 0 \quad (\text{Euler})$$

- For  $n = \dim X$  and  $r = \operatorname{rk} V$ , get a **tower of  $\mathbb{P}^{r-1}$ -bundles**

$$\pi_{k,0} : X_k \xrightarrow{\pi_k} X_{k-1} \rightarrow \cdots \rightarrow X_1 \xrightarrow{\pi_1} X_0 = X$$

with  **$\dim X_k = n + k(r - 1)$ ,  $\operatorname{rk} V_k = r$ ,**

and **tautological line bundles  $\mathcal{O}_{X_k}(1)$  on  $X_k = P(V_{k-1})$ .**

- For  $n = \dim X$  and  $r = \operatorname{rk} V$ , get a **tower of  $\mathbb{P}^{r-1}$ -bundles**

$$\pi_{k,0} : X_k \xrightarrow{\pi_k} X_{k-1} \rightarrow \cdots \rightarrow X_1 \xrightarrow{\pi_1} X_0 = X$$

with  $\dim X_k = n + k(r - 1)$ ,  $\operatorname{rk} V_k = r$ ,

and **tautological line bundles  $\mathcal{O}_{X_k}(1)$**  on  $X_k = P(V_{k-1})$ .

- **Theorem.**  $X_k$  is a smooth compactification of

$$X_k^{\text{GG,reg}} / G_k = J_k^{\text{GG,reg}} / G_k$$

where  $G_k$  is the group of  $k$ -jets of germs of biholomorphisms of  $(\mathbb{C}, 0)$ , acting on the right by reparametrization:  $(f, \varphi) \mapsto f \circ \varphi$ , and  $J_k^{\text{reg}}$  is the space of  $k$ -jets of regular curves.

- For  $n = \dim X$  and  $r = \operatorname{rk} V$ , get a **tower of  $\mathbb{P}^{r-1}$ -bundles**

$$\pi_{k,0} : X_k \xrightarrow{\pi_k} X_{k-1} \rightarrow \cdots \rightarrow X_1 \xrightarrow{\pi_1} X_0 = X$$

with  **$\dim X_k = n + k(r - 1)$** ,  $\operatorname{rk} V_k = r$ ,

and **tautological line bundles  $\mathcal{O}_{X_k}(1)$**  on  $X_k = P(V_{k-1})$ .

- **Theorem.**  $X_k$  is a smooth compactification of

$$X_k^{\operatorname{GG}, \operatorname{reg}} / G_k = J_k^{\operatorname{GG}, \operatorname{reg}} / G_k$$

where  $G_k$  is the group of  $k$ -jets of germs of biholomorphisms of  $(\mathbb{C}, 0)$ , acting on the right by reparametrization:  $(f, \varphi) \mapsto f \circ \varphi$ , and  $J_k^{\operatorname{reg}}$  is the space of  $k$ -jets of regular curves.

- **Direct image formula.**  $(\pi_{k,0})_* \mathcal{O}_{X_k}(m) = E_{k,m} V^* =$   
invariant algebraic differential operators  $f \mapsto P(f_{[k]})$   
acting on germs of curves  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ .

- Although very interesting, results are currently limited by **lack of knowledge on jet bundles and differential operators**
- **Theorem (Bérczi-Kirwan, 2009).** *The ring of germs of invariant differential operators on  $(\mathbb{C}^n, T_{\mathbb{C}^n})$  at the origin*

$$\mathcal{A}_{k,n} = \bigoplus_m E_{k,m} T_{\mathbb{C}^n}^* \quad \text{is finitely generated.}$$

- Although very interesting, results are currently limited by **lack of knowledge on jet bundles and differential operators**
- **Theorem (Bérczi-Kirwan, 2009).** *The ring of germs of invariant differential operators on  $(\mathbb{C}^n, T_{\mathbb{C}^n})$  at the origin*

$$\mathcal{A}_{k,n} = \bigoplus_m E_{k,m} T_{\mathbb{C}^n}^* \quad \text{is finitely generated.}$$

- Checked by direct calculations  $\forall n, k \leq 2$  and  $n = 2, k \leq 4$ :

$$\mathcal{A}_{1,n} = \mathcal{O}[f'_1, \dots, f'_n]$$

$$\mathcal{A}_{2,n} = \mathcal{O}[f'_1, \dots, f'_n, W^{[ij]}], \quad W^{[ij]} = f'_i f''_j - f'_j f''_i$$

$$\mathcal{A}_{3,2} = \mathcal{O}[f'_1, f'_2, W_1, W_2][W]^2, \quad W_i = f'_i DW - 3f''_i W$$

$$\mathcal{A}_{4,2} = \mathcal{O}[f'_1, f'_2, W_{11}, W_{22}, S][W]^6, \quad W_{ii} = f'_i DW_i - 5f''_i W_i$$

where  $W = f'_1 f''_2 - f'_2 f''_1$ ,  $S = (W_1 DW_2 - W_2 DW_1)/W$ .



- **Generalized GGL conjecture.** *If  $(X, V)$  is directed manifold of general type, i.e.  $\det V^*$  big, then  $\exists Y \subsetneq X$  such that  $\forall f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  non const.,  $f(\mathbb{C}) \subset Y$ .*

- **Generalized GGL conjecture.** *If  $(X, V)$  is directed manifold of general type, i.e.  $\det V^*$  big, then  $\exists Y \subsetneq X$  such that  $\forall f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  non const.,  $f(\mathbb{C}) \subset Y$ .*
- **Remark.** Elementary by Ahlfors-Schwarz if  $r = \text{rk } V = 1$ .  $t \mapsto \log \|f'(t)\|_{V, h}$  is strictly subharmonic if  $r = 1$  and  $(V^*, h^*)$  has  $> 0$  curvature in the sense of currents.

- **Generalized GGL conjecture.** *If  $(X, V)$  is directed manifold of general type, i.e.  $\det V^*$  big, then  $\exists Y \subsetneq X$  such that  $\forall f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  non const.,  $f(\mathbb{C}) \subset Y$ .*
- **Remark.** Elementary by Ahlfors-Schwarz if  $r = \text{rk } V = 1$ .  $t \mapsto \log \|f'(t)\|_{V, h}$  is strictly subharmonic if  $r = 1$  and  $(V^*, h^*)$  has  $> 0$  curvature in the sense of currents.
- **Strategy.** Try some sort of induction on  $r = \text{rk } V$ . First try to get differential equations  $f_{[k]}(\mathbb{C}) \subset Z \subsetneq X_k$ . Take minimal such  $k$ . If  $k = 0$ , we are done! Otherwise  $k \geq 1$  and  $\pi_{k, k-1}(Z) = X_{k-1}$ , thus  $V' = V_k \cap T_Z$  has  $\text{rank} < \text{rk } V_k = r$  and should have again  $\det V'^*$  big (unless some improbable geometry situation occurs?).

- Generalized GGL conjecture.** *If  $(X, V)$  is directed manifold of general type, i.e.  $\det V^*$  big, then  $\exists Y \subsetneq X$  such that  $\forall f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$  non const.,  $f(\mathbb{C}) \subset Y$ .*
- Remark.** Elementary by Ahlfors-Schwarz if  $r = \text{rk } V = 1$ .  $t \mapsto \log \|f'(t)\|_{V,h}$  is strictly subharmonic if  $r = 1$  and  $(V^*, h^*)$  has  $> 0$  curvature in the sense of currents.
- Strategy.** Try some sort of induction on  $r = \text{rk } V$ . First try to get differential equations  $f_{[k]}(\mathbb{C}) \subset Z \subsetneq X_k$ . Take minimal such  $k$ . If  $k = 0$ , we are done! Otherwise  $k \geq 1$  and  $\pi_{k,k-1}(Z) = X_{k-1}$ , thus  $V' = V_k \cap T_Z$  has  $\text{rank} < \text{rk } V_k = r$  and should have again  $\det V'^*$  big (unless some unprobable geometry situation occurs?).
- Needed induction step.** *If  $(X, V)$  has  $\det V^*$  big and  $Z \subset X_k$  irreducible with  $\pi_{k,k-1}(Z) = X_{k-1}$ , then  $(Z, V')$ ,  $V' = V_k \cap T_Z$  has  $\mathcal{O}_{Z_\ell}(1)$  big on  $(Z_\ell, V'_\ell)$ ,  $\ell \gg 0$ .*

**Holomorphic Morse inequalities** (D-, 1985) Let  $L \rightarrow X$  be a holomorphic line bundle on a compact complex manifold  $X$ ,  $h$  a smooth hermitian metric on  $L$  and

$$\Theta_{L,h} = \frac{i}{2\pi} \nabla_{L,h}^2 = -\frac{i}{2\pi} \partial \bar{\partial} \log h$$

its curvature form. Then  $\forall q = 0, 1, \dots, n = \dim_{\mathbb{C}} X$

$$\sum_{j=0}^q (-1)^{q-j} h^j(X, L^{\otimes k}) \leq \frac{k^n}{n!} \int_{X(L,h,\leq q)} (-1)^q \Theta_{L,h}^n + o(k^n).$$

where

$$X(L, h, q) = \{x \in X; \Theta_{L,h}(x) \text{ has signature } (n - q, q)\}$$

( $q$ -index set), and

$$X(L, h, \leq q) = \bigcup_{0 \leq j \leq q} X(L, h, \leq j)$$

As a consequence, one gets an upper bound

$$h^0(X, L^{\otimes k}) \leq \frac{k^n}{n!} \int_{X(L,h,0)} \Theta_{L,h}^n + o(k^n)$$

and a lower bound

$$\begin{aligned} h^0(X, L^{\otimes k}) &\geq h^0(X, L^{\otimes k}) - h^1(X, L^{\otimes k}) \geq \\ &\geq \frac{k^n}{n!} \left( \int_{X(L,h,0)} \Theta_{L,h}^n - \int_{X(L,h,1)} |\Theta_{L,h}^n| \right) - o(k^n) \end{aligned}$$

As a consequence, one gets an upper bound

$$h^0(X, L^{\otimes k}) \leq \frac{k^n}{n!} \int_{X(L,h,0)} \Theta_{L,h}^n + o(k^n)$$

and a lower bound

$$\begin{aligned} h^0(X, L^{\otimes k}) &\geq h^0(X, L^{\otimes k}) - h^1(X, L^{\otimes k}) \geq \\ &\geq \frac{k^n}{n!} \left( \int_{X(L,h,0)} \Theta_{L,h}^n - \int_{X(L,h,1)} |\Theta_{L,h}^n| \right) - o(k^n) \end{aligned}$$

and similar bounds for the higher cohomology groups  $H^q$ :

$$h^q(X, L^{\otimes k}) \leq \frac{k^n}{n!} \int_{X(L,h,q)} |\Theta_{L,h}^n| + o(k^n)$$

$$h^q(X, L^{\otimes k}) \geq \frac{k^n}{n!} \left( \int_{X(L,h,q)} - \int_{X(L,h,q-1)} - \int_{X(L,h,q+1)} |\Theta_{L,h}^n| \right) - o(k^n)$$

Let  $J_k V$  be the bundle of  $k$ -jets of curves  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$



Let  $J_k V$  be the bundle of  $k$ -jets of curves  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$   
Assuming that  $V$  is equipped with a hermitian metric  $h$ , one defines a "weighted Finsler metric" on  $J^k V$  by taking  $p = k!$  and

$$\Psi_{h_k}(f) := \left( \sum_{1 \leq s \leq k} \varepsilon_s \|\nabla^s f(0)\|_{h(x)}^{2p/s} \right)^{1/p}, \quad 1 = \varepsilon_1 \gg \varepsilon_2 \gg \cdots \gg \varepsilon_k.$$

Let  $J_k V$  be the bundle of  $k$ -jets of curves  $f : (\mathbb{C}, T_{\mathbb{C}}) \rightarrow (X, V)$ . Assuming that  $V$  is equipped with a hermitian metric  $h$ , one defines a "weighted Finsler metric" on  $J^k V$  by taking  $p = k!$  and

$$\Psi_{h_k}(f) := \left( \sum_{1 \leq s \leq k} \varepsilon_s \|\nabla^s f(0)\|_{h(x)}^{2p/s} \right)^{1/p}, \quad 1 = \varepsilon_1 \gg \varepsilon_2 \gg \dots \gg \varepsilon_k.$$

Letting  $\xi_s = \nabla^s f(0)$ , this can actually be viewed as a metric  $h_k$  on  $L_k := \mathcal{O}_{X_k^{\text{GG}}}(1)$ , with curvature form  $(x, \xi_1, \dots, \xi_k) \mapsto$

$$\Theta_{L_k, h_k} = \omega_{\text{FS}, k}(\xi) + \frac{i}{2\pi} \sum_{1 \leq s \leq k} \frac{1}{s} \frac{|\xi_s|^{2p/s}}{\sum_t |\xi_t|^{2p/t}} \sum_{i, j, \alpha, \beta} c_{ij\alpha\beta} \frac{\xi_{s\alpha} \bar{\xi}_{s\beta}}{|\xi_s|^2} dz_i \wedge d\bar{z}_j$$

where  $(c_{ij\alpha\beta})$  are the coefficients of the curvature tensor  $\Theta_{V^*, h^*}$  and  $\omega_{\text{FS}, k}$  is the vertical Fubini-Study metric on the fibers of  $X_k^{\text{GG}} \rightarrow X$ . The expression gets simpler by using polar coordinates  $x_s = |\xi_s|^{2p/s}$ ,  $u_s = \xi_s / |\xi_s|_h = \nabla^s f(0) / \|\nabla^s f(0)\|$ .

In such polar coordinates, one gets the formula

$$\Theta_{L_k, h_k} = \omega_{\text{FS}, p, k}(\xi) + \frac{i}{2\pi} \sum_{1 \leq s \leq k} \frac{1}{s} x_s \sum_{i, j, \alpha, \beta} c_{ij\alpha\beta}(z) u_{s\alpha} \bar{u}_{s\beta} dz_i \wedge d\bar{z}_j$$

where  $\omega_{\text{FS}, k}(\xi)$  is positive definite in  $\xi$ . The other terms are a weighted average of the values of the curvature tensor  $\Theta_{V, h}$  on vectors  $u_s$  in the unit sphere bundle  $SV \subset V$ . The weighted projective space can be viewed as a circle quotient of the pseudosphere  $\sum |\xi_s|^{2p/s} = 1$ , so we can take here  $x_s \geq 0$ ,  $\sum x_s = 1$ . This is essentially a sum of the form  $\sum \frac{1}{s} \gamma(u_s)$  where  $u_s$  are random points of the sphere, and so as  $k \rightarrow +\infty$  this can be estimated by a “Monte-Carlo” integral

$$\left(1 + \frac{1}{2} + \dots + \frac{1}{k}\right) \int_{u \in SV} \gamma(u) du.$$

As  $\gamma$  is quadratic here,  $\int_{u \in SV} \gamma(u) du = \frac{1}{r} \text{Tr}(\gamma)$ .

It follows that the leading term in the estimate only involves the trace of  $\Theta_{V^*, h^*}$ , i.e. the curvature of  $(\det V^*, \det h^*)$ , which can be taken to be  $> 0$  if  $\det V^*$  is big.

**Corollary (D-, 2010)** Let  $(X, V)$  be a directed manifold,  $F \rightarrow X$  a  $\mathbb{Q}$ -line bundle,  $(V, h)$  and  $(F, h_F)$  hermitian. Define

$$L_k = \mathcal{O}_{X_k^{\text{GG}}}(1) \otimes \pi_k^* \mathcal{O}\left(\frac{1}{kr} \left(1 + \frac{1}{2} + \dots + \frac{1}{k}\right) F\right),$$

$$\eta = \Theta_{\det V^*, \det h^*} + \Theta_{F, h_F}.$$

Then for all  $q \geq 0$  and all  $m \gg k \gg 1$  such that  $m$  is sufficiently divisible, we have

$$h^q(X_k^{\text{GG}}, \mathcal{O}(L_k^{\otimes m})) \leq \frac{m^{n+kr-1} (\log k)^n}{(n+kr-1)! n! (k!)^r} \left( \int_{X(\eta, q)} (-1)^q \eta^n + \frac{C}{\log k} \right)$$

It follows that the leading term in the estimate only involves the trace of  $\Theta_{V^*, h^*}$ , i.e. the curvature of  $(\det V^*, \det h^*)$ , which can be taken to be  $> 0$  if  $\det V^*$  is big.

**Corollary (D-, 2010)** Let  $(X, V)$  be a directed manifold,  $F \rightarrow X$  a  $\mathbb{Q}$ -line bundle,  $(V, h)$  and  $(F, h_F)$  hermitian. Define

$$L_k = \mathcal{O}_{X_k^{\text{GG}}}(1) \otimes \pi_k^* \mathcal{O}\left(\frac{1}{kr} \left(1 + \frac{1}{2} + \dots + \frac{1}{k}\right) F\right),$$

$$\eta = \Theta_{\det V^*, \det h^*} + \Theta_{F, h_F}.$$

Then for all  $q \geq 0$  and all  $m \gg k \gg 1$  such that  $m$  is sufficiently divisible, we have

$$h^q(X_k^{\text{GG}}, \mathcal{O}(L_k^{\otimes m})) \leq \frac{m^{n+kr-1}}{(n+kr-1)!} \frac{(\log k)^n}{n! (k!)^r} \left( \int_{X(\eta, q)} (-1)^q \eta^n + \frac{C}{\log k} \right)$$

$$h^0(X_k^{\text{GG}}, \mathcal{O}(L_k^{\otimes m})) \geq \frac{m^{n+kr-1}}{(n+kr-1)!} \frac{(\log k)^n}{n! (k!)^r} \left( \int_{X(\eta, \leq 1)} \eta^n - \frac{C}{\log k} \right).$$

Using the above cohomological estimate, we obtain:

**Theorem (D-, 2010)** Let  $(X, V)$  be of general type, i.e.  $K_V = (\det V)^*$  is a big line bundle. Then there exists  $k \geq 1$  and an algebraic hypersurface  $Z \subsetneq X_k$  such that every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \mapsto (X, V)$  satisfies  $f_{[k]}(\mathbb{C}) \subset Z$  (in other words,  $f$  satisfies an algebraic differential equation of order  $k$ ).

Using the above cohomological estimate, we obtain:

**Theorem (D-, 2010)** Let  $(X, V)$  be of general type, i.e.  $K_V = (\det V)^*$  is a big line bundle. Then there exists  $k \geq 1$  and an algebraic hypersurface  $Z \subsetneq X_k$  such that every entire curve  $f : (\mathbb{C}, T_{\mathbb{C}}) \mapsto (X, V)$  satisfies  $f_{[k]}(\mathbb{C}) \subset Z$  (in other words,  $f$  satisfies an algebraic differential equation of order  $k$ ).

Another important consequence is:

**Theorem (D-, 2012)** A generic hypersurface  $X \subset \mathbb{P}^{n+1}$  of degree  $d \geq d_n$  with

$$d_2 = 286, \quad d_3 = 7316, \quad d_n = \left\lfloor \frac{n^4}{3} (n \log(n \log(24n)))^n \right\rfloor$$

(for  $n \geq 4$ ) satisfies the Green-Griffiths conjecture.

The proof of the last result uses an important idea due to Yum-Tong Siu, itself based on ideas of Claire Voisin and Herb Clemens, and then refined by M. Păun [Pau08], E. Rousseau [Rou06b] and J. Merker [Mer09].

The idea consists of studying vector fields on the **relative jet space of the universal family of hypersurfaces of  $\mathbb{P}^{n+1}$** .



The proof of the last result uses an important idea due to Yum-Tong Siu, itself based on ideas of Claire Voisin and Herb Clemens, and then refined by M. Păun [Pau08], E. Rousseau [Rou06b] and J. Merker [Mer09].

The idea consists of studying vector fields on the **relative jet space of the universal family of hypersurfaces of  $\mathbb{P}^{n+1}$** .

Let  $\mathcal{X} \subset \mathbb{P}^{n+1} \times \mathbb{P}^{N_d}$  be the universal hypersurface, i.e.

$$\mathcal{X} = \{(z, a); a = (a_\alpha) \text{ s.t. } P_a(z) = \sum a_\alpha z^\alpha = 0\},$$

let  $\Omega \subset \mathbb{P}^{N_d}$  be the open subset of  $a$ 's for which  $X_a = \{P_a(z) = 0\}$  is smooth, and let

$$p : \mathcal{X} \rightarrow \mathbb{P}^{n+1}, \quad \pi : \mathcal{X}|_\Omega \rightarrow \Omega \subset \mathbb{P}^{N_d}$$

be the natural projections.

Let

$$p_k : \mathcal{X}_k \rightarrow \mathcal{X} \rightarrow \mathbb{P}^{n+1}, \quad \pi_k : \mathcal{X}_k \rightarrow \Omega \subset \mathbb{P}^{N_d}$$

be the relative Green-Griffiths  $k$ -jet space of  $\mathcal{X} \rightarrow \Omega$ . Then J. Merker [Mer09] has shown that global sections  $\eta_j$  of

$$\mathcal{O}(T_{\mathcal{X}_k}) \otimes p_k^* \mathcal{O}_{\mathbb{P}^{n+1}}(k^2 + 2k) \otimes \pi_k^* \mathcal{O}_{\mathbb{P}^{N_d}}(1)$$

generate the bundle at all points of  $\mathcal{X}_k^{\text{reg}}$  for  $k = n = \dim X_a$ . From this, it follows that if  $P$  is a non zero global section over  $\Omega$  of  $E_{k,m}^{\text{GG}} T_{\mathcal{X}}^* \otimes p_k^* \mathcal{O}_{\mathbb{P}^{n+1}}(-s)$  for some  $s$ , then for a suitable collection of  $\eta = (\eta_1, \dots, \eta_m)$ , the  $m$ -th derivatives

$$D_{\eta_1} \dots D_{\eta_m} P$$

yield sections of  $H^0(\mathcal{X}, E_{k,m}^{\text{GG}} T_{\mathcal{X}}^* \otimes p_k^* \mathcal{O}_{\mathbb{P}^{n+1}}(m(k^2 + 2k) - s))$  whose joint base locus is contained in  $\mathcal{X}_k^{\text{sing}}$ , whence the result.

## References

**[Dem85]** Demailly, J.-P.: *Champs Magnétiques et Inégalités de Morse pour la  $d''$ -cohomologie*. Ann. Inst. Fourier (Grenoble) **35** (1985), no. 4, 189–229.

**[Dem95]** Demailly, J.-P.: *Algebraic Criteria for Kobayashi Hyperbolic Projective Varieties and Jet Differentials*. Algebraic geometry – Santa Cruz 1995, 285–360, Proc. Sympos. Pure Math., 62, Part 2, Amer. Math. Soc., Providence, RI, 1997.

**[Dem10]** Demailly, J.-P.: *Holomorphic Morse inequalities and the Green-Griffiths-Lang conjecture*, November 2010, arXiv: math.AG/1011.3636, dedicated to the memory of Eckart Viehweg; Pure and Applied Mathematics Quarterly **7** (2011) 1165–1208

**[Dem12]** Demailly, J.-P.: *Hyperbolic algebraic varieties and holomorphic differential equations*, lecture given at the VIASM Annual Meeting 2012, Hanoi 26 August 2012, 63p, to appear

**[D-EG00]** Demailly, J.-P., El Goul, J.: *Hyperbolicity of Generic Surfaces of High Degree in Projective 3-Space*. Amer. J. Math. **122** (2000), no. 3, 515–546.

**[Div09]** Diverio, S.: *Existence of global invariant jet differentials on projective hypersurfaces of high degree*. Math. Ann. **344** (2009) 293–315.

**[DMR09]** Diverio, S., Merker, J., Rousseau, E.: *Effective algebraic degeneracy*. e-print arXiv:0811.2346v5.

**[DT9]** Diverio, S., Trapani, T.: *A remark on the codimension of the Green-Griffiths locus of generic projective hypersurfaces of high degree*. e-print arXiv:0902.3741v2.

**[F-H91]** Fulton, W., Harris, J.: *Representation Theory: A First Course*. Graduate Texts in Mathematics, 129. Readings in Mathematics. Springer-Verlag, New York, 1991, xvi+551 pp.

**[G-G79]** Green, M., Griffiths, P.: *Two Applications of Algebraic Geometry to Entire Holomorphic Mappings*. The Chern Symposium 1979 (Proc. Internat. Sympos., Berkeley, Calif., 1979), pp. 41–74, Springer, New York-Berlin, 1980.

**[Kobayashi70]** Kobayashi S.: *Hyperbolic Manifolds and Holomorphic Mappings*. Marcel Dekker, Inc., New York 1970 ix+148 pp.

**[Lang86]** Lang S.: *Hyperbolic and Diophantine analysis*, Bull. Amer. Math. Soc. **14** (1986), no. 2, 159–205.

**[Mer08]** An algorithm to generate all polynomials in the  $k$ -jet of a holomorphic disc  $D \rightarrow \mathbb{C}^n$  that are invariant under source reparametrization, arxiv.org:math.CV/0808.3547.

**[Mer09]** Merker, J.: *Low pole order frames on vertical jets of the universal hypersurface*. Ann. Inst. Fourier (Grenoble) **59** (2009) 1077-1104.

**[Pau08]** Păun, M.: *Vector fields on the total space of*

*hypersurfaces in the projective space and hyperbolicity*. Math. Ann. **340** (2008), 875–892.

**[Rou05]** Rousseau, E: *Weak Analytic Hyperbolicity of Generic Hypersurfaces of High Degree in the Complex Projective Space of Dimension 4*. arXiv:math/0510285v1 [math.AG].

**[Rou06a]** Rousseau, E.: *Étude des Jets de Demailly-Semple en Dimension 3*. Ann. Inst. Fourier (Grenoble) **56** (2006), no. 2, 397–421.

**[Rou06b]** Rousseau, E: *Équations Différentielles sur les Hypersurfaces de  $\mathbb{P}^4$* . J. Math. Pures Appl. (9) **86** (2006), no. 4, 322–341.

**[Siu04]** Siu, Y.-T.: *Hyperbolicity in Complex Geometry*. The legacy of Niel Henrik Abel, 543–566, Springer, Berlin, 2004.

**[SY96]** Siu, Y.T., Yeung, S.K.: *Hyperbolicity of the complement of a generic smooth curve of high degree in the complex projective plane*. Invent. Math. **124** (1996), 573–618



**[SY97]** Siu, Y.T., Yeung, S.K.: *Defects for ample divisors of Abelian varieties, Schwarz lemma and hyperbolic surfaces of low degree*. Amer. J. Math. **119** (1997), 1139–1172

**[Tra95]** Trapani, S.: *Numerical criteria for the positivity of the difference of ample divisors*, Math. Z. **219** (1995), no. 3, 387–401.

**[Voj87]** Vojta, P.: *Diophantine Approximations and Value Distribution Theory*, Springer-Verlag, Lecture Notes in Mathematics no. 1239, 1987.