

A NOTE ON ASYMPTOTIC SERRE DUALITY FOR JET DIFFERENTIALS

MOHAMMAD REZA RAHMATI

ABSTRACT. We formulate and prove the existence of an asymptotic duality along the fibers of the Green-Griffiths jet bundles over projective manifolds. The existence of global sections for these bundles and also for their dual sheaves has been established by the author and J. P. Demailly in a previous article, [3].

1. INTRODUCTION

The main result of this paper is a Serre duality for asymptotic sections of jet bundles as formulated in Theorem (2.1) below. An application is also given for partial resolution of the Green-Griffiths conjecture.

P. Griffiths and M. Green [4] provided several equivalent methods for characterizing the bundle of jets over a projective variety X . Later, the techniques developed by Semple, Demailly, McQuillan, Siu and Green-Griffiths led to new conjectures in Kahler geometry. Demailly [1], [2] shows the existence of global sections of the Green-Griffiths bundles with sufficiently high degree.

Any entire curve $f : \mathbb{C} \rightarrow X$ satisfies $Q(f_{[k]}) = 0$, where $f_{[k]}$ a lift of f and Q a global section of the jet bundle (see [1]). As the section Q involves both the manifold and jet coordinates, it is desirable to show that there are enough dual global sections along the jet fibers to pair with the given section to give equations on the variety X .

The Green-Griffiths bundle associated with a pair (X, V) , where X is a projective variety and $V \subset T_X$ is a holomorphic sub-bundle of rank r , is a sequence of projective bundles over X which are defined inductively as in [[1], page 20]. Alternatively, one can define the Green-Griffiths bundle X_k as

$$(1) \quad X_k := (J_k V \setminus 0) / \mathbb{C}^*$$

where J_k is the bundle of germs of k -jets of Taylor expansion for f . The bundles $\pi_k : X_k \rightarrow X$ provide a tool to study entire holomorphic curves $f : \mathbb{C} \rightarrow X$, since such a curve has a lift to $f_{[k]} : \mathbb{C} \rightarrow X_k$ for every k . Using the notation in [1], we write $E_{k,m} V^* = (\pi_k)_* \mathcal{O}_{X_k}(m)$.

1991 *Mathematics Subject Classification.* 32Q45, 30D35, 14E99.

Key words and phrases. Green-Griffiths jet bundle, Serre duality, Entire Curve Locus, Holomorphic Morse Inequalities, Weighted Projective Space, Relative Canonical Sheaf.

In [1], [2], the existence of asymptotic global sections for Green-Griffiths jets on a projective variety X has been proved, by using Morse estimates for the curvature of suitable metrics on these bundles. It is shown in [1] that

$$(2) \quad H^0(X, E_{k,m}V^* \otimes A^{-1}) = H^0(X_k, \mathcal{O}_{X_k}(m) \otimes \pi_k^*A^{-1})$$

are non-trivial when $m \gg k \gg 0$, where A is a hermitian ample line bundle on X . In fact the factor A can be absorbed in the other over X_k if we assume $m \in \mathbb{Q}$.

The formulation of Serre duality is based on the existence of the canonical sheaf. A technical difficulty arises here that in the definition of the relative canonical sheaf, the fibers of $X_k \rightarrow X$ have singularities. However, there is a generalization of the definition of the canonical sheaf for singular varieties as explained in [1], [3].

The relative canonical sheaf $K_{X_k/X}$ can be defined in a manner similar to the definition of the canonical sheaf K_V , where V is a holomorphic subbundle of T_X . Serre duality can then be written as

$$(3) \quad H^0(X_k, (\pi_k)_*\mathcal{O}_{X_k}(m)) \otimes H^{k(r-1)}(X_k, K_{X_k/X} \otimes (\pi_k)_*\mathcal{O}_{X_k}(-m')) \longrightarrow \mathcal{O}_X$$

for $m, m' \gg k \gg 0$. The non-triviality of the cohomology groups has been proved by Demailly and the author, [1], [2] et [3].

2. SERRE DUALITY FOR JET BUNDLES

As explained above the existence of the relative canonical sheaves $K_{X_k/X}$ provides the possibility to discuss Serre duality along fibers of jet bundles. Because the fibers in the Green-Griffiths bundle are weighted projective spaces of appropriate weights $(1, 2, \dots, n)$ the relative Serre duality can be interpreted as the duality for coherent sheaves on weighted projective spaces. On account of this we review the formulation of the adjoint pair along the fibers of $E_{k,m}V^*$. Then, according to the classical Serre duality the dual pair associated to $H^0(F_x, (\pi_k)_*\mathcal{O}_{X_k,x}(m))$ is $H^{k(r-1)}(F_x, K_{F_x} \otimes (\pi_k)_*\mathcal{O}_{X_k,x}(-m))$, where K_{F_x} is the canonical sheaf of the fiber. In other words we have

$$(4) \quad H^0(\pi_k^{-1}(x), (\pi_k)_*\mathcal{O}_{X_k}(m))^\vee = H^{k(r-1)}(\pi_k^{-1}(x), K_{F_x} \otimes (\pi_k)_*\mathcal{O}_{X_k}(-m))$$

By the Leray spectral sequence for $\pi_k : X_k \rightarrow X$ we get the following

$$(5) \quad H^{k(r-1)}(X_k, K_{X_k/X} \otimes (\pi_k)_*\mathcal{O}_{X_k}(-m')) = H^0(X, R^{k(r-1)}(\pi_k)_*(K_{X_k/X} \otimes (\pi_k)_*\mathcal{O}_{X_k}(-m')))$$

On account of the formalism of Serre duality for coherent sheaves on projective spaces, the candidate for the duality is

$$(6) \quad H^0(X_k, (\pi_k)_* \mathcal{O}_{X_k}(m)) \otimes H^{k(r-1)}(X_k, K_{X_k/X} \otimes (\pi_k)_* \mathcal{O}_{X_k}(-m')) \longrightarrow \mathcal{O}_X$$

We have to make sure that the sheaf $H^{k(r-1)}(X_k, K_{X_k/X} \otimes (\pi_k)_* \mathcal{O}_{X_k}(-m'))$ is non-trivial i.e. has enough sections for $m \gg k \gg 0$.

Theorem 2.1. (*Serre Duality for Jet Fibers*) *There is a Serre duality for asymptotic cohomologies, $m, m' \gg k \gg 0$,*

$$(7) \quad H^0(X_k, (\pi_k)_* \mathcal{O}_{X_k}(m)) \otimes H^{k(r-1)}(X_k, K_{X_k/X} \otimes (\pi_k)_* \mathcal{O}_{X_k}(-m')) \longrightarrow \mathcal{O}_X$$

Proof. The duality in construction is the duality on each fiber of the jet bundle, all glued by the spectral sequence of Leray. In that sense, it is a duality of coherent sheaves on the weighted projective spaces [8]. The adjoint pair on the fiber along the fiber $\pi_k^{-1}(x)$ is given by the formula

$$(8) \quad H^0(\pi_k^{-1}(x), (\pi_k)_* \mathcal{O}_{X_k}(m))^\vee = H^{k(r-1)}(\pi_k^{-1}(x), K_{F_x} \otimes (\pi_k)_* \mathcal{O}_{X_k}(-m))$$

The degeneration of the Leray spectral sequence of the fibration $\pi_k : X_k \rightarrow X$ provides

$$(9) \quad H^0(X_k, (\pi_k)_* \mathcal{O}_{X_k}(m))^\vee = H^0(X, R^{k(r-1)}(\pi_k)_*(K_{X_k/X} \otimes (\mathcal{O}_{X_k}(-m))))$$

which is equivalent to (7). The non-triviality of the first factor in (7) is proved in [1]. The nontriviality of the adjoint cohomology group in the pairing is obtained from estimates

$$(10) \quad H^q(X_k, K_{X_k/X} \otimes (\pi_k)_* \mathcal{O}_{X_k}(-m')) \geq \sum_{q-1, q, q+1} \frac{rm^n}{r!} \int_{X(\Theta, j)} (-1)^{q-j} \Theta^n - o(m^n)$$

where Θ is the curvature of suitable k-jet metric on X_k and $X(\Theta, j) = \{x \in X; \Theta \text{ has signature } (n-j, j)\}$. We have considered the inequality for $q = k(r-1)$. It follows that when the $K_{X_k/X}$ is big, both of the factors in the pairing (7) are nontrivial for $m, m' \gg 0$, [cf. [3] section 4]. The theorem follows. \square

In [5], J. Merker shows that when X is a hypersurface of degree d in \mathbb{P}^{n+1} and is a generic member of the universal family $\mathfrak{X} \subset \mathbb{P}^{n+1} \times \mathbb{P}^{N_d}$, the Green-Griffiths conjecture holds for X . His method uses ideas of Y. T. Siu, on existence of slanting vector fields, see [9], [7]. The proof establishes the conjecture outside a certain algebraic subvariety $\Sigma \subset J_{\text{vert}}^n(\mathfrak{X})$ defined by Wronskians. An implication of Merker's result [5] is the following;

$$(11) \quad H^0(\mathfrak{X}_k, (\pi_k)_* \mathcal{O}_{\mathfrak{X}_k}(m)) \otimes H^{k(r-1)}(\mathfrak{X}_k, K_{\mathfrak{X}_k/\mathfrak{X}} \otimes \langle J_{\text{vert}}^k(\mathfrak{X}) \rangle^{m'}) \longrightarrow \mathcal{O}_{\mathfrak{X}}$$

where $m', m \gg k \gg 0$ and $\langle J_{\text{vert}}^k(\mathfrak{X}) \rangle^{m'}$ means the ring of operators generated by the $J_{\text{vert}}^k(\mathfrak{X})$ of degree m' .

Application to Green-Griffiths Conjecture: We note that any entire curve $f : \mathbb{C} \rightarrow X$ is satisfied by the sections $Q \in H^0(X_k, (\pi_k)_* \mathcal{O}_{X_k}(m))$ i.e $Q(f_{[k]}) = 0$ for $f_{[k]} : \mathbb{C} \rightarrow X_k$ a lift of f , cf. [1]. It follows that, in (7) after the composition for $Q \in H^0(X_k, (\pi_k)_* \mathcal{O}(X_k(m)))$ and $P \in H^0(X_k, K_{X_k/X} \otimes (\pi_k)_* \mathcal{O}(X_k(-m'))$, the resulting function also satisfies

$$(12) \quad \langle Q, P \rangle(f_{[k]}) = 0, \quad Q = \sum_{|\alpha|=m} A_\alpha(z) \xi^\alpha, \quad P = \sum_{|\beta|=m'} B_\beta(z) \partial_\xi^\beta$$

Therefore, the issue is to show that the image of (7) is a non-trivial ideal of \mathcal{O}_X . The above procedure plays a similar task as the idea on the existence of slanting vector fields along jet fibers, due to Siu [9]. In fact the slanting vector fields play the role of generating sections in the adjoint fiber rings of jet bundles.

ACKNOWLEDGEMENT: I thank the Abdus Salam School of Mathematical Sciences , GCU, for its research facilities and financial support.

REFERENCES

- [1] J. P. Demailly, Hyperbolic algebraic varieties and holomorphic differential equations, Acta Math. Vietnam, 37(4) 441-512, 2012
- [2] J. P. Demailly, Holomorphic Morse inequalities and Green-Griffiths-Lang conjecture, preprint
- [3] Jean-Pierre Demailly, Mohammad Reza Rahmati, Morse cohomology estimates for jet differential operators, Bollettino dell'Unione Matematica Italiana, <https://doi.org/10.1007/s40574-018-0172-2>
- [4] M. Green, P. Griffiths, Two applications of algebraic geometry to entire holomorphic mappings, The Chern Symposium 1979. (Proc. Intern. Sympos., Berkeley, California, 1979) 41-74, Springer, New York, 1980.
- [5] J. Merker, low pole order frameson vertical jets of the universal hypersurface, Ann. Institut Fourier (Grenoble), 59, 2009, 861-933
- [6] J. Merker, Application of computational invariant theory to Kobayashi hyperbolicity and to GreenGriffiths algebraic degeneracy, Journal of Symbolic Computation 45 (2010) 9861074
- [7] M. Paun, Vector fields on the total space of hypersurfaces in the projective space and hyperbolicity, Math. Ann. (2008) 340:875-892
- [8] M. Rossi, L. Terracini, Weighted projective spaces from toric point of view and computational aspects, preprint
- [9] Y. T. Siu, Hyperbolicity in complex geometry, The legacy of Niels Henrik Abel, Springer, Berlin, 2004, 543-566

ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES , GCU, LAHORE, PAKISTAN

,

E-mail address: rahmati@sms.edu.pk