Edge currents in the absence of edges

P. Exner[a,b], A. Joye[c], and H. Kovařík[a,d],

a) Department of Theoretical Physics, Nuclear Physics Institute, Academy of Sciences, 25068 Řež near Prague
b) Doppler Institute, Czech Technical University, Břehová 7, 11519 Prague, Czech Republic
c) Institut Fourier, Université de Grenoble 1, 38402 Saint-Martin d'Heres, France
d) Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, 18000 Prague
exner@ujf.cas.cz, joye@ujf-grenoble.fr, kovarik@ujf.cas.cz

Abstract

We investigate a charged two-dimensional particle in a homogeneous magnetic field interacting with a periodic array of point obstacles. We show that while Landau levels remain to be infinitely degenerate eigenvalues, between them the system has bands of absolutely continuous spectrum and exhibits thus a transport along the array. We also compute the band functions and the corresponding probability current.

The fact that the presence of boundaries can induce a transport in a system with a homogeneous magnetic field is known for long [1, 2] and has numerous consequences in solid state physics. The mentioned pioneering papers were followed by tremendous number of studies in which the magnetic transport was analyzed numerically in various models as well as experimentally. The obstacles at which the particle "bounces" need not at that be hard walls but also objects with openings such as various antidot lattices; for a sample of literature see, e.g., Refs. [3, 4], the following papers by the same authors, and references therein. The closest to the subject of this letter is a very recent

paper [5] where the array of obstacles is strictly one-dimensional and there are no boundaries to help the transport.

One can say therefore that at the heuristic level most aspects of the two-dimensional magnetic transport are understood. With this fact and the extensive literature in mind it is a bit surprising how little attention was paid during a decade and a half to a strict derivation of transport properties from the first principles – which is after all the *raison d'être* of theoretical physics. The situation concerning this aspect of the problem changed recently where several rigorous studies appeared. It was shown, e.g., that the edge currents in a halfplane survive a mild disorder so that away of the Landau levels the spectrum remains absolutely continuous [6, 7], and that the result extends to planar domains containing an open wedge [8].

Recall that the wall producing the edge states need not be of a potential type. It is known, e.g., that a step of the magnetic field or another variation exhibiting a translational symmetry will smear again the Landau levels into a continuous spectrum [9, 10]. Similarly to the usual edge states, this type of propagation allows for a classical explanation in terms of the cyclotronic radius changing with the magnetic field – see Ref. [11], Sec. 6.5.

The aim of the present letter is to contribute to this series of rigorous studies with a simple solvable model in which a charged quantum particle in the plane exposed to a homogeneous magnetic field of intensity B perpendicular to the plane interacts with a periodic array of point obstacles situated at the x axis and described by δ potentials. We shall show that the model exhibits a magnetic transport of which we can with a license say that it is a purely quantum effect in the sense that a quantum particle propagates despite the fact that its classical counterpart moves on localized circular trajectories – apart of a zero-measure family of the initial conditions. The delta potentials are certainly an idealization; a more realistic model with potential-type obstacles will be discussed in a forthcoming paper.

Using the Landau gauge, we can write the Hamiltonian formally as

$$H_{\alpha,\ell} = (-i\partial_x + By)^2 - \partial_y^2 + \sum_j \tilde{\alpha}\delta(x - x_0 - j\ell)\,, \tag{1}$$

where $\ell > 0$ is the array spacing. Since we are interested mainly in the essence of the effect, we use everywhere rationalized units $\hbar = c = e = 2m = 1$. The interaction term, in particular the formal coupling constant $\tilde{\alpha}$, needs an explanation, since the two-dimensional δ potential is an involved object. We

adopt here the conventional definition from Ref. [12] which determines the latter by means of the boundary conditions

$$L_1(\psi,\vec{a}_j)+2\pi\alpha L_0(\psi,\vec{a}_j)=0\,,\quad j=0,\pm1,\pm2,\dots \tag{2}$$

with $\vec{a}_j:=(x_0+j\ell,0)$, where L_k are the generalized boundary values

$$L_0(\psi,\vec{a}):=\lim_{|\vec{x}-\vec{a}|\to0}\frac{\psi(\vec{x})}{\ln|\vec{x}-\vec{a}|}\,,\ L_1(\psi,\vec{a}):=\lim_{|\vec{x}-\vec{a}|\to0}\Big[\psi(\vec{x})-L_0(\psi,\vec{a})\,\ln|\vec{x}-\vec{a}|\Big]\,, \tag{3}$$

and α is the (rescaled) coupling constant; the free (Landau) Hamiltonian corresponds to $\alpha=\infty$. Recall that since the magnetic field amounts locally to a regular potential in the s-wave subspace, the non-magnetic boundary conditions of Ref. [12] need not be modified – see, e.g., Ref. [13].

The difference between α and $\tilde{\alpha}$ reflects the nontrivial way in which the two-dimensional point interaction arises in the limit of scaled potentials. Due to the coupling constant renormalization, a caution is required when interpreting spectral properties of such a Hamiltonian. On the other hand, the two-dimensional point interaction yields a *generically* correct description of low-energy scattering which can be tested in experiments – see references in [12] or a fresh example in [14]. To understand the above remark about the purely quantum nature of the transport here, recall the well-known expression for the scattering length which shows that that obstacles have a "point" character if $e^{2\pi\alpha}\ll\ell$, i.e., if the point interaction is strong enough. Below we shall show that the transport exists here for *any* finite value of α.

Using the periodicity, we can perform the Bloch decomposition in the x direction writing

$$H_{\alpha,\ell}=\frac{\ell}{2\pi}\int_{|\theta\ell|\le\pi}^{\oplus}H_{\alpha,\ell}(\theta)\,d\theta\,, \tag{4}$$

where the fiber operator $H_{\alpha,\ell}(\theta)$ is of the form (1) on the strip $0\le x\le\ell$ with the boundary conditions

$$\partial_x^i\psi(\ell-,y)=e^{i\theta\ell}\partial_x^i\psi(0+,y)\,,\quad i=0,1\,. \tag{5}$$

The Green's function of the operator $H_{\alpha,\ell}(\theta)$ is given by means of the Krein formula [12, App.A],

$$\begin{aligned}(H_{\alpha,\ell}(\theta)-z)^{-1}(\vec{x},\vec{x}')&=G_0(\vec{x},\vec{x}';\theta,z)\\&+(\alpha-\xi(\vec{a}_0;\theta,z))^{-1}G_0(\vec{x},\vec{a}_0;\theta,z)G_0(\vec{a}_0,\vec{x}';\theta,z)\,,\end{aligned} \tag{6}$$

where G_0 is the free Green's function and

$$\xi(\vec{a};\theta,z) := \lim_{|\vec{x}-\vec{a}|\to 0} \left(G_0(\vec{a},\vec{x};\theta,z) - \frac{1}{2\pi} \ln|\vec{x}-\vec{a}| \right) \tag{7}$$

is its regularized value at the point $\vec{a}$. The Bloch conditions (5) determine eigenvalues and eigenfunctions of the transverse part of the free operator,

$$\mu_m(\theta) = \left(\frac{2\pi m}{\ell} + \theta \right)^2, \ \eta_m^\theta(x) = \frac{1}{\sqrt{\ell}} \, \mathrm{e}^{i(2\pi m + \theta\ell)x/\ell} \,, \tag{8}$$

where m runs through integers. Then we have

$$G_0(\vec{x},\vec{x}';\theta,z) = -\sum_{m=-\infty}^{\infty} \frac{u_m^\theta(y_<) v_m^\theta(y_>)}{W(u_m^\theta, v_m^\theta)} \, \eta_m^\theta(x) \overline{\eta_m^\theta}(x') \,, \tag{9}$$

where $y_<, y_>$ is the smaller and larger value, respectively, of y, y', and u_m^θ, v_m^θ are solutions to the equation

$$-u''(y) + \left(By + \frac{2\pi m}{\ell} + \theta \right)^2 u(y) = z u(y) \tag{10}$$

such that u_m^θ is L^2 at $-\infty$ and v_m^θ is L^2 at $+\infty$; in the denominator we have their Wronskian. By the argument shift we get

$$u_m^\theta(y) = u\left(y + \frac{2\pi m + \theta\ell}{B\ell} \right) \tag{11}$$

and a similar relation for v_m^θ, where u, v are the corresponding oscillator solutions. Of course, we have $W(u_m^\theta, v_m^\theta) = W(u,v)$. The functions u, v express in terms of the confluent hypergeometric functions [15, Chap. 13]:

$$v(y) = e^{-By^2/2} U\left(\frac{B-z}{4B}, \frac{1}{2}; By^2 \right) \tag{12}$$

away from zero, and u is obtained by analytical continuation in the y^2 variable; together we have

$$\left\{ \begin{array}{c} u \\ v \end{array} \right\}(y) = \sqrt{\pi} \, e^{-By^2/2} \left[\frac{M\left(\frac{B-z}{4B}, \frac{1}{2}; By^2\right)}{\Gamma\left(\frac{3B-z}{4B}\right)} \pm 2\sqrt{B} y \, \frac{M\left(\frac{3B-z}{4B}, \frac{3}{2}; By^2\right)}{\Gamma\left(\frac{B-z}{4B}\right)} \right] . \tag{13}$$

From here and Ref. [15, Chap. 6] we compute the Wronskian; in combination with (8) we get

$$G_0(\vec{x},\vec{x}';\theta,z) = -\frac{2^{(z/2B)-(3/2)}}{\sqrt{\pi B}\ell}\,\Gamma\left(\frac{B-z}{2B}\right)\,\mathrm{e}^{i\theta(x-x')} \times \sum_{m=-\infty}^{\infty} u\left(y_< + \frac{2\pi m+\theta\ell}{B\ell}\right) v\left(y_> + \frac{2\pi m+\theta\ell}{B\ell}\right)\,\mathrm{e}^{2\pi i m(x-x')/\ell}\,. \quad (14)$$

As expected the function has singularities which are independent of θ and coincide with the Landau levels, i.e., $z_n = B(2n+1),\ n = 0,1,2,\ldots$. Let us observe first that each z_n remains to be infinitely degenerate eigenvalue of the "full" fiber operator $H_{\alpha,\ell}(\theta)$. To this end, one has to adapt the argument of Refs. [16, 17] to the set of functions $w^k \sin\left(\frac{\pi w}{\ell}\right) e^{-B|w|^2/4}$, $k = 0,1,\ldots$, with $w := x+iy$ which vanish at the points of the array so the conditions (2) are satisfied for them automatically.

On the other hand, $H_{\alpha,\ell}(\theta)$ has also eigenvalues away of z_n which we denote as $\epsilon_n(\theta) \equiv \epsilon_n^{(\alpha,\ell)}(\theta)$. In view of (6) they are given by the implicit equation

$$\alpha = \xi(\vec{a}_0;\theta,\epsilon) \quad (15)$$

and the corresponding eigenfunctions are

$$\psi_n^{(\alpha,\ell)}(\vec{x};\theta) = G_0(\vec{x},\vec{a}_0;\theta,\epsilon_n(\theta))\,. \quad (16)$$

In order to evaluate them, we have to assess the convergence of the series in (14). Using the asymptotic behavior

$$\left\{\begin{array}{c} u \\ v \end{array}\right\}(y) = e^{\mp\{\pm\}By^2/2}\left(\mp\sqrt{B}y\right)^{\frac{z-B}{2B}}\left(1+\mathcal{O}(|y|^{-2})\right) \quad (17)$$

for $y \to \mp\infty$, we find that the product

$$s_m := u\left(y_< + \frac{2\pi m+\theta\ell}{B\ell}\right) v\left(y_> + \frac{2\pi m+\theta\ell}{B\ell}\right)$$

is for $y \neq y'$ governed by the exponential term,

$$s_m \sim \exp\left\{\frac{B}{2}\left(y_<^2 - y_>^2\right) + \left(\theta - \frac{2\pi|m|}{\ell}\right)(y_> - y_<)\right\}\left(|m|^{-1} + \mathcal{O}(|m|^{-2})\right) \quad (18)$$

as $|m| \to \infty$, while for $y = y'$ we have

$$s_m = -\frac{1}{4\pi}\,|m|^{-1} + \mathcal{O}(|m|^{-2})\,,$$

so the series (14) is not absolutely convergent. Summing now the contributions from $\pm m$ we see that in the limit $x' \to x$ it diverges at the same rate as the Taylor series of $-(1/2\pi)\ln\zeta$ does for $\zeta \to 0+$. Hence we get

$$\xi(\vec{x};\theta,z) = \sum_{m=-\infty}^{\infty} \left\{ \frac{1-\delta_{m,0}}{4\pi|m|} - \frac{2^{-2\zeta-1}}{\sqrt{\pi B}\ell}\,\Gamma(2\zeta)\,(uv)\left(y + \frac{2\pi m + \theta\ell}{B\ell}\right) \right\}\,, \quad (19)$$

where $\zeta := \frac{B-z}{4B}$. The expression is independent of x, because the regularized resolvent does not change if the array is shifted in the x direction. We can write it by means of the first hypergeometric function alone, since

$$(uv)(\xi/\sqrt{B}) = \pi\,\mathrm{e}^{-\xi^2}\left[\frac{M(\zeta,\frac{1}{2};\xi^2)^2}{\Gamma(\zeta+\frac{1}{2})^2} - 4\xi^2\frac{M(\zeta+\frac{1}{2},\frac{3}{2};\xi^2)^2}{\Gamma(\zeta)^2}\right]\,, \quad (20)$$

where $\xi := \sqrt{B}\,\big(y + \frac{2\pi m+\theta\ell}{B\ell}\big)$.

Spectral bands of our model are given by the ranges of the functions $\epsilon_n(\cdot)$. Solutions of the condition (15) do not cross the Landau levels, because $\xi(\vec{a}_0;\theta,\cdot)$ is increasing in the intervals $(-\infty,B)$ and $(B(2n-1),B(2n+1))$ and diverges at the endpoints; this is a general feature [18]. The spectrum will be continuous away of z_n if the latter are nowhere constant. In view of the spectral condition (15) one has to check that $\xi(\vec{x};\theta,z)$ is nowhere constant as a function of θ. Notice that each term in (19) is real-analytic for real z and the series has a convergent majorant independent of θ; hence $\xi(\vec{x};\cdot,z)$ is real-analytic as well and one has to check that it is non-constant in the whole Brillouin zone $[-\pi/\ell,\pi/\ell)$.

Suppose that the opposite is true. Then the Fourier coefficients

$$c_k := \int_{-\pi/\ell}^{\pi/\ell} \xi(\vec{x};\theta,z)\,e^{ik\ell\theta}\,d\theta \quad (21)$$

should vanish for any non-zero integer k. Since the summand in (19) behaves as $\mathcal{O}(|m|^{-2})$ as $|m| \to \infty$, we may interchange the summation and integration. A simple change of variables then gives

$$c_k = -\frac{2^{-2\zeta-1}}{\sqrt{\pi B}\ell}\,\Gamma(2\zeta)\lim_{M\to\infty}\int_{-\pi(2M+1)}^{\pi(2M+1)} (uv)\left(y+\frac{\vartheta}{B\ell}\right)\,e^{ik\vartheta}\,d\vartheta\,, \quad (22)$$

so

$$\hat{F}_y(k) := \int_{-\infty}^{\infty} F_y(\vartheta)\, e^{ik\vartheta}\, d\vartheta = 0\,, \tag{23}$$

where $F_y(\vartheta) := (uv)\left(y + \frac{\vartheta}{B\ell}\right)$. The same reasoning applies to any finitely periodic extension of $\xi(\vec{x}; \theta, z)$, hence (23) is valid for each non-zero rational k. However, the function decays $\mathcal{O}(|\vartheta|^{-1})$ and the integral makes sense only as the principal value. We shall use the above mentioned asymptotic behavior which implies, in particular,

$$F_y(\vartheta) = -\frac{1}{4\pi\sqrt{1+\vartheta^2}} + f_y(\vartheta)\,, \tag{24}$$

where $f_y(\vartheta) = \mathcal{O}(|\vartheta|^{-2})$ uniformly in $y \in [0, \ell]$. Thus

$$\hat{F}_y(k) = -\frac{1}{2\pi}\, K_0(k) + \hat{f}_y(k)\,, \tag{25}$$

see [19, 3.754.2]. Since $f_y \in L^1$, the second term at the r.h.s. is continuous w.r.t. k and the same is then true for $\hat{F}_y$; this means that the relation (23) is valid for any nonzero k. Furthermore, $\hat{f}_y$ is bounded and K_0 diverges logarithmically at $k = 0$, hence $\int_{-N}^{N} F_y(\vartheta)\, e^{ik\vartheta}\, d\vartheta$ can be bounded by an integrable function independent of N. Then

$$\int_{-\infty}^{\infty} \hat{F}_y(k)\phi(k)\, dk = \int_{-\infty}^{\infty} dk\, \phi(k) \lim_{N\to\infty} \int_{-N}^{N} F_y(\vartheta)\, e^{ik\vartheta}\, d\vartheta = \int_{-\infty}^{\infty} F_y(\vartheta)\, \hat{\phi}(\vartheta)\, d\vartheta \tag{26}$$

holds for any $\phi \in \mathcal{S}(\mathbb{R})$, i.e., $\hat{F}_y(k)$ is the Fourier transform of $F_y(\vartheta)$ in the sense of tempered distributions. Since this is a one-to-one correspondence [20, Thm.IX.2], we arrive at the absurd conclusion that $F_y = 0$. We get thus the following result:

Theorem. For any real α the spectrum of $H_{\alpha,\ell}$ consists of the Landau levels $B(2n+1)$, $n = 0, 1, 2, \ldots$, and absolutely continuous spectral bands situated between adjacent Landau levels and below B.

Let us remark that during the final stage of the work we learned about a similar result for a chain of point scatterers in a three-dimensional space with a homogeneous magnetic field [21]. Due to the higher dimensionality, the spectrum is purely a.c. in that case and has at most finitely many gaps.

The above theorem says a little about the character of the transport. To get a better idea we solve the spectral condition (15) numerically for several



Figure 1: The eigenvalues of $H_{\alpha,l}(\theta)$ in the units of B. (Top) $n = 20$, $B = 4$ (full line), $B = 6$ (full dotted line) and $B = 8$ (dotted line). (Bottom) $n = 1$, $B = 10$ (full line) and $B = 15$ (dotted line). The thick lines represent the Landau levels.

values of the parameters. The results are plotted in Fig. 1 for the second and 21st spectral band. We see that the bands move downwards with decreasing α and their profile becomes more complicated with the band index n; a higher B tends to smear the structure.

The Bloch functions (16) are in general complex-valued and yield thus a nontrivial probability current, $\vec{\jmath}_n(\vec{x};\theta) = 2\,\mathrm{Im}\left(\bar{\psi}_n^{(\alpha,\ell)}(\vec{\nabla} - i\vec{A})\psi_n^{(\alpha,\ell)}\right)(\vec{x};\theta)$. The current pattern changes with θ oscillating between a symmetric "two-way" picture and the situations where one direction clearly prevails, these extremal behaviors occurring at the extrema of the corresponding band func-



Figure 2: Probability current for $n = 20$, $B = 4$, $\alpha = 0.5$ and two different values of θ corresponding to the extremal points of $\epsilon_{20}(\cdot)$: $\theta = 0$ (top), $\theta = 2.2$ (bottom), see the arrows in Fig. 1. The star marks the point perturbation position.

tion. This is illustrated in Fig. 2. In addition, while the pattern has predominantly "laminar" character, in some parts current vortices may form, mainly in low spectral bands as it is illustrated in Fig. 3. Such effects are known from numerical analysis of related models mentioned in the introduction.

To sum up the above discussion, we have analyzed the behavior of a quantum particle in the plane exposed to a homogeneous magnetic field and interacting with a periodic array of point perturbations. We have shown that while the Landau levels survive, the spectrum develops an absolutely continuous part, i.e. a sequence of spectral bands. Depending on the quasi-momentum, the particle is transported along the array with zero or nonzero mean longitudinal momentum, and the probability current pattern may exhibit vortices in some regions.



Figure 3: Probability current for $n = 1$, $B = 20$, $\alpha = -1$ and $\theta = 0$. In the bottom graph the inset shows a vortex between the point perturbations.

Acknowledgments

A.J. wishes to thank his hosts at the Nuclear Physics Institute in Řež, where this work was initiated. The research has been partially supported by the GAAS grant 1048801. We are grateful to V. Geyler for making Ref. 21 available to us prior to publication.

References

[1] B.I. Halperin, *Phys. Rev.* **B25**, 2185 (1982).

[2] A.H. MacDonald, P. Středa, *Phys. Rev.* **B29**, 1616 (1984).

[3] R. Fleischmann, T. Geisel, R. Ketzmerick, *Phys. Rev. Lett.* **68**, 1367 (1992).

[4] I.V. Zozoulenko, F.A. Maaø, E.H. Hauge, *Phys.Rev.* **B53**, 7975 (1996).

[5] T. Ueta, *Phys. Rev.* **B60**, 8213 (1999).

[6] S. De Bièvre, J.V. Pulé, *Math. Phys. El. J.* **5** (1999), No. 3.

[7] J. Fröhlich, G.M. Graf, J. Walcher: On the extended nature of edge states of quantum Hall Hamiltonians, *math-ph/9903014*.

[8] J. Fröhlich, G.M. Graf, J. Walcher: Extended quantum Hall edge states: general domains, *mp_arc 99-327*.

[9] A. Iwatsuka, *Publ. RIMS* **21**, 385 (1985).

[10] M. Mantoiu, R. Purice, *Commun. Math. Phys.* **188**, 691 (1997).

[11] H.L. Cycon, R.G. Froese, W. Kirsch, B. Simon: *Schrödinger Operators with Applications to Quantum Mechanics and Global Geometry,* Springer, Berlin 1987.

[12] S.A. Albeverio, F. Gesztesy, R. Høegh-Krohn, H. Holden: *Solvable Models in Quantum Mechanics,* Springer, Berlin 1988.

[13] F. Gesztesy et al., in *Schrödinger Operators, Standard and Non-Standard*, World Scientific 1989; p. 147.

[14] P. Exner, P. Šeba, Phys. Lett. **A228**, 146 (1997).

[15] M.S. Abramowitz, I.A. Stegun, eds.: *Handbook of Mathematical Functions*, Dover, New York 1965.

[16] Y. Avishai, R.M. Redheffer, Y.M. Band, *J. Phys.* **A25**, 3883 (1992).

[17] T.C. Dorlas, N. Macris, J.V. Pulé: Characterization of the spectrum of the Landau Hamiltonian with point impurities, *math-ph/9904035*.

[18] M.G. Krein, H.K. Langer, *Funct. Anal. Appl.* **5**, 59 (1971).

[19] I.S. Gradshtein, I.M. Ryzhik: *Tables of Integral, Series, and Products*, Academic Press, New York 1980.

[20] M. Reed, B. Simon: *Methods of Modern Mathematical Physics II*, Academic, New York 1975.

[21] S. Albeverio, V.A. Geyler, O.G. Kostov, *Rep. Math. Phys.* **44** (1999), to appear.

Figure captions

Figure 1. The eigenvalues of $H_{\alpha,l}(\theta)$ in the units of B. (Top) $n = 20$, $B = 4$ (full line), $B = 6$ (full dotted line) and $B = 8$ (dotted line). (Bottom) $n = 1$, $B = 10$ (full line) and $B = 15$ (dotted line). The thick lines represent the Landau levels.

Figure 2. Probability current for $n = 20$, $B = 4$, $\alpha = 0.5$ and two different values of θ corresponding to the extremal points of $\epsilon_{20}(\cdot)$: $\theta = 0$ (top), $\theta = 2.2$ (bottom), see the arrows in Fig. 1. The star marks the point perturbation position.

Figure 3. Probability current for $n = 1$, $B = 20$, $\alpha = -1$ and $\theta = 0$. In the bottom graph the inset shows a vortex between the point perturbations.